Wetlands (2019) 39:865–877
https://doi.org/10.1007/s13157-018-1120-8

WETLANDS RESTORATION

Impacts of Shallow Lake Restoration on Vegetation and Breeding Birds in Iowa

Rachel A. Vanausdall[1] · Stephen J. Dinsmore[1]

Received: 6 June 2018 / Accepted: 21 November 2018 / Published online: 14 December 2018
© Society of Wetland Scientists 2018

Abstract

The Shallow Lakes Restoration Project aims to restore eutrophic shallow lakes throughout the Iowa Prairie Pothole Region (PPR). We compared the vegetation taxa richness and frequency of taxa in three vegetation groups surveyed in 2016 and 2017 across non-restored sites, younger restorations, and older restorations. We also assessed the impact of these groups on numbers of breeding marsh birds. Restored wetlands had between 2.7 (95% CI 2.3, 3.2) and 4.7 (95% CI 3.9, 5.5) more species than non-restored sites among the three vegetation groups. *Typha* sp. was the most abundant emergent species. *Lemna minor* and *Ceratophyllum demersum* were the most abundant floating-leaved and submersed species, respectively. The frequency of submersed aquatic vegetation increased with years since restoration, while floating-leaved vegetation and *Typha* sp. peaked at 7 years post-restoration. The frequency of *Typha* sp. positively influenced Marsh Wrens and Yellow-headed Blackbirds. Water depth negatively affected Marsh Wrens, but water depth positively influenced Yellow-headed Blackbirds. Floating-leaved vegetation positively affected Virginia Rails, while water depth had a negative effect on this species. Our results indicate that after about 7 years further management of the vegetation and water levels may need to occur and should include maintaining appropriate conditions for breeding marsh birds.

Keywords Marsh bird · Restoration · Wetland vegetation

Introduction

Wetlands in the Prairie Pothole Region (PPR) have experienced dramatic declines and alterations since European settlement (Dahl 2014). This area was once characterized by expansive wetlands interspersed with grasslands and hosted diverse wildlife and floral populations (Bishop 1981; Tiner Jr 1984; Van Meter and Basu 2015). Due to agricultural expansion more than half of the wetland area has been lost in the conterminous United States alone (Dahl 2014). Many of the wetlands that remain have been altered due to watershed changes, increased fish abundance, increased sedimentation, and excess nutrient loading and chemical drift (Martin and Hartman 1986; Neely and Baker 1989; Baker 1992; Euliss Jr and Mushet 1996; Gleason et al. 2003). Wildlife species, including birds, rely on emergent and submersed aquatic plants for nesting and foraging, and the severe decline in quality wetland habitat has likely led to a decline in many species (Banks and Springer 1994; Page and Gill Jr 1994; Igl and Johnson 1997; Lor and Malecki 2002; Anteau and Afton 2008). In response to these changes, wetland restoration that involves manipulating water levels have become important tools for expanding and improving the quality of habitat for a variety of wildlife species, including birds (Rundle and Fredrickson 1981; Elphick and Oring 1998; Taft et al. 2002).

Historically, wetlands in the PPR experienced varying amounts of vegetation and open water over time, which resulted in seasonal or annual changes in the avian community (Weller and Spatcher 1965; Murkin et al. 1997; Hershey et al. 1999; Murkin and Ross 2000). During drought periods, seeds from vegetation germinate and more nutrients become available as plant litter decomposes (Harris and Marshall 1963; van der Valk and Davis 1976; Bärlocher et al. 1978). The basin gradually refills and submersed aquatic, floating-leaved, and

✉ Rachel A. Vanausdall
rvanausd@iastate.edu

Stephen J. Dinsmore
cootjr@iastate.edu

[1] Department of Natural Resource Ecology and Management, Iowa State University, 2310 Pammel Dr., 339 Science Hall II, Ames, IA 50011, USA

emergent vegetation gradually increases (van der Valk and Davis 1978). Emergent vegetation will eventually die back due to prolonged inundation and muskrat activity, and this will lead to an open lake period once again (van der Valk and Davis 1978). During the vegetated periods, the presence of emergent and aquatic vegetation provide favorable conditions for breeding marsh birds. For example, the growth of tall emergent vegetation, such as *Typha* sp. and *Scheonoplectus* sp., can be used as a substrate for nest building by marsh passerines, particularly Marsh Wrens and Yellow-headed Blackbirds (Kroodsma and Verner 2013; Lupien et al. 2014). Similarly, secretive marsh birds will use *Typha* sp. for nesting, foraging, and displaying (Lor and Malecki 2006; Harms and Dinsmore 2013; Glisson et al. 2015). Submersed aquatic vegetation can be used as a substrate for aquatic invertebrates (Voigts 1976; Driver 1977; Wrubleski 1999; Murkin and Ross 2000), which are important parts of the diets of several marsh passerines and secretive marsh birds (Conway 1995; Twedt and Crawford 1995; Kroodsma and Verner 2013). Floating-leaved vegetation, such as duckweeds, may also provide habitat for invertebrates (Harper and Bolen 1996).

However, agriculture and development have altered water level fluctuations and water quality in remaining wetlands, particularly shallow lakes, which are semi-permanent or permanent wetlands <1 m deep (Cowardin et al. 1979; Miller et al. 2012; Geisthardt et al. 2013). A combination of wetland consolidation and an increase in the abundance of planktivorous and benthivorous fishes contribute to more stable water conditions and increased sedimentation (Peterka 1989; Hanson and Butler 1994; Hanson and Riggs 1995; Euliss Jr and Mushet 1996; Zimmer et al. 2000; Pothoff et al. 2008; Stewart and Downing 2008; McLean et al. 2016). Such factors can lead to increased turbidity and concentrations of nitrogen, phosphorus, and phytoplankton (Hanson and Riggs 1995; Zimmer et al. 2002). Subsequently, submersed aquatic vegetation declines and further allows for increased phytoplankton growth, and the persistent deep water causes emergent vegetation to decline (van der Valk and Davis 1978; Timms and Moss 1984; Scheffer et al. 1993; Sayer et al. 2010). The loss in vegetation and increased turbidity do not provide appropriate habitat for invertebrates (Scheffer et al. 1993; Zimmer et al. 2000) and reduces appropriate foraging and nesting conditions for many breeding birds (Glisson et al. 2015).

In an effort to remedy the degradation of shallow lakes, several restoration projects have focused on implementing the natural wet-dry cycle as a management tool (Chow-Fraser 2005; Geisthardt et al. 2013). Many of these projects focus on maintaining conditions that favor abundant vegetation, particularly emergent vegetation and submersed aquatic plants, and may result in a shift to a clear water state (Scheffer et al. 1993; Chow-Fraser 2005; Søndergaard et al. 2007). Compared to unmanaged wetlands, restored wetlands tend to improve habitat conditions for breeding birds (VanRees-Siewert and Dinsmore 1996; Connor and Gabor 2006; Kaminski et al. 2006). However, several factors may prevent a project from persisting in this desired state for long periods of time (e.g., >10 years), and habitat suitability for marsh birds may actually decline (Glisson et al. 2015). Continuous management, such as water level manipulations and invasive species control, may be necessary (Søndergaard et al. 2007; Hanson et al. 2017). For example, submersed aquatic plants are known to help stabilize the clear water state of shallow lakes (Carpenter and Lodge 1986; Scheffer et al. 1993, Jeppesen et al. 1997), but they may not immediately respond to restoration efforts (Strand 1999; Søndergaard et al. 2007; Bortolotti et al. 2016). Additionally, external nutrient inputs may further prevent the proliferation of submersed aquatic plants (Lauridsen et al. 2003a; Zimmer et al. 2003), so additional management tools may be needed to increase vegetation growth. Monitoring the vegetation community and its impacts on wetland-dependent wildlife may help elucidate when such actions are warranted.

Since 2006 the Iowa Department of Natural Resources and Ducks Unlimited, Inc. have restored over 38 shallow lakes in Iowa PPR through the Shallow Lakes Restoration Project (SLRP). The goal of the SLRP is to improve water quality and the vegetation community to increase the establishment of diverse fish, bird, and invertebrate communities (Geisthardt et al. 2013). Since its implementation, these shallow lakes have shown improvements in water quality and vegetation structure (Geisthardt et al. 2013). However, as they age some shallow lakes appear to be showing a decline in water and vegetation cover (Geisthardt et al. 2013), indicating a need for further management. Our goals were to record the vegetation community of these shallow lakes, determine how they change over time by collecting information from shallow lakes of various states (i.e., years since restoration), and examine how the vegetation affects numbers of wetland-dependent birds. We expected restored shallow lakes to have a greater species richness and frequency of emergent and aquatic vegetation. At the same time, we expected some vegetation (e.g., *Typha* sp., submersed aquatic vegetation) to decline with years since restoration. We also hypothesized that several taxa would influence the abundance of secretive marsh birds. Because they provide nesting substrates and habitat for prey items, we expected emergent vegetation, floating-leaved vegetation, and submersed aquatic vegetation to have positive effects on marsh birds (Verner 1965; Verner and Engelsen 1970; Lor and Malecki 2006; Harms and Dinsmore 2013). Additionally, we hypothesized that the general height and density of vegetation and water depth may influence the counts of marsh birds (Verner 1965; Sayre and Rundle 1984; Twedt and Crawford 1995; Tozer et al. 2010; Lupien et al. 2014).

Methods

Study Area

Our study shallow lakes were located within the Des Moines Lobe region in Iowa, an area formed by the retreating Wisconsin glacial advance 14,000 years ago (IAN 2001; Miller et al. 2009). The PPR covers about 700,000 km^2 in the United States and Canada, and the DML makes up only 800 km^2 of that area (Stewart and Kantrud 1971; Bishop 1981; Kantrud et al. 1989, IAN 2001, Dahl 2014). The PPR is characterized by palustrine and lacustrine wetlands (Kantrud et al. 1989).

In this study, the term "restored" refers to shallow lakes that were once severely degraded and subsequently restored by manipulating the hydrology to improve water quality and vegetation. These shallow lakes were passively restored (i.e., no seed additions), and they were drained using an existing outlet structure to begin the restoration process. Infrastructure, such as water control structures, water channels, pipelines, and fish exclusion structures, were installed into nearly all shallow lakes to manage water levels and exclude rough fish. However, some sites still contained rough fish during the survey periods. Once the restoration process began, sites were refilled gradually over (ideally) a 2-year period to allow vegetation to reestablish. Likewise, the term "non-restored" refers to shallow lakes that were unmanipulated. Most of these shallow lakes were void of emergent vegetation and contained turbid water; some may be restored within the next few years. We considered the date of restoration to be the start of the drawdown, even if it was before completion of the water control structure. The years since restoration ranged between 1 and 12 years, but most restored shallow lakes were restored >2 years prior to this study.

Site Selection

To examine how shallow lakes in different restoration states differ in vegetation composition and influence breeding bird use, 19 restored sites were chosen based on years since restoration, spanning the period from one to eleven years post-restoration. Because only about 38 shallow lakes had been restored at the start of this study, we had a limited number of restorations to choose from. We wanted to survey at least two shallow lakes at each "age". We randomly chose shallow lakes in each age class, but we could not do this for every age group, as there was often only one. Based on recommendation from the Iowa DNR, we also chose 11 non-restored shallow lakes to examine pre-restoration bird use of shallow lakes. One of these shallow lakes was in the early stages of restoration in 2017 and, therefore, considered to be a restored site in that year. Study sites were located in 12 Iowa counties: Buena Vista, Calhoun, Cerro Gordo, Clay, Dickinson, Emmet, Hancock, Palo Alto, Pocahontas, Winnebago, Worth, and Wright. All shallow lakes were surveyed in the summer of 2016 and 2017.

We grouped restored shallow lakes into two categories based on years since restoration for some of the analyses. Several studies have examined the changes in the vegetation community of restored sites at different ages (Galatowitsch and van der Valk 1996a, b; Aronson and Galatowitsch 2008), and others have found that sites considered to be "older" may differ from "younger" sites (Badiou et al. 2011; Bortolotti et al. 2016). For example, previously degraded wetlands restored 1–3 years prior to the study by Bortolotti et al. (2016) had less submersed aquatic vegetation (SAV) indicative of remnant (i.e., undisturbed) wetlands compared to wetlands restored 7–10 years prior. Additionally, Søndergaard et al. (2007) determined that 5–10 years after biomanipulation some shallow lakes returned to a eutrophic, turbid state with little to no vegetation. Based on these findings we examined our sites using the following grouping method (hereafter referred to as restoration state): non-restored, younger (1–5 years since restoration), and older (6–11 years since restoration). This grouping method provided a more balanced design, and we were particularly interested in whether these restorations were showing signs of returning to a eutrophic state >5 years post-restoration.

Vegetation Sampling

We completed vegetation surveys once during each of the two summers in 2016 and 2017. We used a method similar to Webb et al. (2010) and created north-south and east-west transects in each shallow lake. These were situated along the maximum width of each shallow lake for each direction. We used a 1 × 1 m quadrat every 50 m for transects ≤800 m and every 100 m for transects >800 m (Webb et al. 2010). These transects were generated using ArcMap 10.3 (Environmental Systems Research Institute, Inc., Redlands, CA). At each quadrat we recorded the percent cover of plants to genus (e.g., *Typha* sp.; Monfils et al. 2014) or species and their structural definition (wet prairie, sedge meadow, shallow emergent, deep emergent, submersed aquatic, floating-leaved, mudflat annual, and woody shrub; Galatowitsch and van der Valk 1994). For all groups other than submersed aquatics and floating-leaved plants percent cover was calculated using basal cover. We estimated the areal percent cover of submersed aquatics and floating-leaved plants. We used a rake to facilitate the identification of submersed aquatic plants. We assigned each quadrat into one of three size classes based on maximum vegetation height (1 = 0–0.5 m, 2 = 0.5–1 m, 3 = >1 m; Harms and Dinsmore 2013). We used a similar method for vegetation density and assigned each quadrat into one of three density classes (1 = completely open, 2 = anything that falls in between the two extremes, 3 = water not visible

through the base of stems at water level; Conway 2009). Water depth was recorded to the nearest cm at the center of each quadrat.

Bird Surveys

We conducted unlimited-radius, 10-min point counts throughout each site for breeding marsh birds (Ralph et al. 1995). Points were situated randomly in shallow lakes, and the number of points depended on the size of the site. Two points were placed in sites 10.1–20 ha, three points in sites 20.1–30 ha, four points in sites 20.1–40 ha, and five points in sites >40 ha (Harms 2011). Points were situated >400 m apart to avoid double-counting individual birds (Conway 2011). We surveyed each set of points twice during each year to account for any seasonal variation in the vocalization rates of species (Gibbs and Melvin 1993, Conway et al. 2004). Point counts were conducted between a half-hour before sunrise to up to four hours after sunrise. We did not survey in rainy conditions or when winds exceeded 20 km/h (Conway 2009).

To improve detection of secretive marsh birds (i.e., rails, bitterns, grebes), we incorporated call-broadcast surveys into our point count surveys according to methods described by the North American Marsh Bird Monitoring Protocol (Conway 2009, 2011). The first five minutes of the survey period were silent, while the last five minutes were recorded calls. Each minute corresponded to one of five species of secretive marsh birds; the first 30 s included a recording, followed by 30 s of silence. The sequence of calls we used, from first to last, was Least Bittern (*Ixobrychus exilis*), Sora (*Porzana carolina*), Virginia Rail (*Rallus limicola*), King Rail (*Rallus elegans*), and Pied-billed Grebe (*Podilymbus podiceps*; Conway 2011). Except for the King Rail, these are regular breeders in Iowa. In previous call-broadcast surveys conducted in Iowa (Harms 2011), the King Rail did not have many detections, but Sora and Virginia Rail tended to respond to King Rail calls as readily as they responded to intraspecific calls (T. Harms, pers. comm.). We used an MP3 player (SanDisk Sansa Clip 1GB, SanDisk Corporation, Milpitas, California) attached to portable speakers (JBL Flip 3, Harman International Industries, Inc., Stamford, Connecticut) and broadcast at 90 dB from a distance of 1 m in front of the speakers (Conway 2011). The speaker faced the interior of the wetland and was 0.5 m from the ground or water surface (Harms 2011). Because we conducted surveys twice per year, we averaged the number of birds counted between the two visits for each year.

Vegetation Community Analysis

We examined the community structure of three major functional groups of plants (Galatowitsch and van der Valk 1994): emergent vegetation, submersed aquatics, and floating-leaved vegetation. We mostly encountered deep emergent and shallow emergent species and the other functional groups alone were not encountered enough for statistical analyses. We were interested in how several measurements of these groups differed across shallow lakes in different restoration states. Since our study unit was the shallow lake, we averaged the following metrics across the three restoration states: total taxa richness and frequency of occurrence for each taxa. We decided to average most of our calculations across sites ($n = 60$) because these data tended to be zero-inflated at the quadrat level ($n = 1472$). We examined total taxa richness within each of three plant functional groups. We used taxa richness because we included both species and plants identified to genus. Frequency of occurrence is the percentage of points each taxa was encountered at a shallow lake. Because these metrics were non-normal and tended to have unequal variances across the restoration states, we used a Kruskal-Wallis rank sum test to examine differences in taxa richness and frequency of occurrence among restoration states (Kruskal and Wallis 1952). We did not find an effect of year for all metrics, so we combined data across both years. For significant ($P < 0.05$) differences we used a Games-Howell multiple comparisons test to determine which restoration states were different (Games and Howell 1976).

We were also interested in how frequency and percent cover changed with years since restoration and water depth, respectively. For a subset of these taxa, we used two different methods to examine the response of plant taxa to years since restoration and water depth. We assessed years since restoration and water depth separately because we were interested in the separate effects of each covariate on changes in taxa frequency or water depth. Furthermore, frequency of occurrence was a site-level metric, while water depth was measured at the quadrat level. For years since restoration, we used a linear mixed effects models (*lme4* package in R; Bates et al. 2015), with frequency as the response variable and a linear or quadratic effect of years since restoration as the explanatory variable. We also included year as a fixed effect and site as a random effect to account for the repeated measures of frequency at each shallow lake (Zuur et al. 2009). We used a likelihood ratio test to determine the most competitive model (linear vs. quadratic). We had an adequate number of encounters for the following 10 taxa to assess years since restoration: *Bulboschoenus fluviatilis, Phalaris arundinacea, Phragmites australis*, *Schoenoplectus sp.*, *Typha sp.*, total floating-leaved vegetation, *Ceratophyllum demersum*, *Potamogeton sp.*, *Stuckenia pectinata,* and *Urticularia vulgaris*, and total submersed aquatic vegetation. We decided to examine only floating-leaved vegetation because these species showed similar patterns in occurrence among restoration states, and several species had no occurrences at non-restored sites. Because these were frequencies, we transformed them using the

arcsine-square root function (Zar 2010). For these same groups we examined Pearson's correlation coefficients between percent cover of each plant taxa and water depth.

Vegetation Effects on Marsh Bird Abundance

We were interested in the relationship between several vegetation variables and counts of breeding birds. Specifically, we made several hypotheses concerning the directional effects of vegetation variables and water depth on two groups and three species of birds (Table 1). Marsh passerines included four species of obligate wetland breeding passerines (Brown and Smith 1998): Marsh Wren (*Cistothorus palustris*), Swamp Sparrow (*Melospiza georgiana*), Red-winged Blackbird (*Agelaius phoenicus*), and Yellow-headed Blackbird (*Xanthocephalus xanthocephalus*). Secretive marsh birds included American Bittern (*Botaurus lentiginosus*), Least Bittern, Virginia Rail, and Common Gallinule (*Gallinula galeata*). We assessed these groups because they require emergent vegetation for nesting and foraging (Lor and Malecki 2006; Tozer et al. 2010; Kroodsma and Verner 2013). There is also evidence that they are sensitive to certain habitat conditions, such as water quality, and so may be considered wetland quality indicators (Eddleman et al. 1988; Conway 1995; Cumbee Jr et al. 2008; Lowther et al. 2009; Glisson et al. 2015). Additionally, we examined the effect of vegetation and water depth on Marsh Wren, Yellow-headed Blackbird, and Virginia Rail separately; these are relatively common species in Iowa wetlands but are experiencing population declines elsewhere (Conway et al. 1994; Conway 1995; Kroodsma and Verner 2013).

We used linear mixed models to assess the effect of these habitat variables on breeding birds (*lme4* package in R; Bates et al. 2015). We always included year as a fixed effect and site as a random effect to account for any inter-year variation and the repeated measures conducted on wetlands (Zuur et al. 2009). We generated a global model for each group and species, and avoided including variables with a VIF > 5 in the same global model (Montgomery and Peck 1992). We square root transformed any variables that did not meet assumptions of normality and homogeneity (Zar 2010). We used backward elimination of the fixed effects to determine the final model (*lmerTest* package in R; Kuznetsova et al. 2017).

Results

We surveyed a total of 742 quadrats between 5 July and 21 August in 2016 and 730 quadrats between 27 June and 12 August in 2016 at 30 shallow lakes. The number of quadrats in each restoration state was relatively similar, but we had fewer quadrats in older restorations (Table 2).

Species richness differed among restoration states for emergent ($\chi^2_2 = 33.53$, $P < 0.01$), floating-leaved ($\chi^2_2 = 65.63$, $P < 0.01$), and submersed aquatic vegetation ($\chi^2_2 = 26.24$, $P < 0.01$; Fig. 1a). For emergent species, both older ($t_{47} = 4.10$, $P < 0.01$) and younger ($t_{69} = 6.90$, $P < 0.01$) restorations had more species than non-restored sites. There were also more floating-leaved species at older ($t_{45} = 10.60$, $P < 0.01$) and younger ($t_{65} = 9.50$, $P < 0.01$) restorations than at non-restored shallow lakes. Similarly, there were more species of submersed aquatic vegetation at older ($t_{72} = 4.27$, $P < 0.01$) and younger ($t_{82} = 4.71$, $P < 0.01$) restorations.

Frequency of occurrence for species differed for some emergent and floating-leaved vegetation across restoration states (Table 3; Fig. 1). Among emergent species, *Typha* sp. was encountered more frequently than any other species for both younger and older restorations ($\chi^2_2 = 33.18$, $P < 0.01$; Fig. 1b). *Typha* sp. had a greater frequency at older and younger restorations than non-restored shallow lakes. For other emergent species, frequency tended to be greater at restored sites than non-restored sites, but this pattern was not significant. Of the floating-leaved species, *Lemna minor* ($\chi^2_2 = 29.95$, $P < 0.01$), *Lemna trisulca* ($\chi^2_2 = 13.47$, $P < 0.01$), *Spirodela polyrhiza* ($\chi^2_2 = 17.51$, $P < 0.01$), and *Wolffia* sp. ($\chi^2_2 = 20.05$, $P < 0.01$) differed among the restoration states (Fig. 1c). Both *Lemna minor* and *Spirodela polyrhiza* had greater frequencies in older and younger restorations than non-restored sites. *Wolffia* sp. and *Lemna trisulca* had greater frequencies in older restorations than non-restored sites.

Table 1 Predicted effects of habitat covariates on breeding marsh birds surveyed in the PPR of Iowa, summers 2016 and 2017

Habitat variable	Marsh passerines	Secretive marsh birds	Marsh Wren	Yellow-headed Blackbird	Virginia Rail
Schoenoplectus sp.	0	0	+	0	0
Typha sp.	+	+	+	+	+
Submersed aquatic vegetation	+	+	+	+	+
Floating-leaved vegetation	0	+	0	0	+
Medium vegetation (0.5–1 m)	0	+	+	–	+
Highly dense vegetation	+	+	+	+	+
Water Depth (cm)	+	–	+	+	–

We used "+" to represent a positive effect and "-" to represent a negative effect. The variables with "0" were not included in the global models

Table 2 Number of shallow lakes, number of surveyed quadrats, and mean values (with SD) for years since restoration, area, water depth, marsh passerines, Marsh Wrens, Yellow-headed Blackbird, secretive marsh birds, and Virginia Rails for 30 shallow lakes in the Iowa PPR, summers 2016 and 2017

	Non-restored	Younger restorations	Older restorations
No. visits	21	17	22
No. surveyed quadrats	504	569	399
Years since restoration	–	4.14 (0.89)	8.12 (1.64)
Area (ha)	116.87 (107.62)	165.53 (173.65)	138.24 (116.48)
Water depth (cm)	115.64 (55.58)	86.96 (42.70)	83.38 (49.95)
Marsh passerines	12.26 (7.18)	27.55 (10.24)	36.21 (12.94)
Marsh Wrens	3.21 (2.60)	9.20 (6.48)	10.91 (4.36)
Yellow-headed Blackbirds	1.76 (2.06)	7.55 (8.86)	12.15 (5.04)
Secretive marsh birds	0.57 (0.80)	2.23 (2.06)	4.06 (3.28)
Virginia Rails	0.40 (0.58)	1.18 (2.09)	2.09 (1.62)

Although non-restored sites and younger restorations tended to have greater frequencies of submersed aquatic vegetation, this relationship was not significant (Fig. 1d).

Years since restoration influenced the frequency of three species and two functional groups, and water depth showed a significant correlation with percent cover of some taxa. *Typha* sp., floating-leaved vegetation, and *Urticularia vulgaris* increased and then decreased with years since restoration (7-year peak) (Fig. 2b). Submersed aquatic vegetation and *Ceratophyllum demersum* showed a positive linear relationship with years since restoration (Fig. 2a). Three emergent species showed a significant negative relationship with water depth, while all the floating-leaved and submersed aquatic vegetation groups showed a strong positive relationship with water depth (Table 4).

Five habitat variables influenced counts of breeding marsh birds (Table 5). The final model for marsh passerines included a positive effect of *Typha* sp., and a positive effect of submersed aquatic vegetation. *Typha* sp. positively influenced numbers of Marsh Wrens, while *Typha* sp., submersed aquatic vegetation, and water depth all positively influenced numbers of Yellow-headed Blackbirds. The secretive marsh bird final model included a positive effect of moderately dense vegetation. Floating-leaved vegetation positively influenced and water depth negatively influenced counts of Virginia Rails.

Discussion

Restored shallow lakes in the Iowa PPR are showing changes in the vegetation community after restoration, and we found evidence that some of these vegetation changes may influence breeding marsh birds. Within a few years post-restoration, a variety of emergent, floating-leaved, and submersed aquatic vegetation created a more species rich-community, and these vegetation groups showed positive effects on both marsh passerines and secretive marsh birds. However, the frequency of different plant taxa was highly variable among years and restoration states. This is likely due to the timing of the drawdown and the reflooding periods of restoration, when nutrients become available and facilitate vegetation growth (Kadlec et al. 2000; van der Valk et al. 2000). At the same time, we found that parts of the vegetation community are beginning to decline with time since restoration, which may have important implications for management decisions concerning marsh birds. These declines may be a result of prolonged inundation (van der Valk and Davis 1978), increases in planktivorous fishes (Timms and Moss 1984, Jeppesen et al. 1997, Zimmer et al. 2001), sedimentation (Jurik et al. 1994; Euliss Jr and Mushet 1996), or increased nutrient loading from the surrounding landscape (Neely and Baker 1989; Lauridsen et al. 2003b).

Table 3 Summary of Games-Howell multiple comparisons tests for vegetation taxa that differed in frequency of occurrence among three restoration states of shallow lakes surveyed in the Iowa PPR, summers 2016 and 2017

Species/Group	Older:Non-restored				Younger:Non-restored				Younger:Older			
	Estimate	t	df	P	Estimate	t	df	P	Estimate	t	df	P
Typha sp.	0.46	6.51	17	< 0.01	0.38	6.03	23	< 0.01	−0.08	0.88	35	0.66
Lemna minor	0.42	5.19	16	< 0.01	0.38	6.26	22	< 0.01	−0.04	0.43	31	0.90
Lemna trisulca	0.08	2.98	16	0.02	0.09	2.25	21	0.09	0.003	0.07	36	1.00
Spirodela polyrhiza	0.25	3.20	16	0.01	0.16	3.3	22	0.01	−0.09	1.00	28	0.59
Wolffia sp	0.21	3.20	16	0.01	0.09	2.3	21	0.08	−0.12	1.60	27	0.27



Fig. 1 **a** Mean taxa richness of three vegetation types and mean frequency of **b** emergent, **c** floating-leaved, and **d** submersed aquatic vegetation types for shallow lakes surveyed in the Iowa PPR, 2016 and 2017

Overall, restorations are showing increases in taxa richness and frequency of occurrence of vegetation as they progress, but these factors may be declining in older restorations. In particular, we found that younger restorations tended to have more species in the emergent group. This influx of productivity within the first few years of restoration results from exposing the wetland soil, encouraging the germination of seeds already in the seed bank (van der Valk and Davis 1976; Wienhold and van der Valk 1989). Seeds may also be dispersed from nearby wetlands or ditches (Kettenring and Galatowitsch 2011) or from waterbirds (Figuerola et al. 2002). However, the seeds of some species vary in their ability to remain viable after several years of inundation prior to restoration (Wienhold and Galatowitsch 1988) and others may not be so easily dispersed (Kettenring and Galatowitsch 2011). Invasive perennials, such as *Phalaris arundinacea* and *Typha* x *glauca,* can also dominate initial restoration efforts, outcompeting and preventing the establishment of native plants (Aronson and Galatowitsch 2008; Mitchell et al. 2011; Lishawa et al. 2013). This could explain the lower species richness of older restorations; *Typha* sp. was the most frequently encountered emergent species and had the greatest overall frequency in both older and younger restorations. At the same time, *Typha* sp. showed a quadratic relationship with years since restoration and peaked in frequency at around 7–8 years after restoration. This may be due to muskrat activity (pers. obs.) and prolonged inundation (van der Valk and Davis 1978). Indeed, we found that *Typha* sp. showed a negative correlation with water depth.

Floating-leaved and submersed aquatic species also appear to be increasing in frequency, and these species are particularly important in stabilizing the clear water state of restored shallow lakes and providing habitat and forage for wildlife (Carpenter and Lodge 1986; Jeppesen et al. 1997; Lumsden et al. 2015). Total frequency of floating-leaved vegetation appeared to peak within 5 to 6 years after restoration. Species such as *Lemna minor*, *Spirodela polyrhiza*, and *Wolffia* sp. tended to have their greatest frequencies in older restorations, but these were not significantly greater than younger restorations. This variation and decline could be due to grazing from



Fig. 2 **a** Predicted frequency of *Ceratophyllum demersum* and submersed aquatic vegetation and **b** predicted frequency of *Typha* sp., *Urticularia vulgaris*, and floating-leaved vegetation at shallow lakes surveyed in the Iowa PPR, 2016 and 2017

waterfowl (Lauridsen et al. 1993), water level fluctuations (Chow-Fraser 2005), changes in light conditions (Bini et al. 1999), temperature (Minc 1997; Smith and Moelyowati 1998), or changes in nutrient inputs (Lougheed et al. 2001). Floating-leaved vegetation also showed a relationship with years since restoration that was similar to emergent cover. Along with several emergent species, the abundance of floating-leaved species may be driven by high nutrient content (e.g., phosphorous), so perhaps the similarities between the two functional groups are due to changes in the water chemistry (Bini et al. 1999).

Table 4 Pearson's correlation coefficients calculated between percent cover of plant taxa and water depth measured at shallow lakes in the Iowa PPR, summers 2016 and 2017

Taxa	Water depth (cm)
Bulboschoenus fluviatilis	0.02
Phalaris arundinacea	−0.16*
Phragmites australis	−0.11*
Schoenoplectus sp	0.003
Typha sp	−0.10*
Ceratophyllum demersum	0.36*
Potamogeton sp	0.16*
Stuckenia pectinata	0.21*
Urticularia vulgaris	0.17*
Total submersed aquatic vegetation	0.49*
Total floating-leaved vegetation	0.30*

The asterisks (*) indicate a significant ($P < 0.05$) correlation value

In contrast, submersed aquatic species may not have reached their peak in these restorations. There is evidence that submersed aquatic vegetation growth can be delayed in shallow lake restorations (Lauridsen et al. 1994; Søndergaard et al. 2007; Bortolotti et al. 2016) and species composition changes over time (Hansel-Welch et al. 2002) and with sediment type (Lauridsen et al. 1994). However, some species are less tolerant of changes in water condition (e.g., underwater light availability), and the most frequently encountered submersed aquatic vegetation in this study appeared to be species that can tolerate a range of conditions. *Ceratophyllum demersum* and *Stuckenia pectinata* were the most abundant submersed aquatics. These species tend to be more tolerant of slightly turbid waters, partly because each species is structurally adapted to survive in low-light conditions (Bini et al. 1999; Lougheed et al. 2001). *Ceratophyllum demersum* forms dense canopies beneath the water surface, while *Stuckenia pectinata* has leaves that float on or just below the water's surface (Lougheed et al. 2001). They were also present in several non-restored sites. On the other hand, *Urticularia vulgaris, Najas* sp., and some species of *Potamogeton* (e.g., *Potamogeon foliosus*) have smaller leaves that do not reach

Table 5 Coefficient estimates (with SE) for vegetation variables by average counts of breeding marsh passerines, secretive marsh birds, and three species of common, obligate, wetland breeders surveyed in the PPR of Iowa, summers 2016 and 2017

Habitat variable	Typha sp	Submersed aquatic vegetation	Floating-leaved vegetation	Moderately dense vegetation	Water Depth (cm)
Marsh passerines	*2.82 (0.54)	*0.74 (0.35)	NA	NA	NA
Secretive marsh birds	NA	NA	NA	*2.37 (0.50)	NA
Marsh Wren	*1.72 (0.53)	NA	NA	NA	*-0.42 (0.15)
Yellow-headed Blackbird	*2.63 (0.55)	*1.47 (0.32)	NA	NA	*0.40 (0.16)
Virginia Rail	NA	NA	*0.55 (0.23)	NA	*-0.19 (0.11)

The NA represents both variables that were initially in the global model but not included in the final model and variables not considered in the global model. Significant ($P < 0.05$) effects are indicated with an asterisk (*)

the surface, so they are more sensitive to low light availability (Lougheed et al. 2001). These taxa were less abundant in non-restored sites, and *Urticularia vulgaris* showed a quadratic relationship with years since restoration. This, along with the changes in floating-leaved vegetation, may indicate that these sites could revert to a eutrophic, turbid state.

Changes in the macrophyte structure and composition have the potential to influence avian communities in prairie wetlands (Fairbairn and Dinsmore 2001; Rehm and Baldassarre 2007; Harms and Dinsmore 2013), and we found this to be consistent in our study sites. Importance of emergent vegetation, particularly *Typha* sp., was evident for almost all of our focal groups and species. The positive relationship between *Typha* sp. and counts of breeding marsh birds has been found by other studies and in other regions of North America (Tozer et al. 2010; Harms and Dinsmore 2013; Lupien et al. 2014). In particular, the robust leaves of *Typha* sp. provide an appropriate surface for nest building by breeding marsh passerines (Twedt and Crawford 1995; Mowbray 1997; Kroodsma and Verner 2013). Both Marsh Wrens and Yellow-headed Blackbirds use emergent vegetation as a surface on which to build their nests (Verner 1965; Willson 1966). Marsh Wrens primarily used *Typha* sp. in these shallow lakes, although they may use other dominant robust emergent species such as *Schoenoplectus* sp. if there is standing water (Verner 1965; Verner and Engelsen 1970).

Along with *Typha* sp., we found that water depth influenced numbers of Marsh Wrens and Yellow-headed Blackbirds. Both species commonly build nests over deep water (Willson 1966; Minock and Watson 1983; Twedt and Crawford 1995), and we found that water depth had a significant positive effect on Yellow-headed Blackbird numbers. On the other hand, Marsh Wrens showed a negative relationship with water depth. Other studies have found a positive trend between water depth and Marsh Wren abundance or occupancy (Tozer et al. 2010; Lupien et al. 2014). However, unlike these studies, Yellow-headed Blackbirds were common breeders and tended to occur in the same sites as Marsh Wrens in our study area. Both species are aggressive territory defenders and Yellow-headed Blackbirds are known to exclude Marsh Wrens from their territories (Bump 1986; Harms and Dinsmore 2015). In this case, it could be that Yellow-headed Blackbirds exclude Marsh Wrens from areas with deeper water.

We expected secretive marsh birds and Virginia Rails to be influenced by frequency of *Typha* sp., but we found that secretive marsh birds were positively influenced by the amount of moderately dense vegetation. Virginia Rails did not show any significant relationship with emergent vegetation characteristics. Overall vegetation structure and water conditions may be more important than emergent species composition for this group and species (Rundle and Fredrickson 1981). Robust and dense stands of emergent vegetation can provide appropriate habitat conditions for several species of secretive marsh birds (Lor and Malecki 2006; Tozer et al. 2010; Harms and Dinsmore 2013). Several species, including Least Bitterns and Virginia Rails, will build their nests at the base of emergent vegetation and may use the leaves as a canopy (Weller 1961; Conway 1995). At the same time, some degree of interspersion between vegetation and water tends to provide appropriate nesting conditions for a greater variety of secretive marsh birds (Weller and Spatcher 1965; Lor and Malecki 2006), which might explain the importance of moderately dense vegetation. On the other hand, water depth was a more important factor for Virginia Rails, which showed a negative relationship with water depth. Virginia Rails generally use and nest in areas with shallow water (< 20 cm; Rundle and Fredrickson 1981, Sayre and Rundle 1984, Lor and Malecki 2006). Indeed, we found that Virginia Rails had a negative relationship with water depth.

Along with emergent vegetation, submersed and floating aquatic species appeared to be important for some groups. Both marsh passerines and Yellow-headed Blackbirds showed a positive relationship with frequency of submersed aquatic vegetation. This vegetation can provide habitat for a variety of macroinvertebrates (Krull 1970; Voigts 1976; Driver 1977; Hanson and Butler 1994), which are important in the diets of breeding marsh birds (Twedt and Crawford 1995). Yellow-headed Blackbirds and Marsh Wrens will forage on emerging aquatic insects at the water's surface (Twedt and

Crawford 1995; Kroodsma and Verner 2013). These and other breeding marsh birds will also nest over deep water, which can provide appropriate conditions for submersed aquatic vegetation (Sheldon and Boylen 1977). Indeed, some studies have found a positive relationship between the abundance of these species and water depth (Tozer et al. 2010). Similarly, floating-leaved vegetation can also provide habitat for macroinvertebrates (Harper and Bolen 1996), which may also explain this variable's relationship with Virginia Rails. Furthermore, Virginia Rails may use the dense mats of floating-leaved vegetation as a walking surface (Conway 1995).

Management Implications

Our results showed notable differences among the vegetation communities of shallow lakes in different restoration states, which are likely impacting numbers of breeding marsh birds. Because most of our study sites are isolated and exist in a matrix of cropland, they still face pressures from sedimentation, increased nutrient loading, and altered water regimes (Neely and Baker 1989; Anteau 2012; Mushet et al. 2015; Van Meter and Basu 2015). Implementing drawdowns or periodically lowering water levels may be effective management tools for maintaining some years of clear water and abundant submersed aquatic vegetation (Scheffer 1998; Søndergaard et al. 2007). Based on our findings a drawdown every 7 years may be a way to reset the wetland, but further research on restored shallow lakes >7 years of age may be necessary to confirm this timeframe. To increase vegetation species richness, active seeding during drawdown and active removal may prevent monocultures of *Typha* sp. or *Phalaris arundinacea*, particularly in wet prairie or sedge meadow areas (Green and Galatowitsch 2001; Galatowitsch 2006; Lishawa et al. 2010). At the same time, these vegetation changes due to restoration are likely affecting marsh bird abundance. We realize that specific habitat requirements for species in the marsh passerine and secretive marsh bird group are known to differ (e.g., Harms and Dinsmore 2013), but our results may offer some broad implications for improving habitat for the marsh bird community. *Typha* sp. was especially important to several marsh birds, as this emergent species provides shelter and nesting habtiat. At the same time, frequency of submersed aquatic and floating-leaved vegetation also positively influenced marsh bird numbers. Thus, with such large wetlands, maintaining areas with dense emergent cover interspersed with submersed and floating plants will likely provide the greatest variety of habitat types for several species. Furthermore, managing water levels may be necessary to provide a variety of nesting habitats, as water level showed different affects for Yellow-headed Blackbirds and Virginia Rails.

Acknowledgements Funding for this project was provided by the U.S. Fish and Wildlife Service (Prairie Pothole Joint Venture, grant number F15AP00612) and the Iowa Department of Natural Resources (Resource Enhancement and Protection License Plate Funds). We thank Sarah Moodie for assistance with field work. We thank Brian J. Wilsey and Timothy W. Stewart for providing edits and advice on field methods and analyses. We also thank Mark Gulick, Todd Bishop, Doug Janke, Bryan Hellyer, Steve Woodruff, TJ Herrick, Clint Maddix, Chris LaRue, Karen Kinkead, Kevin Murphy, and Tyler Harms for help with planning and logistics.

References

Anteau MJ (2012) Do interactions of land use and climate affect productivity of waterbirds and prairie-pothole wetlands? Wetlands 32:1–9

Anteau MJ, Afton AD (2008) Amphipod densities and indices of wetland quality across the upper-Midwest, USA. Wetlands 28:184–196

Aronson MFJ, Galatowitsch S (2008) Long-term vegetation development of restored priarie pothole wetlands. Wetlands 28:883–895

Badiou P, McDougal R, Pennock D, Clark B (2011) Greenhouse gas emissions and carbon sequestration potential in restored wetlands of the Canadian prairie pothole region. Wetlands Ecology and Management 19:237–256

Baker LA (1992) Introduction to nonpoint source pollution in the United States and prospects for wetland use. Ecological Engineering 1:1–26

Banks RC, Springer PF (1994) A century of population trends of waterfowl in western North America. In: Jehl JR, Johnson NK (eds) A century of avifaunal change in western North America. Allen Press, Inc., Lawrence, pp 143–146

Bärlocher F, Mackay RJ, Wiggins GB (1978) Detritus processing in a temporary vernal pool in southern Ontario Canada. Archiv fuer Hydrobiologie 81:269–295

Bates D, Maechler M, Bolker B, Walker S (2015) Fitting linear mixed-effects models using lme4. Journal of Statistical Software 67(1):1–48. https://doi.org/10.18637/jss.v067.i01

Bini LM, Thomaz SM, Murphy KJ, Camargo AFM (1999) Hydrobiologia 415:147–154

Bishop RA (1981) Iowa's wetlands. Proceedings of the Iowa Academy of Science 88:11–16

Bortolotti LE, Vinebrooke RD, St. Louis VL (2016) Prairie wetland communities recover at different rates following hydrological restoration. Freshwater Biology 61:1874–1890

Brown SC, Smith CR (1998) Breeding season bird use of recently restored versus natural wetlands in New York. The Journal of Wildlife Management 62:1480–1491

Bump SR (1986) Yellow-headed blackbird nest defense: aggressive responses to Marsh Wrens. Condor 88:328–335

Carpenter SR, Lodge DM (1986) Effects of submersed macrophytes on ecosystem processes. Aquatic Botany 26:341–337

Chow-Fraser P (2005) Ecosystem response to changes in water level of Lake Ontario marshes: lessons from the restoration of Cootes Paradise Marsh. Hydrobiologia 539:189–204

Connor KJ, Gabor S (2006) Breeding waterbird wetland habitat availability and response to water-level management in Sain John River floodplain wetlands, New Brunswick. Hydrobiologia 567:169–181

Conway CJ (1995) Virginia rail (*Rallus limicola*). In: Poole Af, Gill FB (eds) The birds of North America online. Ithaca, Cornell lab of ornithology, Retrieved from the birds of North America online: https://doi.org/10.2173/bna.173

Conway CJ (2009) Standardized North American bird monitoring protocols, version 2009–2. Wildlife Research Report #2009–02. U.S. Geological Survey, Arizona Cooperative Fish and Wildlife Research Unit, Tucson, AZ

Conway CJ (2011) Standardized north American marsh bird monitoring protocol. Waterbirds 34:319–346

Conway CJ, Eddleman WR, Anderson SH (1994) Nesting success and survival of Virginia rails and Soras. Wilson Bulletin of Ornithology 106:466–473

Conway CJ, Sulzman C, Raulston BE (2004) Factors affecting detection probability of California black rails. Journal of Wildlife Management. 68:360–370

Cowardin LM, Carter V, Golet FC, LaRoe ET (1979) Classification of wetlands and deepwater habitats of the United States. In: F. a. W. S. U.S. Department of the Interior (ed), Washington D.C

Cumbee JC Jr, Gaines KF, Mills GL, Garvin N, Stephens WL Jr, Novak JM, Brisbin IL Jr (2008) Clapper rails as indicators of mercury and PCB and bioavailability in a Georgia saltmarsh system. Ecotoxicology 17:485–494

Dahl TE (2014) Status and trends of prairie wetlands in the United States 1997 to 2009. U.S. Department of the Interior, Fish and Wildlife Service. Washington D.C

Driver EA (1977) Chironomid communities in small prairie ponds: some characteristics and controls. Freshwater Biology 7:121–133

Eddleman WR, Knopf FL, Meanley B, Reid RA, Zembal R (1988) Conservation of north American Rallids. Wilson Bulletin 100: 458–475

Elphick CS, Oring LW (1998) Winter management of Californian rice fields for waterbirds. Journal of Applied Ecology 35:95–108

Euliss NH Jr, Mushet DM (1996) Water-level fluctuation in wetlands as a function of landscape condition in the prairie pothole region. Wetlands 16:587–593

Fairbairn SE, Dinsmore JJ (2001) Local and landscape-level influences on wetland bird communities of the prairie pothole region of Iowa, USA. Wetlands 21:41–47

Figuerola JA, Green J, Santamaría L (2002) Comparative dispersal effectiveness of wigeongrass seeds by waterfowl wintering in south-West Spain: quantitative and qualitative aspects. Journal of Ecology 90: 989–1001

Galatowitsch SM (2006) Restoring prairie pothole wetlands: does the species pool concept offer decision-making guidance for re-vegetation? Applied Vegetation Science 9:261–270

Galatowitsch SM, van der Valk AG (1994) Restoring prairie wetlands: an ecological approach. Iowa State University Press, Ames

Galatowitsch SM, van der Valk AG (1996a) Characteristics of recently restored wetlands in the prairie pothole region. Wetlands 16:75–83

Galatowitsch SM, van der Valk AG (1996b) The vegetation of restored and natural prairie wetlands. Ecological Applications 6:102–112

Games PA, Howell JF (1976) Pairwise multiple comparison procedures with unequal N's and/or variances: a Monte Carlo study. Journal of Educational Statistics 1:113–125

Geisthardt E, Gautsch J, Harland B (2013) 2013 Shallow lakes summary report. Iowa Department of Natural Resources, Watershed Monitoring and Assessment Section, Des Moines, IA. 65 pages

Gibbs JP, Melvin SM (1993) Call-response surveys for monitoring breeding waterbirds. Journal of Wildlife Management 57:27–34

Gleason RA, Euliss NH Jr, Hubbard DE, Duffy WG (2003) Effects of sediment load on emergence of aquatic invertebrates and plants from wetland soil egg and seed banks. Wetlands 23:26–34

Glisson WJ, Brady RS, Paulios AT, Jacobi SK, Larkin DJ (2015) Sensitivity of secretive marsh birds to vegetation condition in natural and restored wetlands in Wisconsin. Journal of Wildlife Management 79:1101–1116

Green EK, Galatowitsch SM (2001) Differences in wetland plant community establishment with additions of nitrate-N and invasive species (*Phalaris arundinaceae* and *Typha* x *glauca*). Canadian Journal of Botany 79:170–178

Hansel-Welch N, Butler MG, Carlson TJ, Hanson MA (2002) Changes in macrophyte community structure in Lake Christina (Minnesota), a large shallow lake, following biomanipulation. Aquatic Botany 75: 323–337

Hanson MA, Butler MG (1994) Response of plankton, turbidity, and macrophytes to biomanipulation in a shallow prairie lake. Canadian Journal of Fisheries and Aquatic Sciences 51:1180–1188

Hanson MA, Riggs MR (1995) Potential effects of fish predation on wetland invertebrates: a comparison of wetlands with and without fathead minnows. Wetlands 15:167–175

Hanson MA, Herwig BR, Zimmer KD, Hansel-Welch N (2017) Rehabilitation of shallow lakes: time to adjust expectations? Hydrobiologia 787:45–59

Harms TM (2011) Population ecology and monitoring of secretive marsh-birds in Iowa. Thesis. Iowa State University

Harms TM, Dinsmore SJ (2013) Habitat associations of secretive marsh birds in Iowa. Wetlands 33:561–571

Harms TM, Dinsmore SJ (2015) Spatial scale matters when modeling avian co-occurrence. Ecosphere 7(6):e01288. https://doi.org/10.1002/ecs2.1288

Harper CA, Bolen EG (1996) Duckweed (Lemnaceae) as habitat for macroinvertebrates in eastern North Carolina. Wetlands 16:240–244

Harris SW, Marshall WH (1963) Ecology of water-level manipulations on a northern marsh. Ecology 44:331–343

Hershey AE, Shannon L, Niemi GJ, Lima AR, Regal RR (1999) Prairie wetlands of south-central Minnesota: effects of drought on invertebrate communities. In: Batzer DP, Rader RB, Wissinger SA (eds) Invertebrates in freshwater wetlands of North America. Wiley, New York, pp 515–541

Igl LD, Johnson DH (1997) Changes in breeding bird populations in North Dakota: 1967 to 1992-93. The Auk 114:74–92

Iowa Association of Naturalists (IAN) (2001) Iowa Wetlands. http://www.extension.iastate.edu/Publications/IAN204.pdf. Accessed 15 Dec 2015

Jeppesen E, Jensen JP, Søndergaard M, Lauridsen R, Pedersen LJ, Jensen L (1997) Top-down control in freshwater lakes: the role of nutrient state, submerged macrophytes and water depth. Hydrobiologia 342(343):151–164

Jurik TW, Wang SC, van der Valk AG (1994) Effects of sediment load on seedling emergence from wetland seed banks. Wetlands 14:159–165

Kadlec JA, Murkin HR, van der Valk AG (2000) The baseline and deep-flooding years. In: Murkin HR, van der Valk AG, Clark WR (eds) Prairie wetland ecology: the contribution of the marsh ecology research program. Iowa State University Press, Ames, pp 55–74

Kaminski MR, Baldassarre GA, Pearse AT (2006) Waterbird responses to hydrological management of wetlands reserve program habitats in New York. Wildlife Society Bulletin 24:921–926

Kantrud HA, Millar JB, van der Valk AG (1989) Vegetation of wetlands of the prairie pothole region. In: van der Valk AG (ed) Northern prairie wetlands. Iowa State University Press, Ames, pp 132–187

Kettenring KM, Galatowitsch SM (2011) Seed rain of restored and natural prairie wetlands. Wetlands 31:283–294

Kroodsma DE, Verner J (2013) Marsh Wren (*Cistothorus palustris*). In: Poole AF (ed) The birds of North America online. Ithaca, Cornell lab of ornithology, Retrieved from the birds of North America online: https://doi.org/10.2173/bna.308

Krull JN (1970) Aquatic plant-macroinvertebrate associations and waterfowl. The Journal of Wildlife Management 34:707–718

Kruskal WH, Wallis WA (1952) Use of ranks in one-criterion variance analysis. Journal of the American Statistical Association 47:583–621

Kuznetsova A, Brockhoff PB, Christensen RHB (2017) lmerTest package: tests in linear mixed effects models. Journal of Statistical Software 82(13):1–26. https://doi.org/10.18637/jss.v082.i13

Lauridsen TL, Jeppesen E, Anderson FO (1993) Colonization of submerged macrophytes in shallow fish manipulated Lake Væng: impact of sediment composition and waterfowl grazing. Aquatic Botany 46:1–15

Lauridsen TL, Jeppesen E, Søndergaard M (1994) Colonization and succession of submerged macrophytes in shallow Lake Væng during the first five years following fish manipulation. Hydrobiologia 275(276):233–242

Lauridsen TL, Sandsten H, Moller PH (2003a) The restoration of a shallow lake by introducing *Potamogeton* spp.: the impact of waterfowl grazing. Lakes & Reservoirs: Research and Managemetn 8:177–187

Lauridsen TL, Jensen JP, E. Jeppesen, Søndergaard M (2003b) Response of submerged macrophytes in Danish lakes to nutrient loading reductions and biomanipulation. Hydrobiologia 506:641–649

Lishawa SC, Albert DA, Tuchman NC (2010) Water level deviance promotes *Typha x glauca* establishment and vegetation change in Great Lakes coastal wetlands. Wetlands 30:1085–1096

Lishawa SC, Treering DJ, Vail LM, McKenna O, Grimm EC, Tuchman NC (2013) Reconstructing plant invasions using historical aerial imagery and pollen core analysis: *Typha* in the Laurentian Great Lakes. Diversity and Distributions 19:14–28

Lor S, Malecki RA (2002) Call-response surveys to monitor marsh bird population trends. Wildlife Society Bulletin 30:1195–1201

Lor S, Malecki RA (2006) Breeding ecology and nesting habitat associations of five marsh bird species in western New York. Waterbirds 29:427–436

Lougheed VL, Crosbie B, Chow-Fraser P (2001) Primary determinants of macrophyte community structure in 62 marshes across the Great Lakes basin: latitude, land use, and water quality effects. Canadian Journal of Fisheries and Aquatic Sciences 58:1603–1612

Lowther PE, Poole AF, Gibbs JP, Melvin SM, Reid FA (2009) American bittern (*Botaurus lentiginosus*). In: Rodewald PF (ed) The birds of North America online (P. G. Rodewald, Ed.). Ithaca, Cornell lab of ornithology, Retrieved from the birds of norther America online: https://doi.org/10.2173/bna.18

Lumsden HG, Thomas VG, Robinson BR (2015) Response of wild trumpeter swan (*Cygnus buccinators*) broods to wetland drawdown and changes in food abundance. Canadian Field-Naturalist 129:374–378

Lupien NG, Gauthier G, Lavoie C (2014) Effect of the invasive common reed on the abundance, richness and diversity of birds in freshwater marshes. Animal Conservation 18:32–43

Martin DB, Hartman WA (1986) The effect of cultivation on sediment composition and deposition in prairie pothole wetlands. Water, Air, and Soil Pollution 34:45–53

McLean KI, Mushet DM, Renton DA, Stockwell CA (2016) Aquatic-macroinvertebrate communities of prairie-pothole wetlands and lakes under a changed climate. Wetlands 36:S421–S435

Miller BA, Crumpton WG, van der Valk AG (2009) Spatial distribution of historical wetland classes on the Des Moines Lobe, Iowa. Wetlands 29:1146–1152

Miller BA, Crumpton WG, van der Valk AG (2012) Wetland hydrologic class change from prior to European settlement to present on the Des Moines Lobe, Iowa. Wetlands Ecology and Management 20:1–8

Minc LD (1997) Great Lakes coastal wetlands: an overview of controlling abiotic factors, regional distribution and species composition. A report submitted to the Michigan Natural Features Inventory, December 1997. Funded by EPA Great Lakes National Program Office (Federal Grant GL9 95810–02) through The Nature Conservancy's Great Lakes Program Office. Great Lakes National Program Office, U.S. EPA, 77 West Jackson Blvd., Chicago, IL 60604

Minock ME, Watson JR (1983) Red-winged and yellow-headed blackbird nesting habitat in a Wisconsin marsh. Journal of Field Ornithology 54:324–326

Mitchell ME, Lishawa SC, Geddes P, Larkin DJ, Treering D, Tuchman NC (2011) Time-dependent impacts of cattail invasion in a Great Lakes coastal wetland complex. Wetlands 31:1143–1149

Monfils MJ, Brown PW, Hayes DB, Soulliere GJ, Kafcas EN (2014) Breeding bird use and wetland characteristics of diked and undiked coastal marshes in Michigan. The Journal of Wildlife Management 78:79–92

Montgomery DC, Peck EA (1992) Introduction to linear regression analysis. Wiley, New York

Mowbray TB (1997) Swamp sparrow (*Melospiza georgiana*). In: Rodewald PG (ed) The birds of North America online. Ithaca, Cornell lab of ornithology, Retrieved from the birds of North America online: https://doi.org/10.2173/bna.279

Murkin HR, Ross LC (2000) Invertebrates in prairie wetlands. In: Murkin HR, van der Valk AG, Clark WR (eds) Prairie wetland ecology: the contribution of the marsh ecology research program. Iowa State University Press, Ames, pp 201–247

Murkin HR, Murkin EJ, Ball JP (1997) Avian habitat selection and prairie wetland dynamics: a 10-year experiment. Ecological Applications 7: 1144–1159

Mushet DM, Goldhaber MB, Mills CT, McLean KI, Aparicio VM, McCleskey RB, Holloway JM, Stockwell CA (2015) Chemical and biotic characteristics of prairie lakes and large wetlands in south-central North Dakota – effects of a changing climate. US Geological Survey Scientific Investigations Report:2015–5126, 55 p. https://doi.org/10.3133/sir20155126

Neely RK, Baker JK (1989) Nitrogen and phosphorus dynamics and the fate of agricultural runoff. In: van der Valk AG (ed) Northern prairie wetlands. Iowa State University Press, Ames, pp 92–131

Page GW, Gill RE Jr (1994) horebirds in western North America late 1800s to late 1900s. In: Jehl JR, Johnson NK (eds) A century of avifaunal change in western North America. Allen Press, Inc., Lawrence, pp 147–160

Peterka JJ (1989) Fishes in northern prairie wetlands. In: age v d V (ed) Northern prairie wetlands. Iowa State University Press, Ames, pp 302–315

Pothoff AJ, Herwig BR, Hanson MA, Zimmer KD, Butler MG, Reed JR, Parsons BG, Ward MC (2008) Cascading food-web effects of piscivore introductions in shallow lakes. Journal of Applied Ecology 45: 1170–1179

Ralph CJ, Droege S, Sauer JR (1995) Managing and monitoring birds using point counts: standards and applications. In: Ralph CJ, Sauer JR, Droege S (eds) Monitoring bird populations by point counts. USDA Forest Service General Technical Report. PSW-GTR-149. Pp 161–175

Rehm EM, Baldassarre GA (2007) The influence of interspersion on marsh bird abundance in New York. The Wilson Journal of Ornithology 119:648–654

Rundle WD, Fredrickson LH (1981) Managing seasonally flooded impoundments for migrant rails and shorebirds. Wildlife Society Bulletin 9:80–87

Sayer CD, Davidson TA, Jones JI (2010) Seasonal dynamics of macrophytes and phytoplankton in shallow lakes: a eutrophication-driven pathway from plants to plankton? Freshwater Biology 55:500–513

Sayre MW, Rundle WD (1984) Comparison of habitat use by migrant Soras and Virginia rails. The Journal of Wildlife Management 48: 599–605

Scheffer M (1998) Ecology of shallow lakes. Chapman & Hall, Suffolk

Scheffer M, Hosper SH, Meijer M-L, Moss B, Jeppesen E (1993) Alternative equilibria in shallow lakes. Tree 8:275–279

Sheldon RB, Boylen CW (1977) Maximum depth inhabited by aquatic vascular plants. American Midland Naturalist 97:248–254

Smith MD, Moelyowati I (1998) Duckweed based wastewatertreatment (DWWT): design guidelines for hot climates. Water Science and Technology 43:291–299

Søndergaard M, Jeppesen E, Lauridsen TK, Skov C, Van Nes EH, Roijackers R, Lammens E, Portielje R (2007) Lake restoration: successes, failures and long-term effects. Journal of Applied Ecology 44:1095–1105

Stewart TW, Downing JA (2008) Macroinvertebrate communities and environmental conditions in recently constructed wetlands. Wetlands 28:141–150

Stewart RE, Kantrud HA (1971) Classification of natural ponds and lakes in the glaciated prairie region. In: FWS US Department of the Interior (ed) Bureau of Sport Fisheries and Wildlife, Washington D.C.

Strand JA (1999) The development of submerged macrophytes in Lake Ringsjön after biomanipulation. Hydrobiologia 404:113–121

Taft OW, Colwell MA, Isola CR, Safran RJ (2002) Waterbird responses to experimental drawdown: implications for the multispecies management of wetland mosaics. Journal of Applied Ecology 39:987–1001

Timms RM, Moss B (1984) Prevention of growth of potentially dense phytoplankton populations by zooplankton grazing, in the presence of zooplanktivorous fish, in a shallow wetland ecosystem. Limnology and Oceanography 29:472–486

Tiner RW Jr (1984) Wetlands of the United States: Current status and recent trends. National Wetlands Inventory. U.S. Department of the Interior, Fish and Wildlife Service, Washington

Tozer DC, Nol E, Abraham KF (2010) Effects of local and landscape-scale habitat variables on abundance and reproductive success of wetland birds. Wetlands Ecology Management 18:679–693

Twedt DJ, Crawford RD (1995) Yellow-headed blackbird (*Xanthocephalus xanthocephalus*). In: Rodewald PG (ed) The birds of North America online. Ithaca, Cornell lab of ornithology, Retrieved from the birds of North America online: https://doi.org/10.2173/bna.192

van der Valk AG, Davis CB (1976) The seed banks of prairie glacial marshes. The Canadian Journal of Botany 54:1832–1838

van der Valk AG, Davis CB (1978) The role of seed banks in vegetation dynamics of prairie glacial marshes. Ecology 59:322–335

van der Valk AG, Murkin HR, Kadlec JA (2000) The drawdown and reflooding years. In: Murkin HR, van der Valk AG (eds) Prairie wetland ecology: the contribution of the marsh ecology research program. Iowa State University Press, Ames

Van Meter KJ, Basu NB (2015) Signatures of human impact: size distributions and spatial organization of wetlands in the prairie pothole landscape. Ecological Applications 25:451–465

VanRees-Siewert KL, Dinsmore JJ (1996) Influence of wetland age on bird use of restored wetlands in Iowa. Wetlands 16:577–582

Verner J (1965) Breeding biology of the long-billed Marsh Wren. Condor 67:6–30

Verner J, Engelsen GH (1970) Territories, multiple nest building, and polygyny in the long-billed marsh wren. Auk 87:557–567

Voigts DK (1976) Aquatic invertebrate abundance in relation to changing marsh vegetation. The American Midland Naturalist 95:313–322

Webb EB, Smith LM, Vrtiska MP, Lagrange TG (2010) Effects of local and landscape variables on wetland bird habitat use during migration through the Rainwater Basin. Journal of Wildlife Management 74: 109–119

Weller MW (1961) Breeding biology of the least bittern. The Wilson Bulletin 73:11–35

Weller MW, Spatcher CS (1965) Role of habitat in the distribution and abundance of marsh birds. Iowa Agriculture and Home Economics Experiment Station Report 43, Ames, Iowa

Wienhold CE, van der Valk AG (1989) The impact of duration of drainage on the seed banks of northern prairie wetlands. Canadian Journal of Botany 67:1878–1884

Willson MF (1966) Breeding ecology of the yellow-headed blackbird. Ecological Monographs 36:51–77

Wrubleski DA (1999) Northern prairie marshes (Delta marsh, Manitoba): II. Chironomidae (Diptera) responses to changing plant communities in newly flooded habitats. In: Batzer DP, Rader RB, Wissinger SA (eds) Invertebrates in freshwater wetlands of North America. Wiley, New York, pp 571–601

Zar JH (2010) Biostatistical analysis, 5th edn. Pearson Prentice Hall, Upper Saddle River

Zimmer KD, Hanson MA, Butler MG (2000) Factors influencing invertebrate communities in prairie wetlands: a multivariate approach. Canadian Journal of Fisheries and Aquatic Sciences 57:76–85

Zimmer, KD, Hanson MA, Butler MG (2001) Effects of fathead minnow colonization and removal on a prairie wetland ecosystem. Ecosystems 4:346–357

Zimmer KD, Hanson MA, Butler MG (2002) Effects of fathead minnows and restoration on prairie wetland ecosystems. Freshwater Biology 47:2071–2086

Zimmer KD, Hanson MA, Butler MG (2003) Relationships among nutrients, phytoplankton, macrophytes, and fish in prairie wetlands. Canadian Journal of Fisheries and Aquatic Sciences 60:721–730

Zuur AF, Ieno EN, Walker NJ, Saveliev AA, Smith GM (2009) Mixed effects models and extensions in ecology with R. Springer, New York