

A note to paper: Axial symmetry and classification of stationary solutions of Doi-Onsager equation on the sphere with Maier-Saupe potential

Hailiang Liu¹, Hui Zhang², and Pingwen Zhang³

ABSTRACT. This note serves to provide additional details for the proof of Lemma 3.6 in our paper [Liu, Zhang and Zhang, Comm. Math. Sci., 3(2005), pp.201-218]. Moreover, we will also present an alternative, yet simpler, proof based on arguments in [Wang, Zhang and Zhang, CPAM, 68(2015), no. 8, 1326-1398].

In [1] we showed that in order to determine the number of solutions to the Doi-Onsager equation on the sphere with Maier-Saupe potential, it suffices to determine the number of zeros of $B(\eta, \alpha)$ in term of the intensity parameter $\alpha > 0$, where

$$B(\eta, \alpha) = \frac{3e^{-\eta}}{\int_0^1 e^{-\eta z^2} dz} - \left(3 - 2\eta + \frac{4\eta^2}{\alpha} \right). \quad (3.40)$$

Let us state the original Lemma 3.6 in [1], and reproduce its proof with supplemental details mainly in steps 2 and 5. The equation numbers follow those numbered in [1].

Lemma 3.6. *The number of zeros of $B(\eta, \alpha)$ is determined by the intensity α as follows:*

- (i). *If $\alpha > 7.5$, $B(\eta, \alpha)$ has three zeros $\eta_1^* < 0, \eta_2^* > 0$ and $\eta_0 = 0$.*
- (ii). *If $\alpha = 7.5$, $B(\eta, \alpha)$ has two zeros $\eta_1^* < 0$ and $\eta_0 = 0$.*
- (iii). *There exists an $\alpha^* \in (20/3, 7.5)$ such that $B(\eta, \alpha)$ has three zeros $\eta_1^* < 0, \eta_2^* < 0$ and $\eta_0 = 0$ for $\alpha^* < \alpha < 7.5$.*
- (iv). *If $\alpha = \alpha^*$, $B(\eta, \alpha)$ has two zeros $\eta_1^* < 0$ and $\eta_0 = 0$.*
- (v). *If $0 < \alpha < \alpha^*$, $B(\eta, \alpha)$ has one zero $\eta_0 = 0$.*

Proof. This proof is divided into five steps.

Step 1. We first show that

$$B(\pm M, \alpha) < 0 \quad \text{for } M \gg 1. \quad (3.41)$$

¹Department of Mathematics, Iowa State University, Ames, IA 50011-2064, U.S., hliu@iastate.edu

²LMCS and School of Mathematical Sciences, Beijing Normal University, Beijing, 100875, P.R. China, hzhang@bnu.edu.cn

³LMAM and School of Mathematical Sciences, Peking University, Beijing, 100871, P.R. China, pzhang@pku.edu.cn

Note that, for $\eta > 0$, the mean value theorem gives $3e^{-\eta}/\int_0^1 e^{-\eta z^2} dz = 3e^{-\eta(1-\gamma^2)} \in (0, 3)$ for some $\gamma \in (0, 1)$. For $\eta < 0$, we have from (3.39)

$$e^{-\eta} = \int_0^1 e^{-\eta z^2} dz - 2\eta \int_0^1 z^2 e^{-\eta z^2} dz \leq (1 - 2\eta) \int_0^1 e^{-\eta z^2} dz.$$

This shows $0 < \frac{3e^{-\eta}}{\int_0^1 e^{-\eta z^2} dz} \leq 3(1 - 2\eta)$. Thus for large $|\eta| \gg 1$, B is determined by the quadratic term $-4\eta^2/\alpha$. (3.41) follows.

Step 2. If $\alpha > 7.5$ we will show that $B(\eta, \alpha)$ has at least two zeros $\eta_1 < 0$, and $\eta_2 > 0$. We know that $\eta = 0$ is always a zero since $B(0, \alpha) = 0$. Moreover, let $A_k = \int_0^1 z^k e^{-\eta z^2} dz$, then $\partial_\eta A_k = -A_{k+2}$ for any integer $k \geq 0$. By using (3.40) we have

$$B_\eta(\eta, \alpha) = 3e^{-\eta} \frac{A_2 - A_0}{A_0^2} - \left(-2 + \frac{8\eta}{\alpha} \right). \quad (3.42)$$

This yields $B_\eta(0, \alpha) = 0$. Therefore, $\eta = 0$ is a double zero of B for every α . Hence the local shape of B hinges on the sign of $B_{\eta\eta}(0, \alpha)$. From (3.42), we have

$$\begin{aligned} B_{\eta\eta}(\eta, \alpha) &= \frac{3e^{-\eta}}{A_0^3} (A_0^2 - 2A_0A_2 - A_0A_4 + 2A_2^2) - \frac{8}{\alpha} \\ &= \frac{3e^{-\eta}}{A_0^3} [A_0(A_0 - 3A_2) + A_0(A_2 - A_4) + 2A_2^2] - \frac{8}{\alpha}. \end{aligned} \quad (3.43)$$

Therefore a further calculation from (3.43) gives

$$B_{\eta\eta}(0, \alpha) = \frac{16}{15\alpha} \left(\alpha - \frac{15}{2} \right). \quad (3.44)$$

Thus $B_{\eta\eta}(0, \alpha) > 0$ for $\alpha > 7.5$. B is locally convex near $\eta = 0$. This together with (3.41) implies that there exist at least two zeros $\eta_1^* < 0$ and $\eta_2^* > 0$ besides $\eta = 0$. Now we assume η^* is a zero of B , i.e.,

$$\frac{3e^{-\eta^*}}{\int_0^1 e^{-\eta^* z^2} dz} = 3 - 2\eta^* + \frac{4\eta^{*2}}{\alpha}. \quad (3.45)$$

In virtue of (3.38), i.e. $A_2 = \frac{\alpha - 2\eta}{3\alpha} A_0$, and (3.42) we have

$$\begin{aligned} B_\eta(\eta^*, \alpha) &= \frac{3e^{-\eta^*}}{\int_0^1 e^{-\eta^* z^2} dz} \left(-1 + \frac{\alpha - 2\eta^*}{3\alpha} \right) + 2 \left(1 - \frac{4\eta^*}{\alpha} \right) \\ &= \frac{-2e^{-\eta^*}}{\int_0^1 e^{-\eta^* z^2} dz} \left(1 + \frac{\eta^*}{\alpha} \right) + 2 \left(1 - \frac{4\eta^*}{\alpha} \right). \end{aligned} \quad (3.46)$$

Inserting (3.45) into (3.46) gives

$$B_\eta(\eta^*, \alpha) = -\frac{8\eta^*}{3\alpha^2} \left(\eta^{*2} + \frac{\eta^* \alpha}{2} + \frac{\alpha}{2} \left(\frac{15}{2} - \alpha \right) \right) \quad (3.47)$$

$$= \begin{cases} -\frac{8\eta^*}{3\alpha^2} (\eta^* - \bar{\eta}_1)(\eta^* - \bar{\eta}_2), & \text{if } \alpha > 20/3, \\ -\frac{8\eta^*}{3\alpha^2} (\eta^* + \frac{\alpha}{4})^2, & \text{if } \alpha = 20/3, \\ -\frac{8\eta^*}{3\alpha^2} [(\eta^* + \frac{\alpha}{4})^2 + \frac{15\alpha}{4}(1 - \frac{3\alpha}{20})], & \text{if } \alpha < 20/3, \end{cases} \quad (3.48)$$

where

$$\bar{\eta}_1 = -\frac{\alpha}{4} \left(1 + 3\sqrt{1 - \frac{20}{3\alpha}} \right), \quad \bar{\eta}_2 = -\frac{\alpha}{4} \left(1 - 3\sqrt{1 - \frac{20}{3\alpha}} \right). \quad (3.49)$$

From (3.48), we see that $\eta_1^* \leq \bar{\eta}_1 < -\alpha/2$, and $\eta_2^* \geq \bar{\eta}_2 > 0$.

In fact it is impossible that $\eta_1^* = \bar{\eta}_1$ or $\eta_2^* = \bar{\eta}_2$. If $\eta_1^* = \bar{\eta}_1$ or $\eta_2^* = \bar{\eta}_2$, then $B_\eta(\eta^*, \alpha) = 0$ from (3.48); we would have to calculate $B_{\eta\eta}(\eta^*, \alpha)$ to reach a contradiction. For $\eta = \eta^*$, from (3.38) we have $\alpha = \frac{2\eta^* A_0}{A_0 - 3A_2}$, that gives $A_2/A_0 = \frac{\alpha - 2\eta^*}{3\alpha}$. Using the fact $B_\eta(\eta^*, \alpha) = 0$ we also obtain $\alpha = \frac{A_0}{A_2 - A_4}$. Thus

$$\begin{aligned} B_{\eta\eta}(\eta^*, \alpha) &= \frac{3e^{-\eta^*}}{A_0} \left[\frac{A_0 - 3A_2}{A_0} + \frac{A_2 - A_4}{A_0} + 2\frac{A_2^2}{A_0^2} \right] - \frac{8}{\alpha} \\ &= \frac{3e^{-\eta^*}}{A_0} \left(\frac{2\eta^*}{\alpha} + \frac{1}{\alpha} + \frac{2(\alpha - 2\eta^*)^2}{9\alpha^2} \right) - \frac{8}{\alpha} \\ &= \frac{1}{\alpha} \left(3 - 2\eta^* + \frac{4\eta^{*2}}{\alpha} \right) \left(2\eta^* + 1 + \frac{2(\alpha - 2\eta^*)^2}{9\alpha} \right) - \frac{8}{\alpha}, \end{aligned}$$

where we have used (3.45). Now we denote

$$A = \eta^2 + \frac{\eta\alpha}{2} + \frac{\alpha}{2} \left(\frac{15}{2} - \alpha \right) = (\eta - \bar{\eta}_1)(\eta - \bar{\eta}_2).$$

If $\eta^* = \bar{\eta}_1$ or $\bar{\eta}_2$, then $A = 0$ at $\eta = \eta^*$. Therefore we get

$$\begin{aligned} B_{\eta\eta}(\eta^*, \alpha) &= \frac{1}{\alpha} (3 - 2\eta^* - 2\eta^* - 15 + 2\alpha) \\ &\quad \cdot \left(2\eta^* + 1 + \frac{2\alpha}{9} - \frac{8\eta^*}{9} + \frac{8}{9} \left[-\frac{\eta^*}{2} - \frac{1}{2} \left(\frac{15}{2} - \alpha \right) \right] \right) - \frac{8}{\alpha} \\ &= \frac{2}{3\alpha} ((\alpha - 2\eta^* - 6)(2\alpha + 2\eta^* - 7) - 12) \\ &= \frac{2}{3\alpha} (2\alpha^2 - 2\alpha\eta^* - 4\eta^{*2} - 19\alpha + 2\eta^* + 30) \\ &= \frac{4}{3\alpha} (\eta^* + 15 - 2\alpha). \end{aligned} \quad (3.50)$$

Apparently, for $\alpha > 7.5$, $B_{\eta\eta}(\eta_1^*, \alpha) < 0$ as $\eta_1^* < 0$. For $\eta_2^* = \bar{\eta}_2 = -\frac{\alpha}{4} \left(1 - 3\sqrt{1 - \frac{20}{3\alpha}} \right)$, it holds (recalling $\alpha > 15/2$)

$$B_{\eta\eta}(\eta_2^*, \alpha) = -\frac{1}{3} \left(1 - 3\sqrt{1 - \frac{20}{3\alpha}} \right) + \frac{8}{3\alpha} \left(\frac{15}{2} - \alpha \right) = \sqrt{\frac{3\alpha - 20}{3\alpha}} \left(1 - \sqrt{\frac{3(3\alpha - 20)}{\alpha}} \right) < 0.$$

These facts imply that B is locally concave near η_i^* if $\eta_i^* = \bar{\eta}_i$ for $i = 1$ or 2 .

On the other hand, if $\eta_2^* = \bar{\eta}_2$ we know that B is locally convex near $\eta = 0$. Note $B_\eta(0, \alpha) = B_\eta(\eta_2^*, \alpha) = 0$. Therefore there is at least one zero point $\eta_3^* \in (0, \bar{\eta}_2)$ of $B(\eta, \alpha)$ which satisfies

$$B_\eta(\eta_3^*, \alpha) \leq 0.$$

However, this is impossible by (3.48). Therefore $\eta_2^* > \bar{\eta}_2$. Similarly we can get $\eta_1^* < \bar{\eta}_1$.

Step 3. We now show that $B(\eta, \alpha)$ has at most two zeros besides 0 for $\alpha > 7.5$. From (3.48), $B_\eta(\eta^*, \alpha) < 0$ for $\eta^* \in (\bar{\eta}_2, \infty)$. This implies that there is at most one zero of B in $(0, \infty)$. Otherwise $B_\eta(\eta^*, \alpha)$ has to be negative at another zero. Similarly, there exists at most one $\eta^* \in (-\infty, \bar{\eta}_1)$. The claim in (i) is thus proved.

Step 4. We now consider the case $\alpha = 7.5$, for which we show that there exist two zeros $\eta_1^* < 0$ and 0. In this case, $B_\eta(0, \alpha) = B_{\eta\eta}(0, \alpha) = 0$. In order to see the local shape of B at $\eta = 0$, we calculate $B_{\eta\eta\eta}(0, \alpha)$. From (3.43) we have by a careful calculation that

$$B_{\eta\eta\eta}(\eta, \alpha) = \frac{3e^{-\eta}}{A_0^4} [6A_2^3 - 6A_0A_2(A_2 + A_4) + A_0^2(3A_2 + 3A_4 + A_6) - A_0^3]. \quad (3.51)$$

So

$$B_{\eta\eta\eta}(0, \alpha) = -\frac{32}{105} < 0. \quad (3.52)$$

This means

$$\eta B(\eta, \alpha) < 0 \quad \text{for } |\eta| \ll 1. \quad (3.53)$$

(3.47) with $\alpha = 7.5$ gives

$$B_\eta(\eta^*, \alpha) = -\frac{8\eta^{*2}}{3\alpha^2} \left(\eta^* + \frac{\alpha}{2} \right), \quad (3.54)$$

where η^* is assumed to be a zero of B . The local behavior implied from (3.53) and the negative sign of $B_\eta(\eta^*, \alpha)$ for $\eta^* > 0$ shows that no zero of B exists in $(0, \infty)$. On the other hand, (3.53), together with $B(-M, \alpha) < 0$ shows that there exists at least one zero in $(-\infty, 0)$. We denote it by η_1^* . By (3.54), we know B has no zeros in $(-\alpha/2, 0)$ (otherwise consider the one closest to 0). Moreover, $\eta_1^* \neq -\alpha/2$ (otherwise $B_\eta(\eta_1^*, \alpha) = 0$ and by (3.50) $B_{\eta\eta}(\eta_1^*, \alpha) < 0$ which is impossible since B has no zeros in $(-\alpha/2, 0)$). Thus $\eta_1^* < -\alpha/2$ and $B_\eta(\eta_1^*, \alpha) > 0$. Now we claim η_1^* is a unique zero of B in $(-\infty, 0)$. Otherwise, there should appear at least two more zeros in $(-\infty, 0)$, which is not allowed by (3.54). This proves (ii).

Step 5. We can show that $B(\eta, \alpha)$ has no zero in $(0, \infty)$ for $\alpha < 7.5$. Otherwise, as $B(0, \alpha) = B_\eta(0, \alpha) = 0$ and $B_{\eta\eta}(0, \alpha) < 0$ (by (3.44)), there must be a positive zero of $B(\eta, \alpha)$ satisfying $B_\eta(\eta_1, \alpha) \geq 0$, which is impossible by (3.47). Moreover, if $\alpha \leq 20/3$, $B(\eta, \alpha)$ even has no zero in $(-\infty, 0)$. This is ensured by the fact $B(-M, \alpha) < 0$ and the sign constrained by (3.48), as argued in step 4.

In order to identify the second critical value $\alpha^* \in (20/3, 7.5)$, we need to use a continuity argument.

First for $7.5 - \delta < \alpha < 7.5$, $\delta > 0$ small, there are at least two zeros $\eta_1^*, \eta_2^* < 0$ of B . In fact for this range of α , $B_{\eta\eta}(0, \alpha) < 0$. Thus $B(\eta, \alpha)$ is locally concave near $\eta = 0$. We also know that $B(\eta, 7.5) > 0$ for $\eta \in (-\delta_1, 0)$ from (3.53). This implies that there exists a point $\eta_0 \in (-\delta_1, 0)$ such that $B(\eta_0, \alpha) > 0$ for $7.5 - \delta < \alpha < 7.5$ by the continuity of B in α . This together with $B(-M, \alpha) < 0$ shows that there are two zeros of B in $(-\infty, 0)$ for $7.5 - \delta < \alpha < 7.5$.

Secondly, we claim that for $\alpha < 7.5$, B has at most two zeros in $(-\infty, 0)$. This can be concluded from the following facts (see (3.49) for definitions of $\bar{\eta}_1, \bar{\eta}_2$, both are negative for $\alpha < 7.5$):

- (1). $B(\cdot, \alpha)$ has no zero point in $(\bar{\eta}_2, 0)$. Otherwise, let $\eta^* \in (\bar{\eta}_2, 0)$ be the largest zero point. Then $B_\eta(\eta^*, \alpha) < 0$ which is not allowed by (3.48);
- (2). $\bar{\eta}_2$ is not a zero point. Otherwise, by (3.50) we have $B_{\eta\eta}(\bar{\eta}_2, \alpha) > 0$ which contradicts to (3.48);
- (3). $B(\cdot, \alpha)$ has at most one zero point in $(\bar{\eta}_1, \bar{\eta}_2)$. Otherwise there is a zero point $\eta^* \in (\bar{\eta}_1, \bar{\eta}_2)$ satisfying $B_\eta(\eta^*, \alpha) \geq 0$ which is impossible by (3.48) again;
- (4). $B(\cdot, \alpha)$ has at most one zero point in $(-\infty, \bar{\eta}_1)$. Otherwise there is a zero point $\eta^* \in (-\infty, \bar{\eta}_1)$ satisfying $B_\eta(\eta^*, \alpha) \leq 0$ which also contradicts to (3.48);
- (5). If $\bar{\eta}_1$ is a zero point, then $B(\bar{\eta}_1, \alpha) = B_\eta(\bar{\eta}_1, \alpha) = 0, B_{\eta\eta}(\bar{\eta}_1, \alpha) < 0$ (using (3.50)). Thus repeating the argument in (3) or (4), we know there is no zero point of B in $(\bar{\eta}_1, \bar{\eta}_2)$ or $(-\infty, \bar{\eta}_1)$.

Now, choose $\alpha_0 \in (7.5 - \delta, 7.5)$ such that $B(\eta, \alpha_0)$ has exactly two zeros $\eta_{0,1} < \eta_{0,2}$ on $(-\infty, 0)$. Then $B(\cdot, \alpha_0)$ is negative on $(-\infty, \eta_{0,1}) \cup (\eta_{0,2}, 0)$. As $B(\eta, \cdot)$ is a monotonically increasing function of α , we know that for all $\alpha < \alpha_0$, $B(\eta, \alpha) < 0$ for $\eta \in (-\infty, \eta_{0,1}) \cup (\eta_{0,2}, 0)$. Let

$$\alpha^* = \sup\{\alpha < \alpha_0 \mid B(\eta, \alpha) < 0 \text{ for all } \eta \in (\eta_{0,1}, \eta_{0,2})\}. \quad (3.55)$$

Then for $\alpha < \alpha^*$, B has no zeros on $(-\infty, 0)$. Now we prove B has two zeros for $\alpha \in (\alpha^*, 7.5)$ and has one zero for $\alpha = \alpha^*$. Thus $\alpha^* \in (20/3, 7.5)$ is the corresponding critical value.

Apparently, $B(\eta, \alpha^*)$ must have (at least) a zero η^* in $(\eta_{0,1}, \eta_{0,2})$. Moreover, $B_\eta(\eta^*, \alpha^*) = 0$, and thus, (3.48) implies $\eta^* \in \{\bar{\eta}_1(\alpha^*), \bar{\eta}_2(\alpha^*)\}$. However, $\eta^* \neq \bar{\eta}_2$ by the fact (2), thus $\eta^* = \bar{\eta}_1(\alpha)$ is the unique zero of $B(\cdot, \alpha^*)$ in $(-\infty, 0)$. Since $B(\eta^*, \cdot)$ is monotonically increasing function of α , we know $B(\eta^*, \alpha) > 0$ for all $\alpha > \alpha^*$. Thus, there are exactly two zeros of B with one of them,

say η_1^* , belongs to $(-\infty, \eta^*)$ and another one $\eta_2^* \in (\eta^*, 0)$ for all $\alpha \in (\alpha^*, 7.5)$. Using again the fact that $B(\eta^*, \alpha)$ is monotonically increasing function of α , we know η_1^* is a decreasing function of α , while η_2^* is a increasing function of α .

These all together finish the proof of (iii)-(v). \square

An alternative proof:

The core part of this proof can be found in Lemma A.1 in [2].

Step 1: an equivalent formulation

Let $A_k(\eta) = \int_0^1 z^k e^{-\eta z^2} dz$. Then the use of integration by parts gives

$$(k+1)A_k - 2\eta A_{k+2} = e^{-\eta}, \text{ for } k \geq 0.$$

Consequently, we have

$$\begin{aligned} \frac{3e^{-\eta}}{\int_0^1 e^{-\eta z^2} dz} = 3 - 2\eta + \frac{4\eta^2}{\alpha} &\iff \frac{3(A_0 - 2\eta A_2)}{A_0} = 3 - 2\eta + \frac{4\eta^2}{\alpha} \\ \iff 4\eta^2 \left(\frac{A_2 - A_4}{A_0} - \frac{1}{\alpha} \right) = 0 &\iff \eta = 0, \text{ or } \alpha = \frac{A_0}{A_2 - A_4}. \end{aligned} \quad (3.56)$$

(This formulation was also shown in the end of [1], see the last line of pp. 217)

Step 2: number of solutions

It suffices to explore the solutions to $\alpha = f(\eta) := \frac{A_0(\eta)}{A_2(\eta) - A_4(\eta)}$. Using $A'_k(\eta) = -A_{k+2}$, we have

$$f'(\eta) = \left(\frac{A_0}{A_2 - A_4} \right)' = \frac{-A_2(A_2 - A_4) + A_0(A_4 - A_6)}{(A_2 - A_4)^2}.$$

Using the fact that

$$\begin{aligned} &\frac{\partial}{\partial \eta} \left(e^\eta (A_0(A_4 - A_6) - A_2(A_2 - A_4)) \right) \\ &= \frac{1}{2} \frac{\partial}{\partial \eta} \int_0^1 \int_0^1 (x^2 y^4 + x^4 y^2 - x^6 - y^6 + x^4 + y^4 - 2x^2 y^2) e^{-\eta(x^2 + y^2 - 1)} dx dy \\ &= \frac{1}{2} \int_0^1 \int_0^1 (x^2 - y^2)^2 (1 - x^2 - y^2)^2 e^{\eta(x^2 + y^2 - 1)} dx dy > 0, \end{aligned}$$

we deduce that $A_0(A_4 - A_6) - A_2(A_2 - A_4) = 0$ has only one root η_* , hence $f'(\eta)(\eta - \eta^*) > 0$ for $\eta \neq \eta^*$. Note that from $f'(0) = 5/7 > 0$ it follows that $\eta^* < 0$. Hence we have

Conclusion 1: $f(\eta)$ is monotonically decreasing (increasing) on $(-\infty, \eta_*]$ ($[\eta_*, +\infty)$).

Thus $f(\eta)$ has a unique global minimizer $\alpha_* = f(\eta_*) > 0$. In addition, we have

$$\frac{\eta}{f(\eta)} = \frac{\eta(A_2 - A_4)}{A_0} = \frac{A_0 - 3A_2}{2A_0} \in \left(-1, \frac{1}{2}\right) \quad (\text{since } 0 < A_2 < A_0).$$

Therefore $f(\eta) > 2\eta$ and $f(\eta) > -\eta$, which implies

Conclusion 2: $f(\eta) \rightarrow +\infty$ as $\eta \rightarrow \pm\infty$.

Combining the above conclusions, we have that:

- For $\alpha > \alpha_*$, the equation $\alpha = f(\eta)$ has exactly two solutions $\eta_1 < \eta_* < \eta_2$; If $\alpha > (<)7.5$, we have $\eta_2 > (<)0$ (since $f(0) = 7.5$).
- For $\alpha = \alpha_*$, the equation $\alpha = f(\eta)$ has only one solution $\eta = \eta_*$;
- For $\alpha < \alpha_*$, the equation $\alpha = f(\eta)$ has no solution.

These yield the conclusions in Lemma 3.6 except for the justification of $\alpha_* > 20/3$.

Acknowledgments: We are very grateful to Professor John Ball for pointing out to us a lack of clarity in some parts of Step 2 and 5 in the original proof of Lemma 3.6. We also thank Wei Wang for his help on the second proof.

References

- [1] H. Liu, H. Zhang and P.W. Zhang. Axial symmetry and classification of stationary solutions of Doi-Onsager equation on the sphere with Maier-Saupe potential. *Comm. Math. Sci.*, 3(2):201–218, 2005.
- [2] W. Wang, P.W. Zhang and Z. F. Zhang. The small Deborah number limit of the Doi–Onsager equation to the EricksenLeslie equation *Comm. Pure Appl. Math.*, 68(8):1326–1398, 2015.