

# A Balance Between Euchromatic (*JIL-1*) and Heterochromatic [*SU(VAR)2-5* and *SU(VAR)3-9*] Factors Regulates Position-Effect Variegation in *Drosophila*

Chao Wang, Jack Girton, Jørgen Johansen, and Kristen M. Johansen<sup>1</sup>

Department of Biochemistry, Biophysics, and Molecular Biology, Iowa State University, Ames, Iowa 50011

**ABSTRACT** In this study, we show that the haplo-enhancer effect of *JIL-1* has the ability to counterbalance the haplo-suppressor effect of both *Su(var)3-9* and *Su(var)2-5* on position-effect variegation, providing evidence that a finely tuned balance between the levels of *JIL-1* and the major heterochromatin components contributes to the regulation of gene expression.

**T**HE essential *JIL-1* histone H3S10 kinase (Jin *et al.* 1999; Wang *et al.* 2001) is a major regulator of chromatin structure (Deng *et al.* 2005, 2008) that functions to maintain euchromatic domains while counteracting heterochromatinization and gene silencing (Ebert *et al.* 2004; Lerach *et al.* 2006; Zhang *et al.* 2006; Bao *et al.* 2007). In the absence of the *JIL-1* kinase, the major heterochromatin markers H3K9me<sub>2</sub>, HP1a [*Su(var)2-5*], and *Su(var)3-7* spread to ectopic locations on the chromosome arms (Zhang *et al.* 2006; Deng *et al.* 2007, 2010). These observations suggested a model for a dynamic balance between euchromatin and heterochromatin (Ebert *et al.* 2004; Zhang *et al.* 2006; Deng *et al.* 2010), where, as can be monitored in position-effect variegation (PEV) arrangements, the boundary between these two states is determined by antagonistic functions of a euchromatic regulator (*JIL-1*) and the major determinants of heterochromatin assembly, *e.g.*, *Su(var)3-9*, HP1a, and *Su(var)3-7* (for review see Weiler and Wakimoto 1995; Girton and Johansen 2008). In support of this model, Deng *et al.* (2010) recently showed that *Su(var)3-7* and *JIL-1* loss-of-function mutations have an antagonistic and counterbalancing effect on gene expression using PEV assays; however, potential dynamic interactions between *JIL-1* and the other two heterochromatin genes, *Su(var)3-9* and *Su(var)2-5*, were not

addressed in this study. Interestingly, in other genetic interaction assays monitoring the lethality as well as the chromosome morphology defects associated with the null *JIL-1* phenotype, only a reduction in the dose of the *Su(var)3-9* gene (Zhang *et al.* 2006; Deng *et al.* 2007) rescued both phenotypes. In contrast, in the same assays a reduction of *Su(var)3-7* rescued the lethality, but not the chromosome defects (Deng *et al.* 2010), and no genetic interactions were detectable between *JIL-1* and *Su(var)2-5* (Deng *et al.* 2007). Thus, these findings indicate that while *Su(var)3-9* activity may be a major factor in the lethality and chromatin-structure perturbations associated with loss of the *JIL-1* histone H3S10 kinase, these effects are likely to be uncoupled from HP1a and, to a lesser degree, from *Su(var)3-7*. This raises the question of whether *JIL-1* dynamically interacts with the two other heterochromatin genes, *Su(var)2-5* and *Su(var)3-9*, in regulating gene expression, as it does with *Su(var)3-7*.

To answer this question, we explored the effect of various combinations of loss-of-function alleles of *JIL-1* and *Su(var)3-9* or *Su(var)2-5* on PEV caused by the *P*-element insertion line *118E-10* (Wallrath and Elgin 1995; Wallrath *et al.* 1996). Insertion of this *P* element (*P*[*hsp26-pt, hsp70-w*]) into euchromatic sites results in a uniform red-eye phenotype whereas insertion into a known heterochromatin region of the fourth chromosome results in a variegating eye phenotype (Cryderman *et al.* 1998; Bao *et al.* 2007) (Figures 1 and 2). It has been demonstrated that loss-of-function *JIL-1* alleles can act as haplo-enhancers of PEV, resulting in increased silencing of gene expression (Deng *et al.* 2010), whereas loci for structural components of heterochromatin


Copyright © 2011 by the Genetics Society of America

doi: 10.1534/genetics.111.129353

Manuscript received January 13, 2011; accepted for publication April 11, 2011

Supporting information is available online at <http://www.genetics.org/cgi/content/full/genetics.111.129353/DC1>.

<sup>1</sup>Corresponding author: Department of Biochemistry, Biophysics, and Molecular Biology, 3154 Molecular Biology Bldg., Iowa State University, Ames, IA 50011. E-mail: kristen@iastate.edu




**Figure 1** Counterbalancing effect of *JIL-1* and *Su(var)3-9* loss-of-function alleles on the PEV of the *P*-element insertion line *118E-10*. (A) Examples of the degree of PEV in the eyes of wild-type *JIL-1* and *Su(var)3-9* (+/+), *JIL-1<sup>z2</sup>/+* (*z2/+*), *Su(var)3-9<sup>06</sup>/+* (*3-9<sup>06</sup>/+*), and *JIL-1<sup>z2</sup>/Su(var)3-9<sup>06</sup>* (*z2/3-9<sup>06</sup>*) flies in a *118E-10/+* background. All images are from male flies. (B) Histograms of the amount of eye pigment in +/+, *JIL-1<sup>z2</sup>/+* (*z2/+*), *Su(var)3-9<sup>06</sup>/+* (*3-9<sup>06</sup>/+*), and *JIL-1<sup>z2</sup>/Su(var)3-9<sup>06</sup>* (*z2/3-9<sup>06</sup>*) male flies heterozygous for *118E-10*. Fly stocks were maintained according to standard protocols (Roberts 1998). Oregon-R was used for wild-type preparations. The *JIL-1<sup>z2</sup>* allele is described in Wang *et al.* (2001) and in Zhang *et al.* (2003). The *Su(var)3-9<sup>06</sup>* and *Su(var)2-5<sup>05</sup>* alleles are described in Schotta *et al.* (2002) and in Eissenberg *et al.* (1992). The *P*-element *P[hsp26-pt, hsp70-w]* insertion line *118E-10* was the generous gift of L. Wallrath. The *hsp70* promoter is leaky and promotes sufficient expression to generate a variegated

eye phenotype under non-heat-shock conditions (Wallrath and Elgin 1995; Bao *et al.* 2007). PEV assays were performed as previously described in Lerach *et al.* (2006) and in Bao *et al.* (2007). In short, various combinations of *JIL-1*, *Su(var)3-9*, or *Su(var)2-5* alleles were introduced into the *118E-10* or *w<sup>md</sup>* PEV arrangements by standard crossing. To quantify the variegated phenotype, adult flies were collected from the respective crosses at eclosion, aged 6 days at 25°, frozen in liquid nitrogen, and stored at -80°C until assayed. The pigment assays were performed essentially as in Kavi and Birchler (2009) using three sets of 10 fly heads of each genotype collected from males and females. For each sample, the heads from the 10 flies were homogenized in 200  $\mu$ l of methanol with 0.1% hydrochloric acid and centrifuged, and the OD of the supernatant was spectrophotometrically measured at a wavelength of 480 nm.

such as *Su(var)3-9*, *Su(var)2-5*, and *Su(var)3-7* act as strong haplo-suppressors (Eissenberg *et al.* 1990; Reuter *et al.* 1990; Tschiersch *et al.* 1994). In the experiments, the transgenic reporter line *118E-10* was crossed into *JIL-1<sup>z2</sup>/+*, *Su(var)3-9<sup>06</sup>/+*, and *Su(var)2-5<sup>05</sup>/+* mutant backgrounds as well as into *JIL-1<sup>z2</sup>/Su(var)3-9<sup>06</sup>* and *JIL-1<sup>z2</sup>/Su(var)2-5<sup>05</sup>* double-mutant backgrounds. The *JIL-1<sup>z2</sup>* allele is a true null allele (Wang *et al.* 2001; Zhang *et al.* 2003), the loss-of-function *Su(var)3-9<sup>06</sup>* allele is due to a DNA insertion (Schotta *et al.* 2002), and the *Su(var)2-5<sup>05</sup>* loss-of-function allele is associated with a frameshift resulting in a nonsense peptide containing only the first 10 amino acids of HP1a (Eissenberg *et al.* 1992). Thus, to test whether the heterozygous *JIL-1<sup>z2</sup>* allele could counterbalance the suppression of the *Su(var)3-9<sup>06</sup>* or *Su(var)2-5<sup>05</sup>* loss-of-function alleles of the PEV of *118E-10*, we compared the eye pigment levels of the various genotypes (Figures 1 and 2 and Table 1). Pigment assays were performed essentially as in Kavi and Birchler (2009) using

three sets of 10 pooled fly heads from each genotype. Although both male and female flies were scored, due to sex differences only results from male flies are shown. However, the trend observed in female flies was identical to that in male flies (supporting information, Figure S1). As illustrated in Figures 1 and 2, the heterozygous *JIL-1<sup>z2</sup>/+* genotype enhances PEV as indicated by the increased proportion of white ommatidia and a 59% decrease in the optical density (OD) of the eye pigment levels ( $0.0083 \pm 0.0015$ ;  $n = 3$ ) as compared to +/+ flies ( $0.0203 \pm 0.0021$ ;  $n = 3$ ). This reduction was statistically significant (Table 1). In contrast, the heterozygous *Su(var)3-9<sup>06</sup>/+* and *Su(var)2-5<sup>05</sup>/+* genotypes suppress PEV as indicated by an increase of the proportion of red ommatidia and a statistically significant (Table 1) 384% ( $0.078 \pm 0.001$ ;  $n = 3$ ) and 330% ( $0.067 \pm 0.004$ ;  $n = 3$ ) increase, respectively, in the OD of the eye pigment levels. However, in the *JIL-1<sup>z2</sup>/Su(var)3-9<sup>06</sup>* and *JIL-1<sup>z2</sup>/Su(var)2-5<sup>05</sup>* double-mutant backgrounds, variegation



**Figure 2** Counterbalancing effect of *JIL-1* and *Su(var)2-5* loss-of-function alleles on the PEV of the *P*-element insertion line *118E-10*. (A) Examples of the degree of PEV in the eyes of wild-type *JIL-1* and *Su(var)2-5* (+/+), *JIL-1<sup>z2</sup>/+* (*z2/+*), *Su(var)2-5<sup>05</sup>/+* (*2-5<sup>05</sup>/+*), and *JIL-1<sup>z2</sup>/Su(var)2-5<sup>05</sup>* (*z2/2-5<sup>05</sup>*) flies in a *118E-10/+* background. All images are from male flies. (B) Histograms of the levels of eye pigment in +/+, *JIL-1<sup>z2</sup>/+* (*z2/+*), *Su(var)2-5<sup>05</sup>/+* (*2-5<sup>05</sup>/+*), and *JIL-1<sup>z2</sup>/Su(var)2-5<sup>05</sup>* (*z2/2-5<sup>05</sup>*) male flies heterozygous for *118E-10*.

**Table 1** Statistical comparison of eye pigment assays

Genotype <sup>a</sup>	<i>JIL-1<sup>z2</sup>/+</i>	<i>3-9<sup>06</sup>/+</i>	<i>JIL-1<sup>z2</sup>/3-9<sup>06</sup></i>
<i>118E-10/+</i>			
+/+	$P < 0.005$	$P < 0.0001$	$P > 0.8$
<i>JIL-1<sup>z2</sup>/+</i>	—	$P < 0.0001$	$P < 0.005$
<i>3-9<sup>06</sup>/+</i>	—	—	$P < 0.0001$
Genotype <sup>a</sup>	<i>JIL-1<sup>z2</sup>/+</i>	<i>2-5<sup>05</sup>/+</i>	<i>JIL-1<sup>z2</sup>/2-5<sup>05</sup></i>
<i>118E-10/+</i>			
+/+	$P < 0.005$	$P < 0.0001$	$P > 0.06$
<i>JIL-1<sup>z2</sup>/+</i>	—	$P < 0.0001$	$P < 0.005$
<i>2-5<sup>05</sup>/+</i>	—	—	$P < 0.0005$
Genotype	<i>JIL-1<sup>z2</sup>/+</i>	<i>3-9<sup>06</sup>/+</i>	<i>JIL-1<sup>z2</sup>/3-9<sup>06</sup></i>
<i>w<sup>m4</sup>/Y</i>			
+/+	$P < 0.005$	$P < 0.0001$	$P < 0.0001$
<i>JIL-1<sup>z2</sup>/+</i>	—	$P < 0.0001$	$P < 0.0001$
<i>3-9<sup>06</sup>/+</i>	—	—	$P < 0.001$
Genotype	<i>JIL-1<sup>z2</sup>/+</i>	<i>2-5<sup>05</sup>/+</i>	<i>JIL-1<sup>z2</sup>/2-5<sup>05</sup></i>
<i>w<sup>m4</sup>/Y</i>			
+/+	$P < 0.005$	$P < 0.0001$	$P < 0.0001$
<i>JIL-1<sup>z2</sup>/+</i>	—	$P < 0.0001$	$P < 0.0001$
<i>2-5<sup>05</sup>/+</i>	—	—	$P < 0.001$

For each genotype, the average pigment level from three sets of measurements from 10 pooled fly heads were compared using a two-tailed Student's *t*-test.


<sup>a</sup> Only male flies were scored.

of the proportion of red ommatidia was intermediate, and the eye pigment levels ( $0.0207 \pm 0.0031$  and  $0.0263 \pm 0.0035$ , respectively;  $n = 3$ ) were statistically indistinguishable from genotypes with +/+ levels of *JIL-1*, *Su(var)3-9*, and *Su(var)2-5* proteins (Figures 1 and 2 and Table 1).

To test whether a heterozygous *JIL-1* null allele also could counterbalance the suppression of the *Su(var)3-9<sup>06</sup>* or *Su(var)2-5<sup>05</sup>* alleles of the PEV of *w<sup>m4</sup>*, we performed experiments similar to those described above for *118E-10*. As illustrated in Figure 3 for male flies, the heterozygous *JIL-1<sup>z2</sup>/+* genotype enhances PEV as indicated by a 67% decrease in the OD of the eye pigment levels ( $0.008 \pm 0.002$ ;  $n = 3$ ) as compared to +/+ flies ( $0.024 \pm 0.003$ ,  $n = 3$ ). This reduction was statistically significant (Table 1).

In contrast, the heterozygous *Su(var)3-9<sup>06</sup>/+* and *Su(var)2-5<sup>05</sup>/+* genotypes suppress PEV as indicated by a statistically significant (Table 1) 1115% ( $0.2677 \pm 0.0061$ ;  $n = 3$ ) and 1023% ( $0.2454 \pm 0.0103$ ;  $n = 3$ ) increase, respectively, in the OD of the eye pigment levels. However, in the *JIL-1<sup>z2</sup>/Su(var)3-9<sup>06</sup>* and *JIL-1<sup>z2</sup>/Su(var)2-5<sup>05</sup>* double-mutant backgrounds, the eye pigment levels ( $0.2337 \pm 0.0011$  and  $0.2043 \pm 0.0037$ , respectively;  $n = 3$ ) were significantly reduced (Table 1) by 13% and 17% as compared to heterozygous *Su(var)3-9<sup>06</sup>/+* and *Su(var)2-5<sup>05</sup>/+* genotypes, indicating that a heterozygous *JIL-1* null allele has the ability to counterbalance the suppression of the *Su(var)3-9<sup>06</sup>* or *Su(var)2-5<sup>05</sup>* alleles of the PEV of *w<sup>m4</sup>*. However, it should be noted that it has been demonstrated that *JIL-1* can act both as an enhancer and as a suppressor of *w<sup>m4</sup>* PEV, depending on the precise levels of *JIL-1* (Lerach *et al.* 2006; Deng *et al.* 2010). Thus, the genetic interactions between *JIL-1* and the *Su(var)3-9* and *Su(var)2-5* alleles in regulating the PEV of *w<sup>m4</sup>* are likely to be more complex than in the case of *118E-10* where reduced levels of *JIL-1* always act as an enhancer (Bao *et al.* 2007). In females where the enhancer effect of the heterozygous *JIL-1<sup>z2</sup>* allele is less pronounced than in males, a statistically significant counterbalancing effect was detected only in flies of the *JIL-1<sup>z2</sup>/Su(var)2-5<sup>05</sup>* genotype (Figure S2).

These results demonstrate that the haplo-enhancer effect of *JIL-1* has the ability to counterbalance the haplo-suppressor effect of both *Su(var)3-9* and *Su(var)2-5* on the PEV of two different alleles. In previous experiments, a genetic interaction between *JIL-1* and *Su(var)2-5* was not detected (Deng *et al.* 2007). However, the assays used to probe for interactions were viability and rescue of polytene chromosome morphology. As indicated by the experiments presented here, these parameters are likely to be independent of and separate from the mechanisms contributing to epigenetic regulation of PEV and gene silencing. Consequently, the present experiments, taken together with those of Deng *et al.* (2010) using a *JIL-1* null allele, provide strong evidence that a finely tuned balance between the levels of *JIL-1* and all of the major heterochromatin components *Su(var)3-9*, *HP1a*, and



**Figure 3** Counterbalancing effect of *JIL-1* with *Su(var)3-9* and *Su(var)2-5* loss-of-function alleles on the PEV of *w<sup>m4</sup>*. (A) Histograms of the levels of eye pigment in +/+, *JIL-1<sup>z2</sup>/+* (*z2/+*), *Su(var)3-9<sup>06</sup>/+* (*3-9<sup>06</sup>/+*), and *JIL-1<sup>z2</sup>/Su(var)3-9<sup>06</sup>* (*z2/3-9<sup>06</sup>*) *w<sup>m4</sup>* male flies. (B) Histograms of the levels of eye pigment in +/+, *JIL-1<sup>z2</sup>/+* (*z2/+*), *Su(var)2-5<sup>05</sup>/+* (*2-5<sup>05</sup>/+*), and *JIL-1<sup>z2</sup>/Su(var)2-5<sup>05</sup>* (*z2/2-5<sup>05</sup>*) *w<sup>m4</sup>* male flies.

Su(var)3-7 contributes to the regulation of PEV and gene expression.

## Acknowledgments

We thank members of the laboratory for discussion, advice, and critical reading of the manuscript. We also acknowledge V. Lephart for maintenance of fly stocks and Kevin Bienik for technical assistance. We especially thank L. Wallrath for providing fly stocks. This work was supported by National Institutes of Health grant GM062916 (to K.M.J. and J.J.).

## Literature Cited

- Bao, X., H. Deng, J. Johansen, J. Girton, and K. M. Johansen, 2007 Loss-of-function alleles of the JIL-1 histone H3S10 kinase enhance position-effect variegation at pericentric sites in *Drosophila* heterochromatin. *Genetics* 176: 1355–1358.
- Cryderman, D. E., M. H. Cuaycong, S. C. R. Elgin, and L. L. Wallrath, 1998 Characterization of sequences associated with position-effect variegation at pericentric sites in *Drosophila* heterochromatin. *Chromosoma* 107: 277–285.
- Deng, H., W. Zhang, X. Bao, J. N. Martin, J. Girton *et al.*, 2005 The JIL-1 kinase regulates the structure of *Drosophila* polytene chromosomes. *Chromosoma* 114: 173–182.
- Deng, H., X. Bao, W. Zhang, J. Girton, J. Johansen *et al.*, 2007 Reduced levels of Su(var)3-9 but not Su(var)2-5 (HP1) counteract the effects on chromatin structure and viability in loss-of-function mutants of the JIL-1 histone H3S10 kinase. *Genetics* 177: 79–87.
- Deng, H., X. Bao, W. Cai, M. J. Blacketer, A. S. Belmont *et al.*, 2008 Ectopic histone H3S10 phosphorylation causes chromatin structure remodeling in *Drosophila*. *Development* 135: 699–705.
- Deng, H., W. Cai, C. Wang, S. Lerach, M. Delattre *et al.*, 2010 *JIL-1* and *Su(var)3-7* interact genetically and counteract each other's effect on position-effect variegation in *Drosophila*. *Genetics* 185: 1183–1192.
- Ebert, A., G. Schotta, S. Lein, S. Kubicek, V. Krauss *et al.*, 2004 *Su(var)* genes regulate the balance between euchromatin and heterochromatin in *Drosophila*. *Genes Dev.* 18: 2973–2983.
- Eissenberg, J. C., T. James, D. Foster-Hartnett, D. Hartnett, V. Ngan *et al.*, 1990 Mutation in a heterochromatin-specific chromosomal protein is associated with suppression of position-effect variegation in *Drosophila melanogaster*. *Proc. Natl. Acad. Sci. USA* 87: 9923–9927.
- Eissenberg, J. C., G. D. Morris, G. Reuter, and T. Hartnett, 1992 The heterochromatin-associated protein HP-1 is an essential protein in *Drosophila* with dosage-dependent effects on position-effect variegation. *Genetics* 131: 345–352.
- Girton, J., and K. M. Johansen, 2008 Chromatin structure and regulation of gene expression: the lessons of PEV in *Drosophila*. *Adv. Genet.* 61: 1–43.
- Jin, Y., Y. Wang, D. L. Walker, H. Dong, C. Conley *et al.*, 1999 JIL-1: a novel chromosomal tandem kinase implicated in transcriptional regulation in *Drosophila*. *Mol. Cell* 4: 129–135.
- Kavi, H. H., and J. A. Birchler, 2009 Interaction of RNA polymerase II and the small RNA machinery affects heterochromatic silencing in *Drosophila*. *Epigenetics Chromatin* 2: 15–30.
- Lerach, S., W. Zhang, X. Bao, H. Deng, J. Girton *et al.*, 2006 Loss-of-function alleles of the JIL-1 kinase are strong suppressors of position effect variegation of the *w<sup>m4</sup>* allele in *Drosophila*. *Genetics* 173: 2403–2406.
- Reuter, G., M. Giarre, J. Farah, J. Gausz, A. Spierer *et al.*, 1990 Dependence of position-effect variegation in *Drosophila* on dose of a gene encoding an unusual zinc-finger protein. *Nature* 344: 219–223.
- Roberts, D. B., 1998 *Drosophila: A Practical Approach*. IRL Press, Oxford.
- Schotta, G., A. Ebert, V. Krauss, A. Fischer, J. Hoffmann *et al.*, 2002 Central role of *Drosophila* SU(VAR)3-9 in histone H3-K9 methylation and heterochromatic gene silencing. *EMBO J.* 21: 1121–1131.
- Tschiersch, B., A. Hofmann, V. Krauss, R. Dorn, G. Korge *et al.*, 1994 The protein encoded by the *Drosophila* position-effect variegation suppressor gene *Su(var)3-9* combines domains of antagonistic regulators of homeotic gene complexes. *EMBO J.* 13: 3822–3831.
- Wallrath, L. L., and S. C. R. Elgin, 1995 Position effect variegation in *Drosophila* is associated with altered chromatin structure. *Genes Dev.* 9: 1263–1277.
- Wallrath, L. L., V. P. Guntur, L. E. Rosman, and S. C. R. Elgin, 1996 DNA representation of variegating heterochromatic P-element inserts in diploid and polytene tissues of *Drosophila melanogaster*. *Chromosoma* 104: 519–527.
- Wang, Y., W. Zhang, Y. Jin, J. Johansen, and K. M. Johansen, 2001 The JIL-1 tandem kinase mediates histone H3 phosphorylation and is required for maintenance of chromatin structure in *Drosophila*. *Cell* 105: 433–443.
- Weiler, K. S., and B. T. Wakimoto, 1995 Heterochromatin and gene expression in *Drosophila*. *Annu. Rev. Genet.* 29: 577–605.
- Zhang, W., Y. Jin, Y. Ji, J. Girton, J. Johansen *et al.*, 2003 Genetic and phenotypic analysis of alleles of the *Drosophila* chromosomal JIL-1 kinase reveals a functional requirement at multiple developmental stages. *Genetics* 165: 1341–1354.
- Zhang, W., H. Deng, X. Bao, S. Lerach, J. Girton *et al.*, 2006 The JIL-1 histone H3S10 kinase regulates dimethyl H3K9 modifications and heterochromatic spreading in *Drosophila*. *Development* 133: 229–235.

Communicating editor: J. Tamkun


# GENETICS

Supporting Information


<http://www.genetics.org/cgi/content/full/genetics.111.129353/DC1>

## **A Balance Between Euchromatic (JIL-1) and Heterochromatic [SU(VAR)2-5 and SU(VAR)3-9] Factors Regulates Position-Effect Variegation in *Drosophila***

Chao Wang, Jack Girton, Jørgen Johansen, and Kristen M. Johansen



**Figure S1** Counterbalancing effect of *JIL-1* with *Su(var)3-9* and *Su(var)2-5* loss-of-function alleles on PEV of *118E-10* in female flies. (A) Histograms of the amount of eye pigment in +/+, *JIL-1*<sup>z2</sup>/+ (*z2*/+), *Su(var)3-9*<sup>06</sup>/+ (*3-9*<sup>06</sup>/+), and *JIL-1*<sup>z2</sup>/*Su(var)3-9*<sup>06</sup> (*z2/3-9*<sup>06</sup>) female flies heterozygous for *118E-10*. (B) Histograms of the levels of eye pigment in +/+, *JIL-1*<sup>z2</sup>/+ (*z2*/+), *Su(var)2-5*<sup>05</sup>/+ (*2-5*<sup>05</sup>/+), and *JIL-1*<sup>z2</sup>/*Su(var)2-5*<sup>05</sup> (*z2/2-5*<sup>05</sup>) female flies heterozygous for *118E-10*. The average pigment level from three sets of measurements from 10 pooled flyheads were compared using a two-tailed Student's t-test.



**Figure S2** Counterbalancing effect of *JIL-1* with *Su(var)3-9* and *Su(var)2-5* loss-of-function alleles on PEV of  $w^{m4}$  in female flies. (A) Histograms of the amount of eye pigment in +/+, *JIL-1<sup>z2</sup>/+* (*z2/+*), *Su(var)3-9<sup>06</sup>/+* (*3-9<sup>06</sup>/+*), and *JIL-1<sup>z2</sup>/Su(var)3-9<sup>06</sup>* (*z2/3-9<sup>06</sup>*) female flies heterozygous for  $w^{m4}$ . (B) Histograms of the levels of eye pigment in +/+, *JIL-1<sup>z2</sup>/+* (*z2/+*), *Su(var)2-5<sup>05</sup>/+* (*2-5<sup>05</sup>/+*), and *JIL-1<sup>z2</sup>/Su(var)2-5<sup>05</sup>* (*z2/2-5<sup>05</sup>*) female flies heterozygous for  $w^{m4}$ . The average pigment level from three sets of measurements from 10 pooled flyheads were compared using a two-tailed Student's t-test.