

Chromatin Structure and the Regulation of Gene Expression: The Lessons of PEV in *Drosophila*

Jack R. Girton and Kristen M. Johansen

Department of Biochemistry, Biophysics, and Molecular Biology,
Iowa State University, Ames, Iowa 50011

- I. Introduction: Position Effect in *Drosophila*
- II. Historical Background of the PEV Phenotype
 - A. The PEV phenotype and heterochromatin
 - B. Heterochromatin and euchromatin
- III. Types of PEV
 - A. Chromosomal rearrangement PEV
 - B. Transposon insertion PEV
 - C. Pairing-dependent dominant PEV: *Trans*-inactivation
- IV. Genome Organization and PEV
 - A. Chromatin structure
 - B. Nuclear organization
- V. Concluding Remarks
 - Acknowledgments
 - References

ABSTRACT

Position-effect variegation (PEV) was discovered in 1930 in a study of X-ray-induced chromosomal rearrangements. Rearrangements that place euchromatic genes adjacent to a region of centromeric heterochromatin give a variegated phenotype that results from the inactivation of genes by heterochromatin spreading from the breakpoint. PEV can also result from P element insertions that place euchromatic genes into heterochromatic regions and rearrangements

that position euchromatic chromosomal regions into heterochromatic nuclear compartments. More than 75 years of studies of PEV have revealed that PEV is a complex phenomenon that results from fundamental differences in the structure and function of heterochromatin and euchromatin with respect to gene expression. Molecular analysis of PEV began with the discovery that PEV phenotypes are altered by suppressor and enhancer mutations of a large number of modifier genes whose products are structural components of heterochromatin, enzymes that modify heterochromatic proteins, or are nuclear structural components. Analysis of these gene products has led to our current understanding that formation of heterochromatin involves specific modifications of histones leading to the binding of particular sets of heterochromatic proteins, and that this process may be the mechanism for repressing gene expression in PEV. Other modifier genes produce products whose function is part of an active mechanism of generation of euchromatin that resists heterochromatization. Current studies of PEV are focusing on defining the complex patterns of modifier gene activity and the sequence of events that leads to the dynamic interplay between heterochromatin and euchromatin.

© 2008, Elsevier Inc.

I. INTRODUCTION: POSITION EFFECT IN *DROSOPHILA*

The discovery of position effects in *Drosophila* began with the efforts of many investigators to generate a coherent picture of the nature of genes and chromosomes and of the relationship between them. Between 1910 and 1930, many genes were discovered in *Drosophila* by the recovery of mutant alleles. Analysis of mutant phenotypes suggested that each gene had a specific function and linkage mapping indicated that each gene had a unique chromosomal location (its locus). Cytological studies of mitotic and salivary gland polytene chromosomes indicated that chromosomes contain two visibly different regions, heterochromatin and euchromatin. Heterochromatin consists of regions that remain condensed and densely stained during interphase, while euchromatin consists of regions that become diffused during interphase and form the visible, banded regions of the polytene chromosomes. Euchromatin contains high concentrations of genes and heterochromatin contains very few genes. Although many believed that differential regulation of gene expression was an underlying feature of eukaryotic development, these investigations did not provide information about how gene activity was regulated. Since chromosomal rearrangements, such as inversions or translocations, that changed gene location but did not eliminate genes often had no visible mutant phenotype, chromosome structure was not believed to have a role in regulating gene function. Chromosomes appeared to simply serve as

vehicles for gene segregation during cell division. Two major findings challenged this view and launched an intensive study of the role of chromosome structure in gene regulation that continues today.

The first finding came in Sturtevant's study of the *Bar* eye mutation (Sturtevant, 1925). *Bar* is a dominant, sex-linked mutation that reduces the number of facets in the compound eye (normally about 800). The *Bar* mutation is a duplication of the *Bar* locus (*B*). Normal females have two copies of the locus, one on each X chromosome (*B/B*), but *Bar*-mutant females have four copies, two on each chromosome (*BB/BB*). Sturtevant described a new mutation (called *ultraBar* or *double-Bar*) that caused an even greater reduction in the number of facets. The *ultraBar* mutation has three copies of the *Bar* locus on one chromosome (*BBB*) and homozygous *ultraBar* females have a total of six copies (*BBB/BBB*). This suggested that increasing the number of copies of the *Bar* gene causes a decrease in eye size. However, Sturtevant noted that homozygous *Bar* females (*BB/BB*) have an average of 68×12 facets per eye, while females heterozygous for *ultraBar* (*BBB/B*) have an average of only 45×42 facets. Since both genotypes contain four copies of the *Bar* locus, this difference could not be due to differences in the number of copies of the locus. Sturtevant concluded that the arrangement of the copies of the *Bar* gene in the chromosome had an influence on *Bar* gene expression. He named this influence "position effect." This suggested that chromosomes are not passive collections of genes, but that they contain an internal structure that has a role in the regulation of gene expression.

The second finding came in 1930, when Muller reported the discovery of a series of X-ray-induced mutations affecting the *white* gene. Previously discovered *white* mutations reduced the amount of eye pigment, changing eye color from the normal uniform red to a uniform orange, yellow, or white (Lindsley and Zimm, 1992; Morgan *et al.*, 1925). Muller's new mutations showed a "mottled" phenotype, with each eye having some white (mutant) and some red (normal) regions (Fig. 1.1). The size, shape, and location of the white regions varied from eye to eye. The mutations causing these mottled phenotypes were chromosomal rearrangements that changed the location of the *white* gene in the chromosome (Muller, 1930). This rearrangement-induced mottled phenotype was named "variegated position effect" or "V-type" to distinguish it from the "stable position effect" or "S-type" position effect of *Bar*. Muller's discovery of the phenomenon of position-effect variegation (PEV) launched an intense study of the role of chromosome structure in gene expression that has continued for more than 75 years. This interest was partly because of the intriguing novel variegated phenotypes and partly because studies of PEV appeared likely to give important information about how gene expression was controlled by chromosomal organization.

PEV in *Drosophila* presents an interesting example of the historical progression of genetic analysis. During the past 75 years, the discovery of new investigative techniques has invariably led to a multitude of PEV studies using




Figure 1.1. Position-effect variegation (PEV) of an inversion chromosome. The mottled pigmentation observed in the eye of an individual carrying the $In(1)w^{m4}$ inversion chromosome shows regions where the *white* gene has been silenced (white) and regions where the *white* gene is actively expressed (dark).

these techniques. The sheer number of studies and the volume of information accumulated about PEV present a formidable task for the reader. There have been a number of extensive reviews of the experimental literature about PEV (Farkas *et al.*, 2000; Huisenga *et al.*, 2006; Lewis, 1950; Schulze and Wallrath, 2007; Spofford, 1976; Spradling and Karpen, 1990; Weiler and Wakimoto, 1995; Zhimulev, 1998). It is not the purpose of this chapter to repeat these works. Our goal is to summarize what has been learned about certain key aspects of PEV that occur reliably and repeatedly and which are considered defining features of the PEV phenomenon. These reflect, in a manner that is not currently understood, fundamental characteristics of underlying molecular/biochemical processes of chromatin organization that affect gene regulation. We will attempt to connect these characteristics of PEV with the results of recent molecular studies of the formation and molecular structure of chromatin. These results reflect the current excitement in the field, as it appears that the long examination of PEV may be approaching the point of finally revealing the role that different forms of

chromatin structure play in the regulation of gene expression, and how that regulation occurs. To cover such a large topic, we will have to be selective in the experiments we discuss and in the original publications we cite.

II. HISTORICAL BACKGROUND OF THE PEV PHENOTYPE

In 1930, Muller described a series of mutants induced by X-ray treatment. Muller originally discovered the first such mutation as a dominant *Notch* wing mutation. Dominant *Notch* mutations produce nicks or “notches” in the wings of heterozygous ($N/+$) flies. Muller knew *Notch* mutations sometimes resulted from deletions of portions of the X chromosome, so he generated flies heterozygous for the new *Notch* and a chromosome containing a recessive mutant allele of the nearby *white* locus ($N (?)/+ w$), expecting to see either a white eye phenotype, if a deletion removing both *Notch* and *white* was present, or a red eye phenotype if it were not. However, what he observed fit neither of these expectations. “To the great surprise of the writer the *Notch*-winged offspring of this cross had neither white nor normal red eyes, nor even eyes of any uniform intermediate colour. They had mottled eyes, and exhibited various grades and sizes of lighter and darker areas” ([Muller, 1930](#)). Further study revealed that the *Notch*-mutant phenotype was also “mottled” and that individuals showing a stronger mutant eye phenotype (lighter eyes) also had a stronger *Notch* phenotype (larger wing notches). This mutation was a translocation that had attached the distal tip of the X chromosome (the normal location of the *Notch* and *white* genes) to the third chromosome. Muller discovered four additional “white-mottled” mutations that did not involve mutations of *Notch* in additional irradiation experiments and all of these also were rearrangements (translocations or inversions) that moved the region of the X chromosome containing the *white* locus. Studying these mutations for more than 50 generations convinced Muller that the white mottled was a permanent mutant phenotype. He concluded that the change in the chromosomal arrangement of the genes on the X chromosome was responsible for the appearance of what he called the “eversporting displacement” phenotype, although he admitted to having no good explanation for how this worked. Muller’s paper, with its striking color diagrams showing the white-mottled phenotype, made an enormous impression and stimulated a number of other investigators to begin studying the relationship between chromosomal rearrangements and “mottled” gene expression. Studies of several genes quickly demonstrated that the mottled phenotype was produced only by certain types of chromosomal rearrangements and that the PEV phenotypes given by all rearrangements had a number of consistent characteristics ([Demerec, 1941](#); [Dobzhansky, 1936](#); [Lewis, 1950](#); [Muller, 1932](#)).

A. The PEV phenotype and heterochromatin

The PEV phenotypic effects Muller and other early investigators recovered were all produced by chromosomal rearrangements with one breakpoint close to the variegating gene and the other in a heterochromatic region that brought the affected gene into close proximity to heterochromatin. Examples of such PEV-inducing rearrangements are shown in Fig. 1.2. For example, the N^{264-52} X chromosome contains an inversion with one breakpoint in the centric heterochromatin and the other between *rst* and *w*. This places normal alleles of the *bi*, *ec*, *dm*, *fa*, and *rst* genes close to the centric heterochromatin. This inversion

Key:

Region of PEV = Breakpoint that gives PEV = Heterochromatin =

Euchromatin = Centromere =




Figure 1.2. A diagram showing some of the X chromosome rearrangements discovered by Muller to give position-effect variegation (PEV). The relationship between the chromosomal breakpoint and several genes that are affected is shown. Adapted from Lewis (1950).

generates a variegating loss-of-function phenotype for each of these genes. Females that are heterozygous for the N^{264-52} chromosome and a normal chromosome containing a recessive loss-of-function mutant allele of any of these genes have a variegating mutant phenotype for that gene. The variegation results from the alleles of the genes on the rearranged chromosome being repressed in some cells, and active in other cells. Females heterozygous for the rearranged chromosome and a chromosome with normal alleles of these genes have a normal phenotype, indicating that the PEV effect is recessive and does not affect the alleles on the normal chromosome. Other rearrangements, such as N^{264-57} or w^{m4} , have breakpoints in different locations and have different genes that show PEV but the effect is the same. PEV is produced in translocations, such as w^{258-18} , that move euchromatic genes on one chromosome adjacent to a heterochromatic region of a different chromosome, or in rearrangements, such as N^{264-57} , that move a region of heterochromatin into a euchromatic location (Fig. 1.2). The requirement for PEV is clearly that a euchromatic gene be moved adjacent to a heterochromatic breakpoint (Spofford, 1976; Zhimulev, 1998). This suggested that PEV phenotypes result from heterochromatic chromatin structure “spreading” from the breakpoint into the adjacent euchromatin and inactivating nearby euchromatic genes. The variegated phenotype was proposed to result from the length of spreading along the chromosome being variable.

B. Heterochromatin and euchromatin

Because heterochromatin plays a central role in any discussion of PEV, it is worthwhile to briefly discuss some key features of heterochromatin and euchromatin. Heitz (1928) discovered that portions of the chromosomes remain highly condensed and heavily stained throughout the cell cycle. He named these regions “heterochromatin” to distinguish them from the regions showing variable staining and condensation, which he called “euchromatin.” The boundaries of heterochromatin as defined by such cytological analyses are not precise and may change in different tissues or with different analytical techniques. In *Drosophila*, heterochromatic regions are located near the centromeres (the pericentric heterochromatin), at the telomeres, interspersed throughout the fourth chromosome, throughout the Y chromosome, and at a number of defined “intercalary heterochromatic” sites scattered among the chromosome arms (Zhimulev, 1998; Zhimulev and Belyaeva, 2003). In most *Drosophila* somatic cells, 30–35% of the genome is heterochromatic. Heterochromatin makes up all of the Y chromosome, 40% of the X, 25% of chromosomes 2 and 3, and more than half of the fourth. These heterochromatic regions contain a high percentage of families of repetitive DNA sequences with specific highly and moderately repetitive sequences clustered at particular chromosomal locations. The arrangement of these clusters of sequences is conserved among populations of *Drosophila*.

melanogaster (Pimpinelli *et al.*, 1995; Weiler and Wakimoto, 1995). In *Drosophila*, the pericentric heterochromatin is divided into two visibly distinct categories, α -heterochromatin and β -heterochromatin. The α -heterochromatin is largely composed of highly repetitive satellite sequences and is located in and around the centromere. It coalesces into a highly condensed, uniform body in the center of the chromocenter in salivary gland nuclei. β -Heterochromatin is more diffuse and expands or “loops out” to form the bulk of the visible chromocenter. β -Chromatin more closely resembles euchromatin. It contains functioning genes interspersed with middle repeated sequences, and a significant percentage consists of sequences derived from transposons (Pimpinelli *et al.*, 1995). More than 40 functioning genes in heterochromatic regions have been identified by mutant phenotypes, including visible phenotypes, lethality, and sterility (Gatti and Pimpinelli, 1992; Weiler and Wakimoto, 1995). *Drosophila* chromosomes contain complex repeated sequences at the telomeres. These include copies of two retrotransposons (HeT-A and TART) at the end of the telomere that have the ability to repair or heal damaged telomeres (Pardue and DeBaryshe, 1999; Wallrath, 2000). Proximal to these sequences, *Drosophila* telomeres contain repeated sequences called telomere-associated sequences (Karpen and Spradling, 1992). These sequences differ in length and sequence from chromosome to chromosome (Walter *et al.*, 1995).

III. TYPES OF PEV

There are a variety of ways in which a mosaic phenotype can be produced in *Drosophila*, including somatic recombination, somatic chromosomal loss, and somatic P element transposition. In this chapter, we will consider as PEV only those variegated phenotypes that are produced by chromosomal rearrangements that share the common feature that they bring euchromatic genes into close association with heterochromatin without altering the gene itself. These can be grouped into three classes or types based on a number of characteristic features.

A. Chromosomal rearrangement PEV

The common feature of all chromosomal rearrangements that give PEV is that they either move a euchromatic gene into a pericentric region or move a block of heterochromatic sequences into a euchromatic region. Chromosomal rearrangements involving the X, the second, the third, and the fourth chromosomes give PEV phenotypes (Demerec, 1941; Kaufmann, 1946; Lewis, 1950; Weiler and Wakimoto, 1995; Zhitulev, 1998). The heterochromatic regions that can produce a PEV phenotype are the pericentric regions of all chromosomes, most of

the fourth chromosome, and all of the Y chromosome. Several experiments in which the variegating allele is removed from the rearranged chromosome show the PEV effect is a repression of gene expression and not a permanent change in the allele. [Dubinin and Sidorov \(1935\)](#) used meiotic recombination to remove the variegated allele of the *hairy* gene from a translocation between the third and fourth chromosome. A double crossover in flies heterozygous for this translocation and normal chromosomes moved the variegating allele to the normal chromosome. In the normal chromosome, the allele gave a normal, *hairy*⁺ phenotype. When this same allele was reinserted into the translocated chromosome by another double crossover event, the allele again showed a variegating phenotype. Similar experiments with other variegating genes gave the same results ([Lewis, 1950](#); [Spofford, 1976](#)). Inducing an additional chromosomal rearrangement has also been used to separate the variegating allele from the heterochromatin. [Panshin \(1938\)](#) used X-rays to induce a second translocation event that returned the portion of the X chromosome containing the variegating *white* allele in the *w^{m11}* translocation to a euchromatic chromosomal location where it again gave a normal phenotype. Several similar experiments demonstrated that variegating alleles of any gene separated from the heterochromatic breakpoint by another chromosomal rearrangement ceases to show the variegating phenotype ([Demerec, 1941](#); [Hinton, 1950](#); [Judd, 1955](#); [Lewis, 1950](#)). Another type of evidence that PEV is not the result of a change in gene structure came from studies of gene products using alleles that produce proteins with detectable structural differences. In one such experiment, [Bahn \(1971\)](#) studied individuals heterozygous for T(1;2)OR32 carrying the *Amy2* allele and a normal chromosome carrying the *Amy4,6* allele using gel electrophoresis to detect which electrophoretic variant proteins were produced. In variegating individuals, the level of the *Amy2* variant was reduced, and the level of reduction was greater when the strength of the PEV was increased, but the structure of the protein product was not altered. The level of the *Amy4,6* enzyme produced by the allele on the normal chromosome was not altered, indicating that the PEV effect acted only on the allele in the rearranged chromosome. In other similar studies, PEV was also found to reduce enzyme levels from alleles of *sc*, *dor*, and *pn* ([Gerasimova et al., 1972](#)).

In a study of PEV reversion, [Tartof et al. \(1984\)](#) induced a series of rearrangements of the *w^{m4}* chromosome. The *w^{m4}* chromosome has an inversion with one breakpoint about 25 kb from the *white* locus and the other in the centric heterochromatin and produces a white PEV phenotype. Tartof recovered three rearrangements that had a normal phenotype. Each had one breakpoint in the heterochromatin beyond the *w^{m4}* breakpoint and they all retained some heterochromatic sequences at the original breakpoint. This indicates that the breakpoint itself did not cause the PEV phenotype. [Reuter et al. \(1985\)](#) studied 51 reversions of the *w^{m4}* chromosome and found only 3 true revertants with no

detectable PEV phenotype. The others showed a reduced PEV phenotype, and each of these retained blocks of heterochromatin at the original breakpoint site. These studies indicate that PEV results from the influence of a significant region of heterochromatic sequence adjacent to euchromatic genes, and that it does not simply result from a breakpoint, or from the gene being moved close to a centromere. They also suggest that the PEV effect is correlated with the amount of heterochromatic sequences and that there is a lower limit on the size of the block of heterochromatin that can cause PEV.

1. The spreading effect

The strength of PEV (the frequency with which a gene is inactivated) is inversely proportional to distance from the breakpoint (Lewis, 1950). For example, as diagrammed in Fig. 1.2, chromosomes containing the inversion N^{264-52} give a variegated phenotype for five genes (*rst*, *fa*, *dm*, *ec*, *bi*). The strength of the PEV effect on each was measured and genes further from the breakpoint showed less frequent or less strong mutant phenotypes, while those closer to the breakpoint showed more frequent or stronger mutant phenotypes (Demerec, 1940, 1941). This type of observation led to the hypothesis that PEV phenotypes resulted from a physical “spreading” of heterochromatin from the breakpoint into the euchromatic regions of the chromosome (Schultz, 1939). The distance between *Notch* and *white* is 330 kb, and in the original mutation of Muller both are frequently inactivated. The *bi* locus inactivated in the N^{264-52} chromosome is located at least 50 salivary gland chromosome bands away from the breakpoint, a molecular distance of more than 1mb of DNA. The PEV effect of $T(1;4)w^{m258-21}$ can extend over 67 bands (Hartman-Goldstein, 1967), that of $Dp(1;F)R$ over 86 bands (Spofford, 1976; Weiler and Wakimoto, 1995). These observations indicate that the spreading effect is able to cover significant chromosomal distances.

Two early studies suggested that the heterochromatic spreading effect in PEV is linear, that is, does not jump over genes. Demerec and Slizynska (1937) studied the PEV given by the w^{258-18} chromosome of *roughest* (*rst*) and *white* (*w*), two genes whose phenotypes can be scored in the eye. The gene order on the chromosome is: breakpoint, *rst*, *w*. An examination of the PEV phenotype showed the eye facets had one of three phenotypes: smooth and red (rst^+, w^+), rough and red ($rst^V w^+$), or rough and white (rst^V, w^V). They observed no facets with a smooth and white (rst^+, w^V) phenotype and concluded that the variegating effect passed in a linear manner from the breakpoint along the chromosome and always inactivated *roughest* before reaching *white*. Similar results in a study by Schultz (1941) of the PEV phenotype given by w^{258-21} on *split* and *white* reached the same conclusion. In a recent study, Talbert and Henikoff (2000) studied the PEV phenotypes given by X chromosome inversions that had

the arrangement of breakpoint, *w*, and *rst*. They reported cases of *w⁺* cells in a rough (*rst^V*) region of the eye. Does this mean that inactivation has “skipped over” *w* to inactivate *rst*? These results must be interpreted with caution. [Talbert and Henikoff \(2000\)](#) themselves noted that the alleles of *rst* and *w* used in their studies are not necessarily affecting the same cells. The *white* mutation is cell autonomous and its phenotype can be seen in each pigment cell in the eye. However, alleles of *roughest* affect the shape of the facet by causing the inclusion of extra cells in the eye facet. Not all cells in the facet need to be *rst⁻* to give a rough facet phenotype. PEV effects of *w^{m4}* and other rearrangements on *white* give a fine-scale variegation and individual facets often contain both white and pigmented cells (Girton, personal observation). It is thus possible that a “rough” facet may be mosaic and that the pigmented cells are actually *w⁺ rst⁺*, and the inactivation did not jump over *w* to inactivate *rst*. A definitive study of this question would need to use markers whose relationship to the breakpoint is known and whose expression can unambiguously be measured in the same cell.

2. All genes may show PEV

The first studies of PEV involved genes whose mutation produced a visible phenotype (such as *w*, *fa*, and *rst*). Later studies showed that PEV can be detected for a wide range of genes including those whose mutation produces a quantitative phenotype such as the number of bristles (*hairy*), the level of a detectable enzyme product (tryptophan pyrolase in *v⁺*, [Tobler et al., 1971](#)), enzymatic variants, (*Amy* and *Pdg*, [Bahn, 1971](#)), the amount of kynurenine production in fat body in *T(1;2)ras^v* and *T(1;Y)y⁺ Yv⁺* ([Rizki, 1961](#); [Tobler et al., 1971](#)), or the rate of ribosomal RNA synthesis in female ovaries (*sc^{S1}* and *sc^{L8}*, [Puckett and Snyder, 1973](#)). [Lewis \(1950\)](#) listed 32 genes shown to variegate. [Spofford \(1976\)](#) concluded that all genes can show a PEV-variegated phenotype in the right rearrangement.

Some genes are normally located in heterochromatic regions of chromosomes. Estimates of how many such genes exist in *Drosophila* range from 40 to 100 or more ([Schulze and Wallrath, 2007](#); [Spofford, 1976](#); [Weiler and Wakimoto, 1995](#)). A question that was raised very early in the study of PEV was how a gene whose normal location is heterochromatic would respond in a rearrangement that moved it to a euchromatic location. The first such gene studied was the light gene (*lt*) studied by [Schultz and Dobzhansky \(1934\)](#). The *lt* maps to a region considered to be β heterochromatin in chromosome 2 ([Hilliker, 1976](#); [Schultz and Dobzhansky, 1934](#)). Rearrangements that move *lt* adjacent to euchromatic regions give a variegated phenotype ([Hessler, 1958](#); [Schultz, 1936](#); [Schultz and Dobzhansky, 1934](#)). The conditions under which *lt* PEV inactivation occurs are the opposite of the conditions under which PEV inactivation of euchromatic genes occurs. At least five other heterochromatic genes normally

located near *lt* show similar position effects when placed next to euchromatin (Wakimoto and Hearn, 1990; Weiler and Wakimoto, 1995). Other heterochromatic genes that show this pattern include *cubitus interruptus* (*ci*) (Dubinin and Sidorov, 1935), *peach* (*pe*) in *Drosophila virilis* (Baker, 1953), male fertility factors on the Y chromosome (Hackstein and Hochstenbach, 1995; Neuhaus, 1939), and the nucleolus organizer region (Baker, 1971). Not all euchromatic regions show the same efficiency in generating heterochromatic PEV. All regions of X chromosome euchromatin can cause heterochromatic gene variegation, but only the distal parts of the arms of chromosome 2 and 3. Breakpoints in the fourth do not induce PEV of heterochromatic genes, perhaps reflecting the fact that the fourth chromosome contains interspersed heterochromatic and euchromatic regions (Spofford, 1976). In two cases, heterochromatic genes showing variegation when removed from surrounding heterochromatin had this variegation eliminated when an additional rearrangement moved them back near a block of heterochromatin (Eberl *et al.*, 1993). This is one of the most intriguing observations of PEV, that genes normally located in heterochromatin can be inactivated by being placed adjacent to euchromatin. Genes normally located in heterochromatic regions have some differences in molecular structure (Reugels *et al.*, 2000; Risinger *et al.*, 1997; Schulze and Wallrath, 2007; Schulze *et al.*, 2005). These observations suggest heterochromatic genes have evolved to not only tolerate being in a heterochromatic region but to actually require it for normal action.

3. Modifiers of the PEV phenotype

PEV induced by chromosomal rearrangements is strongly affected by *trans*-acting modifiers that increase the severity of the mutant phenotype (enhance variegation) or decrease the severity of the mutant phenotype (suppress variegation). Modifiers of PEV fall into two general classes, large-scale effects and single mutations. One large-scale effect is temperature. Rearing flies at higher temperature (29 °C) suppresses variegation (reduces the severity of the mutant phenotype) and rearing at lower temperature (16–18 °C) enhances variegation (increases the severity of the mutant phenotype). Gowen and Gay (1934) studied the phenotypic effects of temperature on PEV of *w*, *spl*, and *ec* caused by *w^{m1}*, *w^{m2}*, and *w^{m3}*. All of these genes showed more extreme mutant phenotypes at lower temperature. Kaufmann (1942) documented similar high- and low-temperature effects on PEV of *rst³*. All other euchromatic genes tested showed similar results, indicating this is a general effect of PEV (Bahn, 1971; Benner, 1971; Schalet, 1969; Spofford, 1976; Wargent, 1972).

PEV phenotypes are modified by changes in the amount of heterochromatic sequences in the genome. This was first observed in studies in which adding an extra copy of the Y chromosome was found to suppress variegation

and eliminating the Y chromosome enhanced variegation (Gowen and Gay, 1934). A PEV phenotype is more severely mutant in XO versus XY males or in XX versus XXY females (Lewis, 1950; Spofford, 1976). This effect is seen in genes producing visible phenotypes and those examined by measuring enzyme phenotypes (Bahn, 1971; Gerasimova *et al.*, 1972). In extreme cases, a rearrangement may not show a variegation phenotype except when enhancing modifiers are present. One example studied by Gersh (1963) involved the inversion *In(1)rst³*, which gives a white variegated phenotype in XO males but not in XY males. Changes in the Y chromosome affect all PEV-inducing rearrangements (Lindsley *et al.*, 1960; Weiler and Wakimoto, 1995; Zhimulev, 1998). Changes in the amount of centric heterochromatin also modify PEV phenotypes in the same manner as the Y chromosome. The deletion of a significant portion of the centric heterochromatic region of the X enhanced PEV of the *y*, *w*, and *ac* loci (Panshin, 1938). The addition or subtraction of an entire X has about the same effect as the addition or subtraction of the Y chromosome (Grell, 1958; Hinton, 1949). Similar effects have been discovered for the centric heterochromatic regions of the autosomes. A second chromosome deficiency that removes heterochromatin from the right arm of the second chromosome, *Df(2R)M-S2¹⁰*, enhances the PEV of several different genes in several different rearrangements (Morgan *et al.*, 1941). Duplications of the heterochromatic region of the right arm of the second chromosome suppress PEV phenotypes (Grell, 1970; Hannah, 1951). The fact that centric heterochromatin and the Y chromosome have similar effects on all euchromatic genes showing PEV indicates that this is a general effect of heterochromatin, and not due to specific sequences on the Y.

4. Single gene *Su(var)* and *E(var)* mutations

In several of the early studies of PEV, the severity of the phenotypes given by a PEV-inducing rearrangement varied from strain to strain, even when the strains did not differ for any obvious modifying factor (Hinton, 1949). A rearrangement, such as *In(1)w^{m4}*, that gives a PEV phenotype of 50% white eye in one strain could give phenotypes of 25% white eye or 75% white eye in other strains. These “genetic background effects” were believed to be due to unmapped dominant, single gene suppressor or enhancer mutations located throughout the genome (Schultz, 1941; Spofford, 1967, 1976). The first such modifier mutation to be isolated and characterized was a dominant suppressor mutation, *Su(var)*, on the third chromosome (Spofford, 1967). Individuals with an *In(1)w^{m4}* PEV-inducing genotype that also had the *Su(var)* allele had almost completely red eyes, while siblings with the *Su(var)⁺* allele had light, variegated eyes, the standard PEV phenotype. The *Su(var)* allele had similar effects on the PEV phenotypes of *w*, *rst*, and *dm* given by the *In(1)w^{m4}*, *Dp(1;3)N²⁶⁴⁻⁵⁸*, and *In(1)rst³* chromosomes

(Spofford, 1967, 1973, 1976). The discovery of the *Su(var)* allele suggested that there might be a number of genes whose mutation could modify PEV phenotypes. This led to several experiments to recover suppressor and enhancer mutations (Dorn *et al.*, 1993; Eissenberg *et al.*, 1990, 1992; Grigliatti, 1991; Locke *et al.*, 1988; Reuter and Wolff, 1981; Sinclair *et al.*, 1983, 2000; Wustmann *et al.*, 1989). A large number of dominant and recessive modifier alleles have been isolated, and it is estimated that there are between 50 and 150 loci that can mutate to give a PEV-modifying phenotype (Weiler and Wakimoto, 1995; Schulze and Wallrath, 2007). The effects of different modifier mutations appear to be similar; all suppressors reduce the severity of the PEV phenotype given by all rearrangements, and all enhancers increase the severity of the PEV phenotype given by all rearrangements (Fig. 1.3).

The PEV phenotype of genes that normally reside in heterochromatic regions shows a characteristic inverse response to modifiers of PEV. Temperature effects on the PEV of the heterochromatic gene *lt* are the reverse of those on euchromatic gene PEV, with higher temperatures producing a more mutant phenotype (Gersh, 1949). While extra Y chromosomes suppress PEV of euchromatic genes, they act as enhancers of variegation of *lt* (Baker and Reim, 1962; Schultz, 1936) and *ci* (Altorfer, 1967). Baker and Spofford (1959) studied the effects of 14 free Y chromosome fragments and reported that the same regions of




Figure 1.3. The effect of *su(var)* and *e(var)* mutations on PEV phenotypes. All individuals are homozygous for the pericentric P insertion line P118E-10 that gives a *white* PEV phenotype (Wallrath and Elgin, 1995). (A) The eye phenotype of an individual with no modifier mutation is variegated, with the *white* marker gene actively expressed in some cells but silenced in others. (B) An individual with a strong *e(var)* genotype, $JIL-1^{z2}/JIL-1^{z60}$, shows complete silencing of *white* gene expression. (C) An individual homozygous for a strong *su(var)* genotype, $Su(var)3-1/Su(var)3-1$, shows high levels of *white* gene expression. Photos: X. Bao.

the Y that suppress the *w* PEV phenotype given by *Dp(1;3)N²⁶⁴⁻⁵⁸* enhanced the *lt* PEV phenotype given by *T(2;3)ltm¹⁰⁰* (Baker and Reim, 1962). Single-gene mutations that suppress euchromatic gene PEV enhance the PEV phenotype of heterochromatic genes, and single-gene mutations that enhance euchromatic gene PEV suppress the PEV phenotype of heterochromatic genes (Schulze and Wallrath, 2007; Spofford, 1976; Weiler and Wakimoto, 1995). The reason why heterochromatic gene PEV and euchromatic gene PEV should show such consistent and opposite reactions is one of the most intriguing characteristics of PEV.

A paradigm in genetics is that the normal function of a gene can be deduced from the phenotype given by mutant alleles. The observation that mutations of modifier genes alter the PEV phenotype, and that heterochromatin is involved in generating PEV phenotypes, suggested that the normal functions of modifier genes are part of a genetic system that is essential for the establishment, maintenance, or function of chromatin structure. This is supported by several observations. The products of most modifier genes that have been studied to date are either structural components of heterochromatin (Baksa et al., 1993; Henikoff, 1979; James and Elgin, 1986; Locke et al., 1988; Reuter and Spierer, 1992; Reuter et al., 1982; Sinclair et al., 1983), enzymes that modify histones or nonhistone chromatin proteins (Ebert et al., 2006; Lerach et al., 2006; Mottes et al., 2000; Nishioka et al., 2002; Schotta et al., 2004), or nuclear architectural proteins (Bao et al., 2007a; Yamaguchi et al., 2001). For example, the product of the *Su(var)205* gene is the HP-1 protein, a heterochromatic protein that is found in high concentrations in the chromocenter in salivary gland nuclei (Eissenberg et al., 1990, 1992; James and Elgin, 1986). The product of the *Su(var)3-7* gene is a large zinc-finger protein that is mainly associated with the pericentric heterochromatin (Reuter et al., 1990). The JIL-1 protein is a tandem kinase that phosphorylates the serine 10 residue in the tail of histone 3 (Jin et al., 1999; Lerach et al., 2006; Zhang et al., 2003) and a dominant gain of function allele of JIL-1 is one of the strongest suppressors of PEV so far described (Ebert et al., 2004). The majority are haplo-abnormal, showing the dominant modifying effect when the gene is present in only single copy, and some (10%) have both haplo- and triplo-abnormal effects, modifying PEV when present in either one or three copies. For example, a deletion of *Su(var)205* suppresses the *In(1)w^{m4}* PEV phenotype, giving a pure red eye, and a duplication giving a genome three copies of the locus enhances the PEV phenotype, giving a nearly pure white eye (Schulze and Wallrath, 2007; Weiler and Wakimoto, 1995). This sensitivity to dosage suggests that a balance in the amount of the products of these genes is important for chromatin structure formation and function.

B. Transposon insertion PEV

Rearrangements that break and rejoin chromosomes are not the only method for placing a euchromatic gene adjacent to heterochromatin. In 1982, a genetically recombinant P element with an inserted cloned reporter gene (*ry*⁺) was induced to transpose into a chromosomal site (Rubin and Spradling, 1982; Spradling and Rubin, 1982). The reporter gene in P element transformation is essentially a small piece of euchromatic sequence that is inserted into different regions of the genome. In an early transformation experiment (Hazelrigg *et al.*, 1984), two insertions of a transposon containing as a reporter a wild-type copy of the *white* gene were recovered that gave a white variegated eye phenotype. There was no indication that the inserted *w*⁺ gene was altered in the transposition process and it was concluded that the variegation was the result of a position effect. One of these variegating insertions was located in the basal region of chromosome 2L, a heterochromatic region that was known to induce PEV in chromosomal rearrangements. The second was inserted near the telomere of the right arm of chromosome 3. This finding that genes inserted in a heterochromatic region by P element transformation can show a variegated phenotype was soon confirmed by other investigators (Steller and Pirrotta, 1985). In a genome-wide screen, Wallrath and Elgin (1995) recovered insertions with a white variegated phenotype in all of the heterochromatic regions known to induce PEV in rearrangements (the pericentric regions of the X, the second, the third, throughout the fourth, and in the Y chromosome). Variegating insertions were also recovered in or near the telomeres of all chromosomes (Wallrath and Elgin, 1995). Ahmad and Golic (1996) also recovered variegating P insertions in euchromatic regions at a site of intercalary heterochromatin. These results suggested that there was more to learn about position effects than had been suspected before, because P insertions can give a variegated phenotype in locations that do not give a PEV phenotype in chromosomal rearrangements (Fig. 1.4).

An important question about variegation produced by P element insertion is whether it is generated by the same mechanism as chromosomal rearrangement-induced PEV. One test of this is to determine the effect of known modifiers of PEV on the transposon insertion phenotype. Talbert *et al.* (1994) observed that the PEV phenotype of a transposon insertion in the telomeric region of 3R was not altered by modifiers of the PEV effect given by *w*^{m4}. Wallrath and Elgin (1995) showed the known PEV-modifying mutations, *Su(var)2-5*⁰² and *Su-var(2)1*⁰¹, suppressed the PEV phenotypes of all pericentric and fourth chromosome insertions, including insertions near the telomere of the fourth chromosome, but did not suppress the variegation produced by six different insertions located near the telomeres of 2R, 2L, or 3R. Comparing flies with XO and XXY genotypes showed that increasing the amount of heterochromatin suppressed the variegation phenotype of 13 inserts in the pericentric




Figure 1.4. A diagram showing the chromosomal regions in which P element insertions showing a position-effect variegation (PEV) phenotype were recovered by [Wallrath and Elgin \(1995\)](#). Insertions were primarily recovered at centromeric and telomeric regions and distributed throughout the fourth chromosome.

heterochromatin or in the fourth chromosome, but had no effect on 7 insertions in or near the telomeres of 2R, 3L, or 3R ([Wallrath and Elgin, 1995](#)). A second test of PEV is to determine whether variegation is eliminated when the variegated allele is removed from its location. [Ahmad and Golic \(1996\)](#) demonstrated that if an inserted P element that is showing variegation in somatic cells is physically removed from the chromosome to an extrachromosomal circle using the flippase (FLP) recombinase system, the gene expression is restored. All of the insert lines they tested responded to modifiers of PEV (enhancement in XO males and suppression by the *Su(var)205⁵⁰²* mutation), but none of their insertions were located at a telomere. These results indicate that PEV generated by P

element insertions in nontelomeric locations behaves like chromosomal rearrangement PEV, but that inserts in the telomeric regions have different properties. These results suggest that there may be differences in the mechanisms that produce telomeric insertion PEV phenotypes.

C. Pairing-dependent dominant PEV: *Trans*-inactivation

The first PEV-inducing rearrangements recovered gave mostly recessive phenotypes. For example, an individual heterozygous for the *In(1)w^{m4}* chromosome and a normal chromosome (*In(1)w^{m4}/w⁺*) has a wild-type phenotype because the breakpoint in the *In(1)w^{m4}* chromosome has no *trans*-inactivation effect on the *w⁺* allele in the normal chromosome. Chromosomal rearrangements that give a dominant PEV phenotype are rare. One well-studied example consists of rearrangements that affect the *brown* (*bw*) gene. Several inversions with one breakpoint in heterochromatin and another in the 59C6–59F3 interval of chromosome 2R have been recovered that give a dominant variegating brown eye PEV phenotype (Schultz, 1936; Slatis, 1955). Flies that are heterozygous for such a rearrangement and a normal chromosome containing a wild-type allele of the *brown* gene have a variegating brown eye color phenotype. This indicates that the breakpoint has a *trans*-inactivation effect on the *bw⁺* allele in the normal chromosome. A dominant *brown* allele (*bw^D*) was discovered that is not associated with an inversion but which carries an insertion of a block of 1.6 mb of centromeric heterochromatin in the *bw* locus (Hinton, 1940, 1942; Hinton and Goodsmith, 1950; Slatis, 1955). Flies with a *bw^D/bw⁺* genotype have a *brown* PEV phenotype. Removal of the heterochromatic block by further rearrangement eliminates the PEV effect (Hinton and Goodsmith, 1950). The dominant *bw^D* PEV phenotype is altered by suppressors of euchromatic gene PEV (Csink and Henikoff, 1996). These findings confirmed that the dominant *trans*-inactivation effect of *bw^D* is due to the presence of adjacent heterochromatin, and that the effect is a PEV effect.

Northern analysis indicates that *bw^D/bw⁺* individuals produce no detectable mRNA from either allele (Dreesen *et al.*, 1988; Henikoff and Dreesen, 1989). Thus, the dominant *brown* effect results from an inactivation of transcription of *brown* alleles in both chromosomes. The obvious question is how the insertion of a block of heterochromatin in one chromosome can affect the transcription of the allele on the other homologue. The key to this *trans*-inactivation effect is homologous chromosome pairing (Sass and Henikoff, 1999). When a fly has two *brown* variegating rearrangements with different breakpoints that disrupt somatic chromosome pairing, the PEV phenotype is also reduced (Henikoff and Dreesen, 1989). Confirmation that pairing is the key to *brown*-dominant *trans*-inactivation came when *bw^D* was combined with duplications

containing extra copies of the bw^+ in other chromosomal locations, where they do not pair with the normal locus. In such cases, the inserted bw^+ is not inactivated by bw^D (Dreesen *et al.*, 1991).

The idea that somatic pairing of chromosomes is important for normal function and might be a factor in PEV is an old one (Ephrussi and Sutton, 1944; Henikoff, 1994). How might pairing with a chromosome containing bw^D cause *trans*-inactivation of the bw^+ allele on the non-rearranged chromosome? The inserted heterochromatic block in bw^D contains multiple copies of the AAGAG sequence, a sequence that is one of the most abundant in the genome and which is found on each chromosome, and which is especially common in the pericentric region of chromosome 2 (Lohe *et al.*, 1993). Tracking the nuclear location of the bw insert using fluorescent *in situ* hybridization (FISH) shows that in the interphase nucleus, this block becomes associated with a region of the nucleus where the 2R-centric heterochromatin is localized (Csink *et al.*, 2002; Dernburg *et al.* 1996). The silencing of bw^D is thus correlated with the localization of the bw^D allele and the paired bw^+ allele to a heterochromatic region of the nucleus. This suggested that the dominant bw^D PEV effect is due to the bw^D allele dragging the paired bw^+ allele on the other homologue to a nuclear compartment where heterochromatic pairing results in gene inactivation. In a further investigation, the expressions of a series of P element insertions at different locations along the normal 2R chromosome were assayed when paired with the bw^D chromosome to determine the chromosomal extent of this *trans*-inactivation effect. An insertion 45 kb from the bw^D insertion was inactivated but inserts 280 and 460 kb from bw were not inactivated (Csink *et al.*, 2002). When the bw^D insert was introgressed into *Drosophila simulans*, which has AAGAG sequences in the centric region of the X chromosome but not the second chromosome, the insert continued to associate with the centric region of the second chromosome, suggesting that the heterochromatic association effect is not sequence specific but is an association with the nearest large block of heterochromatin (Sage and Csink, 2003). Recent studies suggest that this heterochromatic-pairing *brown*-dominant PEV may not be unique, and that heterochromatic insertions in other euchromatic regions may show *trans*-association/inactivation (Sage and Csink, 2003; Thakar and Csink, 2005). This suggests that *trans*-inactivation by heterochromatic insertion may be an important mechanism of gene silencing.

IV. GENOME ORGANIZATION AND PEV

The study of PEV has been instrumental in advancing our appreciation of the role that chromatin structure and genome organization play on the regulation of gene expression. In particular, the ability to isolate modifiers of the PEV-silencing effect in genetic screens resulted in identification of a large number of

chromosomal and nuclear architectural factors that are important for long-range control of gene expression ([Table 1.1](#)). Analysis of these factors has allowed us to develop a molecular understanding of the different processes contributing to the establishment or maintenance of active versus silent chromosomal domains and especially it has illuminated the critical role that heterochromatin plays in genome organization. In this section, we will review key aspects of chromatin structure and will examine some of the genes that affect PEV penetrance and the molecular mechanisms underlying the phenomenon. We will consider recent studies that underscore the link between heterochromatin formation and nuclear organization.

A. Chromatin structure

The primary packaging unit of chromatin is the nucleosome, composed of 147 bp of DNA wrapped around a histone octamer composed of two subunits each of the core histones H2A, H2B, H3, and H4 (reviewed in [Luger, 2006](#)). The histone globular domains are arranged in the interior of the nucleosome, while their unstructured tail domains project outwards where they are targets for a range of posttranslational modifications, including acetylation, phosphorylation, methylation, ADP-ribosylation, ubiquitination, sumoylation, and biotinylation. These modifications have been proposed to regulate genomic function by influencing chromatin structure both by altering the biophysical contacts between DNA and histones and by providing specific binding sites for different classes of chromatin-binding proteins (reviewed in [Kouzarides, 2007](#)). Thus, the modified histone tails provide a signaling platform that integrates output from various signal transduction pathways ultimately specifying the level of higher order chromatin folding ([Cheung et al., 2000; Fischle et al., 2003; Turner et al., 1992](#)).

Hyperacetylation of histones H3 and H4 and methylation of H3K4 are correlated with the establishment of transcriptionally active chromatin across a wide range of species from yeast to *Drosophila* to human ([Bernstein et al., 2005; Pokholok et al., 2005; Schubeler et al., 2004](#)). Acetylation neutralizes the basic charge of the lysine residue, which is thought to reduce histone tail interaction with the acidic phosphate residues of DNA, making it more accessible for transcription ([Wade et al., 1997](#)), and acetylation of the H4K16 residue, in particular, has been shown to physically impede the formation of higher order folded structures as well as block nucleosome sliding by the *Drosophila* ATP-utilizing chromatin assembly and remodeling factor (ACF) chromatin-remodeling complex *in vitro* ([Shogren-Knaak et al., 2006](#)). But besides changing the biophysical properties of chromatin fiber behavior, lysine acetylation also generates specific docking sites for distinct bromodomain proteins that have been found in a variety of different transcriptional activators and chromatin-remodeling complexes; binding to these sites can be influenced

Table 1.1. Nuclear PEV Modifiers of w^{m^4}

Locus	Function	References
Histone-modifying enzymes/histone variants		
JIL-1	H3S10 kinase	Wang <i>et al.</i> , 2001; Ebert <i>et al.</i> , 2004; Bao <i>et al.</i> , 2007a
Su(var)3-9	H3K9 methyltransferase	Reuter <i>et al.</i> , 1986; Schotta <i>et al.</i> , 2003b
G9a	H3K9 methyltransferase	Mis <i>et al.</i> , 2006
HDAC1/RPD3	Histone deacetylase	De Rubertis <i>et al.</i> , 1996; Mottus <i>et al.</i> , 2000
Chameau	Histone acetyltransferase (HAT)	Grienberger <i>et al.</i> , 2002
Reptin	HAT complex	Qi <i>et al.</i> , 2006
H2Av	H2A variant	Swaminathan <i>et al.</i> , 2005
Chromosomal proteins		
Su(var)2-5/HP1	H3K9me-binding	James and Elgin, 1986; Eissenberg <i>et al.</i> , 1990
HP2	AT Hook protein	Shaffer <i>et al.</i> , 2002
D1	AT Hook protein	Aulner <i>et al.</i> , 2002
Su(var)3-7	DNA binding (Zn finger)	Reuter <i>et al.</i> , 1990; Cléard and Spierer, 2001
Modulo	DNA binding	Perrin <i>et al.</i> , 1998; Büchner <i>et al.</i> , 2000
trl/GAGA factor	DNA binding	Dorn <i>et al.</i> , 1993; Farkas <i>et al.</i> , 1994
Bonus	Nuclear receptor cofactor	Beckstead <i>et al.</i> , 2005
mus209	Proliferating cell nuclear antigen (PCNA); DNA replication and repair	Henderson <i>et al.</i> , 1994
Cramped	Pc-G gene; DNA replication	Yamamoto <i>et al.</i> , 1997
Zeste	Transvection	Hazelrigg and Petersen, 1992
E(var)3-95E	E2F transcription factor; cell cycle	Seum <i>et al.</i> , 1996
ORC2	Replication; complexes with HP1	Pak <i>et al.</i> , 1997
HOAP	Complex with HP1/ORC; HMG-like	Shareef <i>et al.</i> , 2001
DRE4/spt16	Facilitates chromatin transcription (FACT) complex	Nakayama <i>et al.</i> , 2007
Nuclear and chromosome architecture		
Lamin Dm0	Nuclear lamina/architecture	Bao <i>et al.</i> , 2007b
Su(var)2-10/PIAS	Nuclear organization	Reuter and Wolff, 1981; Hari <i>et al.</i> , 2001
Piwi	RNAi	Pal-Bhadra <i>et al.</i> , 2004

(Continues)

Table 1.1. (Continued)

Locus	Function	References
Homeless/Spn-E	RNAi	Pal-Bhadra <i>et al.</i> , 2004
Aubergine	RNAi	Pal-Bhadra <i>et al.</i> , 2004
mod/mdg4 [E(var) 3–93D)]	Boundary element	Dorn <i>et al.</i> , 1993; Gerasimova <i>et al.</i> , 1995
BEAF-32	Boundary element	Zhao <i>et al.</i> , 1995; Gilbert <i>et al.</i> , 2006
Lip/Rm62	Helicase	Csink <i>et al.</i> , 1994
dCAP-G	Condensin subunit	Dej <i>et al.</i> , 2004
gluon/SMC4	Condensin subunit	Cobbe <i>et al.</i> , 2006
wapl	Chromosome adhesion?	Verni <i>et al.</i> , 2000

by neighboring posttranslational modifications on the histone tail, thus providing a combinatorial complexity to the signaling process (Dyson *et al.*, 2001; Yang, 2004). Although histone acetylation is generally associated with active chromatin regions, acetylation of H4K12 is found in heterochromatic regions, suggesting that acetylation at particular sites may mediate specific effects on chromatin function (Turner *et al.*, 1992).

Whereas acetylation is, with few exceptions, associated with euchromatin, different methylated histone modifications have been found to play an important role in differentiating active (euchromatic) chromosomal domains from silenced (heterochromatic) domains (reviewed in Lachner *et al.*, 2003). In general, the “activating” modifications consist of methylated H3K4, H3K36, and H3K79, while the “silencing” modifications are composed of methylated H3K9, H3K27, and H4K20. Thus, identification of the enzymes affecting these modifications and elucidating the mechanisms underlying their targeting promised to provide significant insight in the specification of chromatin domains. Given that the early genetic and cytogenetic experiments indicated that PEV occurs when heterochromatin spreads into juxtaposed euchromatic sequences, it was anticipated that many of the modifiers of position effect would fall into the categories of histone-modifying and histone-binding proteins (Schotta *et al.*, 2003a).

1. Methylation and the spread of heterochromatin

Three *Su(var)* gene products that were of early interest due to their heterochromatic localization, their “haplo-suppressor/triplo-enhancer” behavior, and their physical and genetic interactions were SU(VAR)2–5 (also known as SU(VAR)205, heterochromatin protein 1, or HP1) (Eissenberg *et al.*, 1990; James and Elgin, 1986; James *et al.*, 1989), SU(VAR)3–7 (Cléard and Spierer, 2001; Cléard

et al., 1997; Reuter *et al.*, 1990), and SU(VAR)3–9, the latter of which is one of the strongest modifiers of PEV known, showing dominance over nearly all other enhancers of variegation tested (Reuter *et al.*, 1986; Tschiertsch *et al.*, 1994; Wustmann *et al.*, 1989). Interestingly, one of the only two exceptions was *ptn*^D, which is a gain of function allele of *Su(var)*3–9 that causes ectopic localization of the protein to euchromatic sites (Kuhfittig *et al.*, 2001). A critical advance in the field of heterochromatin biology came from the findings that (1) the mammalian orthologs of SU(VAR)3–9 encode a methyltransferase that specifically methylates H3K9 (Rea *et al.*, 2000), an activity that is conserved in *Drosophila* SU(VAR)3–9 (Czernin *et al.*, 2001; Schotta *et al.*, 2002) and (2) the chromo-domain of HP1 binds to the methylated K9 of histone H3 (Bannister *et al.*, 2001; Lachner *et al.*, 2001). The silencing activity of HP1 was disrupted when its ability to bind methylated H3K9 was abolished by mutation (Jacobs *et al.*, 2001). Because the H3K9 methyltransferase SU(VAR)3–9 associates both with the H3K9 deacetylase HDAC1 (Czernin *et al.*, 2001) and with HP1 (Schotta *et al.*, 2002; Yamamoto and Sonoda, 2003), a spreading model for heterochromatin involving these components was proposed in which deacetylation of H3K9 clears the residue for methylation by SU(VAR)3–9 to enable HP1 binding, which, in turn, recruits additional SU(VAR)3–9 to methylate the adjacent nucleosome that provides another HP1-binding site in a self-propagating process (Grewal and Elgin, 2002; Schotta *et al.*, 2003b).

2. Initiation of heterochromatin formation

In recent years, a number of studies have indicated that RNAi-mediated silencing pathways can initiate the formation of heterochromatin (reviewed in Grewal and Rice, 2004; Matzke and Birchler, 2005; Zaratiegui *et al.*, 2007). This pathway was first identified in *S. pombe* where mutations in components of the RNAi pathway were found to disrupt centric heterochromatin silencing (Verdel *et al.*, 2004). Transcripts from repetitive elements in the centromeric region are processed into siRNAs that are incorporated into a RITS (RNAi-induced transcriptional silencing) complex that recognizes and binds homologous regions to initiate gene silencing mediated via H3K9 methylation (Verdel and Moazed, 2005; Verdel *et al.*, 2004). *Drosophila* centromeres are also composed of short satellite- and transposon-fragment repeats (Sun *et al.*, 1997) that are actively transcribed (Lakhotia and Jacob, 1974) and transgenes inserted into these regions show PEV (Cryderman *et al.*, 1998; Wallrath and Elgin, 1995; Wallrath *et al.*, 1996). Mutations in genes encoding the RNAi pathway Argonaute homologues *piwi*, *aubergine*, and the helicase *homeless* disrupt HP1 localization and suppress transgene silencing, indicating a similar mechanism for RNAi-mediated heterochromatin assembly operates in *Drosophila* as well (Pal-Bhadra *et al.*, 2004).

Mutants in Argo2, another RNAi argonaut family member, showed defects in centromeric heterochromatin assembly, including abnormal HP1 localization and defects in H3K9 methylation (Deshpande *et al.*, 2005).

3. Terminating heterochromatic spreading

That PEV is, by definition, a *variegated* phenotype indicates that chromosomal silencing can spread different lengths in different cells, and the question of what terminates the spreading is not well understood. Because of the “haplo-suppressor/triplo-enhancer” behavior of certain key components of heterochromatin, some models have postulated that heterochromatic-promoting factors are present in limiting amounts and spreading continues until components are exhausted (Locke *et al.*, 1988; Zuckerkandl, 1974). Alternatively, a “boundary model” was proposed in which discrete sites promoted initiation and termination of heterochromatin (Tartof *et al.*, 1984), but discrete sites conferring these activities could not be identified. Recently, the examination of PEV in *JIL-1* mutants in *Drosophila* has revealed a role for H3S10 phosphorylation in antagonizing heterochromatic spreading that suggests both models may apply in different genomic contexts (Bao *et al.*, 2007b; Ebert *et al.*, 2004; Lerach *et al.*, 2006; Zhang *et al.*, 2006). The *JIL-1* histone H3S10 tandem kinase localizes specifically to euchromatic interband regions of polytene chromosomes (Jin *et al.*, 1999; Wang *et al.*, 2001) and loss of *JIL-1* results in an array of chromosomal defects, including disruption of banded regions that normally do not contain *JIL-1* (Deng *et al.*, 2005). Alterations in band morphology in these mutants is accompanied by extensive spreading of the major heterochromatin markers H3K9me2 and HP1 to ectopic locations on the chromosome arms with the most pronounced increase on the X chromosomes (Zhang *et al.*, 2006). As would be expected, if *JIL-1* is required to “mark” euchromatic regions and terminate heterochromatic spreading, when a *white* reporter transgene is inserted into pericentric heterochromatin *JIL-1* loss-of-function alleles act as enhancers of variegation (i.e., the *white* gene is silenced; Bao *et al.*, 2007b). However, these very same *JIL-1* loss-of-function alleles act as *sub*suppressors of variegation for the *w^{m4}* inversion chromosome (Lerach *et al.*, 2006). In this case, the redistribution of the major heterochromatin markers to ectopic locations on the chromosome arms in *JIL-1* loss-of-function mutants decreases the concentration of these components at the centromere, and the increased expression of the *white* gene reflects the reduced extent of pericentromeric silencing (Lerach *et al.*, 2006) (Fig. 1.5). A gain-of-function *JIL-1* allele *JIL-1^{Su(var)3-1}*, however, is able to mark the euchromatin and prevent silencing in both instances (Bao *et al.*, 2007b; Ebert *et al.*, 2004). *JIL-1^{Su(var)3-1[3]}* is a C-terminal truncation that retains its ability to phosphorylate H3S10 but is mislocalized, showing a broad distribution on polytene chromosomes




Figure 1.5. Three diagrams illustrating position-effect variegation (PEV) inactivation of genes by heterochromatic spreading. (A) In a genome with normal amounts of JIL-1 protein, heterochromatic spreading from the *w^{m4}* inversion breakpoint often reaches the *white* gene; the gene is heterochromatized and is silenced. (B) In a genotype with significantly reduced JIL-1 levels, ectopic heterochromatization is widespread, resulting in lower levels of heterochromatin factors at the centromere. Therefore, heterochromatic spreading does not extend as far into the euchromatin and usually does not reach the *white* gene, leaving the gene euchromatic and active. (C) If a P element inserts a copy of the *white* gene into heterochromatin in a genotype where JIL-1 is present, JIL-1 often establishes a euchromatic chromatin structure at the *white* gene, the gene is protected from becoming heterochromatized, and the gene remains active. If the same insert is in a genome where JIL-1 is not present, the *white*

(Ebert *et al.*, 2004; Zhang *et al.*, 2006). Ectopic H3S10ph appears to restrict the formation of heterochromatin, both at the w^{m^4} inversion breakpoint and in the pericentric transgenes, resulting in a dominant suppression of variegation (Bao *et al.*, 2007b; Ebert *et al.*, 2004). Recently, H3S10 phosphorylation has been shown to displace HP1 binding to methylated H3K9 during mitosis (Fischle *et al.*, 2005; Hirota *et al.*, 2005). Likewise, JIL-1 kinase phosphorylation of H3S10 may displace or prevent HP1 binding, thereby terminating heterochromatic spreading at that site in order to establish a euchromatic domain (Johansen and Johansen, 2006). Consistent with this model, the *JIL-1^{Su(var)3-1}* C-terminal truncation alleles are dominant over the triplo-enhancer effects of *Su(var)3-9* and *HP1* (Ebert *et al.*, 2004).

A second mechanism that operates to restrict heterochromatic spreading is the incorporation of the histone variant H3.3 into actively transcribed regions (Ahmad and Henikoff, 2002; Schwartz and Ahmad, 2005). Genome-wide profiling of histone H3.3 replacement patterns revealed that H3.3 is also enriched far upstream and downstream of active genes, but it was suggested this may reflect intergenic transcription that is now known to be widespread (Mito *et al.*, 2005). Recently, GAGA factor was found to interact with FACT (facilitates chromatin transcription) complex to direct H3.3 replacement (Nakayama *et al.*, 2007). GAGA factor had originally been identified as the enhancer of variegation E(var)3-93D (Dorn *et al.*, 1993) and in this recent study, mutation in either GAGA factor or in DRE4/spt16, a subunit of FACT, enhanced w^{m^4} PEV (Nakayama *et al.*, 2007). Furthermore, the GAGA factor-FACT complex occupies a site where H3K4 methylation peaks and H3K9 methylation dips, suggesting that the remodeling activity of GAGA-FACT serves a barrier function against heterochromatic spreading and silencing (Nakayama *et al.*, 2007). The absence of such a barrier may result in expanded heterochromatic silencing, thus titrating heterochromatic factors away from the pericentromeric region resulting in an e(var) phenotype in the GAGA mutant.

B. Nuclear organization

The phenomenon of *bw^D* PEV in which chromosomal pairing brings a functional *bw* allele into a “silencing compartment” underscores the impact of nuclear organization on gene expression (Csink and Henikoff, 1996; Csink *et al.*, 2002;

gene is not protected. The establishment of ectopic heterochromatic sites throughout the chromosomes lowers the level of heterochromatin factors in the pericentric region. However, the basal level of pericentric heterochromatin, maintained by the RNAi system that generates heterochromatin, is sufficient to spread into the inserted *white* gene, the gene becomes heterochromatized, and is silenced.

Dernburg *et al.*, 1996). The functional consequences of long-range nuclear reorganization on gene expression were examined by FISH for three known variegating genes located on different chromosomes and in each case a strong correlation was observed between silencing and association with centromeric satellite sequences (Harmon and Sedat, 2005). Thus, changes in nuclear chromosomal organization have clear consequences on gene expression. What establishes the organization of the nucleus, however, is still not well understood (Jackson, 2003; Lanctôt *et al.*, 2007; Misteli, 2007).

Lamins are the best characterized nuclear architectural components to date and their role in chromosomal organization and gene expression is increasingly appreciated (reviewed in Hutchison, 2002; Goldman *et al.*, 2005). Association of genes with the nuclear lamina is generally linked to gene silencing (Andrulis *et al.*, 1998; Taddei and Gasser, 2004), although in some cases interaction with nuclear pore complex components is associated with transcriptional activation (Casolari *et al.*, 2004; Ishii *et al.*, 2002; Mendjan *et al.*, 2006). In a recent study, Pickersgill *et al.* (2006) identified ~500 genes that interact with lamin Dm₀ (the *Drosophila* B-type lamin) and found that they are transcriptionally repressed. Lamin-bound genes tend to be clustered in the genome, developmentally coregulated, have long intergenic domains, and lack active histone marks, suggesting that these regions likely comprise compacted chromatin (Pickersgill *et al.*, 2006). Consistent with lamins playing a role in epigenetic regulation of gene expression, a lamin Dm₀ C-terminal truncation mutant Lam^{Ani3} modified the PEV of *w^{m4}* (Bao *et al.*, 2007a). However, this allele shows an *enhancer* of variegation effect indicating increased silencing when the C-terminus is deleted, indicating that the role of lamin B in organizing chromatin domains is complex. Although interaction with lamin B is associated with gene silencing (Pickersgill *et al.*, 2006), the PEV studies suggest that lamin B's C-terminus may play a role in reversing silencing effects (Bao *et al.*, 2007a).

Interaction between nuclear lamins and chromatin is thought to regulate higher order chromatin organization and a direct physical interaction between lamin Dm₀ and histone H2A has been reported (Mattout *et al.*, 2006; Rzepecki *et al.*, 1998). In addition, different chromatin-organizing proteins have been found to interact with nuclear lamins and nuclear envelope proteins and the nature of these activities may facilitate the complex integration of various signaling pathways to influence genome organization and gene regulation (Gruenbaum *et al.*, 2005). Signaling to the chromatin may be mediated at several different levels besides physical association to the lamina. For example, the nuclear envelope protein LAP2 β recruits a histone deacetyltransferase to the periphery, thus triggering silencing histone modifications in this domain (Somech *et al.*, 2005).

The nuclear lamina has also been found to bind the ubiquitin ligase dTopors (Topoisomerase-I-interacting protein), which functions in concert with proteins that bind to a chromatin regulatory domain known as an insulator or boundary element (Capelson and Corces, 2005; Wei *et al.*, 2005; West *et al.*, 2002). Insulators block enhancer interactions (Geyer and Corces, 1992; Kellum and Schedl, 1992) and block the spreading of one chromatin state into another (Sun and Elgin, 1999). The *gypsy* insulator sequence was originally identified as a *gypsy* retrotransposon insertion that blocked enhancer function at the *yellow* locus by introducing a Su(Hw) (suppressor of Hairy wing)-binding site (Geyer and Corces, 1992; Parkhurst *et al.*, 1988; Roseman *et al.*, 1993). The Su(Hw) protein binds to BTB/POZ-domain proteins (Broad complex Tramtrack, Bric-abrac/Poxvirus and Zinc finger), CP190 (Pai *et al.*, 2004) and mod/mdg4 (Gerasimova *et al.*, 1995; Ghosh *et al.*, 2001) to form a complex with dTopors at the nuclear periphery that is thought to isolate regions of DNA in “insulator bodies” (Capelson and Corces, 2004; Gerasimova *et al.*, 2000). The presence of *gypsy* insulator sequences at endogenous sites in the genome suggests that this may provide a means to compartmentalize the genome and prevent heterochromatic spreading into active euchromatic regions (Parnell *et al.*, 2006; Ramos *et al.*, 2006). Consistent with a role in limiting heterochromatic spreading, *mod/mdg4* was originally identified as the enhancer of variegation *E(var)3–93D* (Dorn *et al.*, 1993).

Because there are significantly more chromosomal-binding sites for Su(Hw) and *mod/mdg4* than genomic consensus Su(Hw) DNA-binding sites (Parnell *et al.*, 2006; Ramos *et al.*, 2006), it is not known whether the PEV modifier effects observed in the *mod/mdg4* mutant reflect activities related directly to *gypsy* insulator function, activities at other as-yet uncharacterized chromosomal sites, or is the indirect consequence of the nuclear reorganization that has been found to occur when *gypsy* insulator components are disrupted (Gerasimova *et al.*, 2000; Pai *et al.*, 2004). One plausible scenario is that if the *mod/mdg4* mutation compromises the insulator’s barrier function to heterochromatic spreading, the resultant increased euchromatic distribution of heterochromatic factors reduces their levels at the pericentromeric regions resulting in decreased spreading at the *w^{m4}* inversion and an E(var) phenotype, much as occurs for JIL-1 (Bao *et al.*, 2007b). Likewise, proper insulator function also depends on the presence of functional RNAi-mediated pathways, perhaps to ensure the generation of heterochromatin and thus the appropriate sequestration of heterochromatic components at centromeric regions. When Argonaute RNAi pathway genes are mutated, nuclear architecture is disrupted and insulator activity is decreased (Lei and Corces, 2006). Furthermore, CP190 interacts with Rm62/Lip (Lei and Corces, 2006), a DEAD-box helicase that has also been identified as a modifier of PEV and is required for dsRNA-mediated silencing, heterochromatin formation, and transposon silencing (Csink *et al.*, 1994; Ishizuka *et al.*, 2002). These

results underscore the relationship between genome organization/chromatin packaging of euchromatic sites and the levels of heterochromatic factors that are available to spread at pericentromeric sites (i.e., the inversion breakpoint).

Another nuclear domain besides the periphery that has been associated with silencing is the nucleolus (Lawrence and Pikaard, 2004; Lewis and Pikaard, 2001; Wang *et al.*, 2005; Yasufzi *et al.*, 2004) and chromosomal inversions that place the *white* gene adjacent to the nucleolar organizer region (NOR) exhibit PEV consistent with different levels of gene silencing (Spofford and DeSalle, 1991). Recently, it was found that heterochromatin-mediated silencing pathways including the RNAi pathway, HP1, and H3K9 methylation regulate nucleolar organization and structural integrity (Peng and Karpen, 2006). In most of the RNAi pathway mutants examined as well as in *Su(var)2-5/HP1* and *Su(var)3-9* mutants, heterochromatic repeated DNAs became dispersed and multiple, ectopic nucleoli were formed. In addition, significant amounts of extrachromosomal circles (ecc) of rDNA accumulated (Peng and Karpen, 2006). It was proposed that loss of heterochromatization in the NOR correlates with increased recombination or repair rates in this repetitive sequence region with subsequent accumulation of ecc rDNAs when errors occurred (Peng and Karpen, 2006). In support of this idea, mutation in *Ligase4*, a DNA repair enzyme, suppressed ecc generation (Peng and Karpen, 2006). A connection between DNA synthesis/repair and heterochromatin behavior has been inferred by a number of studies showing that mutations in different factors involved in these processes act as PEV modifiers (Table 1.1). Another mutation that disrupts nucleolar and chromosomal organization in *Drosophila*, *Su(var)2-10/dPIAS* (Hari *et al.*, 2001), also was originally identified as a modifier of PEV (Reuter and Wolff, 1981). Interestingly, a correlation between deregulation of chromatin-silencing and disruption of nuclear organization has also been observed in yeast (Teixeira *et al.*, 2002), suggesting that the link between heterochromatin behavior and nuclear organization is a general feature of eukaryotes.

V. CONCLUDING REMARKS

Studies of the control of gene regulation that began with the discovery of PEV and the chromosomal conditions that cause it have brought us a long way toward understanding the role of chromatin structure in the regulation of gene expression. Studies of chromosome rearrangements and P element insertions demonstrated that a change in chromatin structure from euchromatic to heterochromatic leads to gene silencing and that this silencing effect has specific characteristics that are amenable to analysis. The identification of mutations that modify the PEV phenotype identified a number of factors that are part of or

regulate the formation or maintenance of heterochromatin and euchromatin. The findings of these studies indicate that control of gene expression is a complex process involving regulatory mechanisms that are integrated at multiple hierarchical levels ranging from the primary DNA sequence to the chromatin packaging of DNA and higher order chromosomal folding to the three-dimensional spatial organization of chromosomes within the nucleus (reviewed in van Driel *et al.*, 2003; Misteli, 2007). A key component of this process may be the modification of histone N-terminal tails to “mark” a region of chromatin for the activation of a number of specific downstream responses (Fischle *et al.*, 2003; Kouzarides, 2007; Strahl and Allis, 2000; Turner, 2002) such as the establishment of transcriptionally active chromatin (Brownell and Allis, 1996; Csordas, 1990; Turner, 2000) or higher level folding associated with silencing (Turner, 2002; Zhang and Reinberg, 2001). However, regulation of gene expression at the level of the nucleosome goes beyond the modification of histone tails. Genes that are actively transcribed are often subject to nucleosomal remodeling by chromatin-remodeling complexes (Becker and Horz, 2002; Workman and Kingston, 1998) and characterized by nucleosome-free or “DNase-hypersensitive” sites, whereas silenced or heterochromatic regions are characterized by tightly packed, ordered nucleosomal arrays (Elgin, 1988; Wallrath and Elgin, 1995). Besides the local remodeling of a small number of nucleosomes in the promoter region, larger scale nucleosomal remodeling is also implicated in the unfolding of large chromatin domains (Dietzel *et al.*, 2004; Neely and Workman, 2002; Peterson, 2003; Tumbar *et al.*, 1999). Current models suggest that within the nucleus, there exist regions of condensed silent chromatin interspersed with regions of decondensed, active chromatin. A clear example of this is found within the band-interband pattern observed in *Drosophila* larval polytene chromosomes (Labrador and Corces, 2002). The challenge is to identify the molecules and molecular mechanisms that determine how this organization is established and maintained and the signal transduction events that regulate this process.

A constructive way to think about heterochromatin is in quantitative terms. The *Drosophila* genome contains a finite number of repeated sequences that interact with the RNAi system to utilize a fixed quantity of heterochromatic factors that can bind to and heterochromatize a set length of chromatin. The concentration of these factors is normally high in the pericentric chromosomal regions, which serve as a reservoir for heterochromatic components. The amount of chromatin that is heterochromatic during interphase is a dynamic balance between the number of heterochromatic factors and the number of chromatin sites available for binding. Anything that increases the number of sites or decreases the number of factors will decrease the pericentric concentration and anything that decreases the number of sites or increases the number of factors will increase the pericentric concentration. Consider three types of

Su(var) phenotypes. (1) The addition of Y chromosomes or heterochromatic regions increases the sequences competing to bind heterochromatic factors. This increased competition lowers the concentration of heterochromatic factors in the pericentric heterochromatin. (2) Mutations that reduce the dose of genes that code for structural components of heterochromatin (e.g., HP1) also reduce the quantity of heterochromatic factors in pericentric regions. (3) JIL-1 proteins establish euchromatic regions that are resistant to heterochromatic factor binding. In loss-of-function JIL-1 mutants, ectopic-binding sites for heterochromatic factors are established throughout the genome that compete for heterochromatic factors. This reduces the concentration of heterochromatic factors in pericentric regions. In all three cases, the lower level of heterochromatic factors in pericentric regions means that heterochromatin cannot spread as far into euchromatin from a breakpoint and adjacent genes are not inactivated as often, resulting in a Su(var) phenotype. E(var) phenotypes result from mutations that increase the concentration of heterochromatic factors in the pericentric regions, leading to increased lengths of spreading from breakpoints.

The details of a number of the most interesting aspects of PEV remain to be explained. How does the regulation of genes normally residing in heterochromatic regions differ from that of genes residing in euchromatic regions? How does the PEV of genes inserted into telomeres by P element transposition differ from that of those inserted into pericentric regions? What controls nuclear compartmentalization and how does this normally act to control gene regulation? These and other questions are currently being actively investigated in a continuation of an examination that began more than 75 years ago with the observation of a novel color of the eyes of a fly.

Acknowledgments

We thank members of the lab for critical reading of the manuscript and for helpful comments. We thank Drs. Lori Wallrath and Diane Cryderman (University of Iowa) for helpful discussion and for providing the 118E-10 P element insertion line. The authors' work on JIL-1 is supported by NIH grant GM62916.

References

- Ahmad, K., and Golic, K. G. (1996). Somatic reversion of chromosomal position effects in *Drosophila melanogaster*. *Genetics* **144**, 657–670.
- Ahmad, K., and Henikoff, S. (2002). The histone variant H3.3 marks active chromatin by replication-independent nucleosome assembly. *Mol. Cell* **9**, 1191–1200.
- Altorfer, N. (1967). Effect of the absence of the Y chromosome on the expression of the cubitus interruptus phenotype in various translocations in *Drosophila melanogaster*. *Genetics* **55**, 755–767.
- Andrulis, E. D., Neiman, A. M., Zappulla, D. C., and Sternglanz, R. (1998). Perinuclear localization of chromatin facilitates transcriptional silencing. *Nature* **394**, 592–595.

- Aulner, N., Monod, C., Mandicourt, G., Julien, D., Cuvier, O., Sall, A., Janssen, S., Laemml, U. K., and Käs, E. (2002). The AT-hook protein D1 is essential for *Drosophila melanogaster* development and is implicated in position-effect variegation. *Mol. Cell. Biol.* **22**, 1218–1232.
- Bahn, E. (1971). Position effect variegation for an isoamylase in *Drosophila melanogaster*. *Hereditas* **67**, 79–82.
- Baker, W. K. (1953). V-type position effects on a gene normally found in heterochromatin. *Genetics* **38**, 328–344.
- Baker, W. K. (1971). Evidence for position effect suppression of the ribosomal RNA cistrons in *Drosophila melanogaster*. *Proc. Natl. Acad. Sci. USA* **68**, 2472–2476.
- Baker, W. K., and Reim, A. (1962). The dichotomous action of Y chromosomes on the expression of position-effect variegation. *Genetics* **47**, 1399–1407.
- Baker, W. K., and Spofford, J. B. (1959). Heterochromatic control of position-effect variegation in *Drosophila*. *Univ. Texas Publ.* **5914**, 135–154.
- Baksa, K., Morawietz, H., Dombradi, V., Axton, M., Taubert, H., Szabo, G., Torol, I., Udvardy, A., Gurkovic, B., Szoor, B., Gilover, D., Reuter, G., et al. (1993). Mutations in the protein phosphatase 1 gene at 87B can differentially affect suppression of position-effect variegation and mitosis in *Drosophila melanogaster*. *Genetics* **135**, 117–125.
- Bannister, A. J., Zegerman, P., Partridge, J. F., Miska, E. A., Thomas, J. O., Allshire, R. C., and Kouzarides, T. (2001). Selective recognition of methylated lysine 9 on histone H3 by the HP1 chromo domain. *Nature* **410**, 120–124.
- Bao, X., Girton, J., Johansen, J., and Johansen, K. M. (2007a). The lamin Dm₀ allele *Ari3* acts as an enhancer of position effect variegation of the *wm4* allele in *Drosophila*. *Genetica* **129**, 339–342.
- Bao, X., Deng, H., Johansen, J., Girton, J., and Johansen, K. M. (2007b). Loss-of-function alleles of the JIL-1 histone H3S10 kinase enhance position-effect-variegation at pericentric sites in *Drosophila* heterochromatin. *Genetics* **176**, 1355–1358.
- Becker, P. B., and Horz, W. (2002). ATP-dependent nucleosome remodeling. *Annu. Rev. Biochem.* **71**, 247–273.
- Beckstead, R. B., Ner, S. S., Hales, K. G., Grigliatti, T. A., Baker, B. S., and Bellen, H. J. (2005). Bonus, a *Drosophila* TIF1 homolog, is a chromatin-associated protein that acts as a modifier of position-effect variegation. *Genetics* **169**(2), 783–794.
- Benner, D. B. (1971). Some evidence against the presence of suppressors of variegation on the Y chromosome. *Dros. Inform. Serv.* **47**, 72.
- Bernstein, B. E., Kamal, M., Lindblad-Toh, K., Bekiranov, S., Bailey, D. K., Huebert, D. J., McMahon, E. K., Karlsson, E. K., Kulbokas, E. J. 3rd., Gingeras, T. R., Schreiber, S. L., and Lander, E. S. (2005). Genomic maps and comparative analysis of histone modifications in human and mouse. *Cell* **120**, 169–181.
- Brownell, J. E., and Allis, C. D. (1996). Special HATs for special occasions: Linking histone acetylation to chromatin assembly and gene activation. *Curr. Opin. Genet. Dev.* **6**, 176–184.
- Büchner, K., Roth, P., Schotta, G., Krauss, V., Saumweber, H., Reuter, G., and Dorn, R. (2000). Genetic and molecular complexity of the position effect variegation modifier mod(mdg4) in *Drosophila*. *Genetics* **155**, 141–157.
- Capelson, M., and Corces, V. G. (2004). Boundary elements and nuclear organization. *Biol. Cell* **96**, 617–629.
- Capelson, M., and Corces, V. G. (2005). The ubiquitin ligase dTopors directs the nuclear organization of a chromatin insulator. *Mol. Cell* **20**, 105–116.
- Casolari, J. M., Brown, C. R., Komili, S., West, J., Hieronymus, H., and Silver, P. A. (2004). Genome-wide localization of the nuclear transport machinery couples transcriptional status and nuclear organization. *Cell* **117**, 427–439.

- Cheung, P., Allis, C. D., and Sassone-Corsi, P. (2000). Signaling to chromatin through histone modifications. *Cell* **103**, 263–271.
- Cléard, F., and Spierer, P. (2001). Position-effect variegation in *Drosophila*: The modifier *Su(var)3-7* is a modular DNA-binding protein. *EMBO Rep.* **21**, 1095–1100.
- Cléard, F., Delattre, M., and Spierer, P. (1997). SU(VAR)3-7: A *Drosophila* heterochromatin-associated protein and companion of HP1 in the genomic silencing of position-effect variegation. *EMBO J.* **16**, 5280–5288.
- Cobbe, N., Savvidou, E., and Heck, M. M. S. (2006). Diverse mitotic and interphase functions of condensins in *Drosophila*. *Genetics* **172**, 991–1008.
- Cryderman, D. E., Cuaycong, M. H., Elgin, S. C., and Wallrath, L. L. (1998). Characterization of sequences associated with position-effect variegation at pericentric sites in *Drosophila* heterochromatin. *Chromosoma* **107**, 277–285.
- Csink, A. K., and Hennikoff, S. (1996). Genetic modification of heterochromatic association and nuclear organization in *Drosophila*. *Nature* **381**, 529–531.
- Csink, A. K., Linsk, R., and Birchler, J. A. (1994). The *Lighten up* (*Lip*) gene of *Drosophila melanogaster*, a modifier of retroelement expression, position effect variegation and *white* locus insertion alleles. *Genetics* **138**, 153–163.
- Csink, A. K., Bounoutas, A., Griffith, M. L., Sabl, J. F., and Sage, B. T. (2002). Differential gene silencing by trans-heterochromatin in *Drosophila melanogaster*. *Genetics* **160**, 257–269.
- Csordas, A. (1990). On the biological role of histone acetylation. *Biochem. J.* **265**, 23–38.
- Czernin, B., Schotta, G., Hulsmann, B. B., Brehm, A., Becker, P. B., Reuter, G., and Imhof, A. (2001). Physical and functional association of SU(VAR)3-9 and HDAC1 in *Drosophila*. *EMBO Rep.* **2**, 915–919.
- Dej, K. J., Ahn, C., and Orr-Weaver, T. L. (2004). Mutations in the *Drosophila* condensin subunit dCAP-G: defining the role of condensin for chromosome condensation in mitosis and gene expression in interphase. *Genetics* **168**, 895–906.
- Demerec, M. (1940). Genetic behavior of euchromatic segments inserted into heterochromatin. *Genetics* **25**, 618–627.
- Demerec, M. (1941). The nature of changes in the *white*–*Notch* region of the X chromosome of *Drosophila melanogaster*. *Proc. 7th Int. Genet. Congr.* 99–103.
- Demerec, M., and Slizynska, H. (1937). Mottled *white*^{258–18} of *Drosophila melanogaster*. *Genetics* **22**, 641–649.
- Deng, H., Zhang, W., Bao, X., Martin, J. N., Girton, J., Johansen, J., and Johansen, K. M. (2005). The JIL-1 kinase regulates the structure of *Drosophila* polytene chromosomes. *Chromosoma* **114**, 173–182.
- Dernburg, A. F., Broman, K. W., Fung, J. C., Marshall, W. F., Philips, J., Agard, D. A., and Sedat, J. W. (1996). Perturbation of nuclear architecture by long-distance chromosome interactions. *Cell* **85**, 745–759.
- De Rubertis, F., Kadosh, D., Henchoz, S., Pauli, D., Reuter, G., Struhl, K., and Spierer, P. (1996). The histone deacetylase RPD3 counteracts genomic silencing in *Drosophila* and yeast. *Nature* **384**, 589–591.
- Deshpande, G., Calhoun, G., and Schedl, P. (2005). *Drosophila* argonaute-2 is required early in embryogenesis for the assembly of centric/centromeric heterochromatin, nuclear division, nuclear migration, and germ cell formation. *Genes Dev.* **19**, 1680–1685.
- Dietzel, S., Zolghadr, K., Hepperger, C., and Belmont, A. S. (2004). Differential large-scale chromatin compaction and intranuclear positioning of transcribed versus non-transcribed transgene arrays containing β -globin regulatory sequences. *J. Cell Sci.* **117**, 4603–4614.
- Dobzhansky, T. (1936). Position effects on genes. *Biol. Rev.* **11**, 364–384.

- Dorn, R., Szidonya, J., Korge, G., Sehnert, M., Taubert, H., Archoukieh, E., Tschiersch, B., Morawietz, H., Wustmann, G., Hoffmann, G., and Reuter, G. (1993). P transposon-induced dominant enhancer mutations of position-effect variegation in *Drosophila melanogaster*. *Genetics* **133**, 279–290.
- Dreesen, T. D., Johnson, D. H., and Henikoff, S. (1988). The brown protein of *Drosophila melanogaster* is similar to the white protein and to components of active transport complex. *Mol. Cell. Biol.* **8**, 5206–5215.
- Dreesen, T. D., Henikoff, S., and Loughney, K. (1991). A pairing-sensitive element that mediates trans-inactivation is associated with *Drosophila* brown gene. *Genes Dev.* **5**, 331–340.
- Dubinin, N. P., and Sidorov, B. N. (1935). Position effect of the gene hairy. *Biol. Zh.* **4**, 555–568.
- Dyson, M. H., Rose, S., and Mahadevan, L. C. (2001). Acetyllysine-binding and function of bromodomain-containing proteins in chromatin. *Front. Biosci.* **6**, D853–D865.
- Eberl, D. F., Dufy, B. J., and Hilliker, A. J. (1993). The role of heterochromatin in the expression of a heterochromatic gene, the *rolled* locus of *Drosophila melanogaster*. *Genetics* **134**, 277–292.
- Ebert, A., Schotta, G., Lein, S., Kubicek, S., Krauss, V., Jenuwein, T., and Reuter, G. (2004). Su(var) genes regulate the balance between euchromatin and heterochromatin in *Drosophila*. *Genes Dev.* **18**, 2973–2983.
- Ebert, A., Lein, S., Schotta, G., and Reuter, G. (2006). Histone modification and the control of heterochromatic gene silencing in *Drosophila*. *Chromosome Res.* **14**, 377–392.
- Eissenberg, J. C., James, T. C., Foster-Hartnett, D. M., Ngan, V., and Elgin, S. C. R. (1990). Mutation in a heterochromatin-specific protein is associated with suppression of position effect variegation in *Drosophila melanogaster*. *Proc. Natl. Acad. Sci. USA* **87**, 9923–9927.
- Eissenberg, J. C., Morris, G. D., Reuter, G., and Hartnett, T. (1992). The heterochromatin-associated protein HP-1 is an essential protein in *Drosophila* with dosage-dependent effects on position effect variegation. *Genetics* **131**, 345–352.
- Elgin, S. C. (1988). The formation and function of DNase I hypersensitive sites in the process of gene activation. *J. Biol. Chem.* **263**, 19259–19262.
- Ephrussi, B., and Sutton, E. (1944). A reconsideration of the mechanism of position effect. *Proc. Natl. Acad. Sci. USA* **30**, 183–197.
- Farkas, G., Gausz, J., Galloni, M., Reuter, G., Gyurkovics, H., and Karch, F. (1994). The *Trithorax-like* gene encodes the *Drosophila* GAGA factor. *Nature* **371**, 806–808.
- Farkas, G., Leibovitch, B. A., and Elgin, S. C. R. (2000). Chromatin organization and transcriptional control of gene expression in *Drosophila*. *Gene* **253**, 117–136.
- Fischle, W., Wang, Y., and Allis, C. D. (2003). Regulation of HP1-chromatin binding by histone H3 methylation and phosphorylation. *Nature* **425**, 475–479.
- Fischle, W., Tseng, B. S., Dormann, H. L., Ueberheide, M. B., Garcia, B. A., Shabanowitz, J., Hunt, D. F., Funabiki, H., and Allis, C. D. (2005). Regulation of HP1-chromatin binding by histone H3 methylation and phosphorylation. *Nature* **438**, 1116–1122.
- Gatti, M., and Pimpinelli, S. (1992). Functional elements in *Drosophila melanogaster* heterochromatin. *Annu. Rev. Genet.* **26**, 239–275.
- Gerasimova, T. I., Gvozdev, V. A., and Birstein, V. (1972). Position effect variegation of *Pgl* locus determining 6-phosphogluconate dehydrogenase in *Drosophila melanogaster*. *Dros. Inform. Serv.* **48**, 81.
- Gerasimova, T. I., Gdula, D. A., Gerasimov, D. V., Simonova, O., and Corces, V. G. (1995). A *Drosophila* protein that imparts directionality on a chromatin insulator is an enhancer of position-effect variegation. *Cell* **82**, 587–597.
- Gerasimova, T. I., Byrd, K., and Corces, V. G. (2000). A chromatin insulator determines the nuclear localization of DNA. *Mol. Cell* **6**, 1025–1035.
- Gersh, E. S. (1949). Influence of temperature on the expression of position effects in the *scute-8* stock of *Drosophila melanogaster* and its relation to heterochromatization. *Genetics* **34**, 701–707.
- Gersh, E. S. (1963). Variegation at the *white* locus in *In(1)rst*. *Dros. Inform. Serv.* **37**, 81.

- Geyer, P. K., and Corces, V. G. (1992). DNA position-specific repression of transcription by a *Drosophila* zinc finger protein. *Genes Dev.* **6**, 1865–1873.
- Ghosh, D., Gerasimova, T. I., and Corces, V. G. (2001). Interactions between the Su(Hw) and Mod(mdg4) proteins required for gypsy insulator function. *EMBO J.* **20**, 2518–2527.
- Gilbert, M. K., Tan, Y. Y., and Hart, C. M. (2006). The *Drosophila* boundary element-associated factors BEAF-32A and BEAF-32B affect chromatin structure. *Genetics* **173**, 1365–1375.
- Goldman, R. D., Goldman, A. E., and Shumaker, D. K. (2005). Nuclear lamins: Building blocks of nuclear structure and function. *Novartis Found. Symp.* **264**, 3–16.
- Gowen, J. M., and Gay, E. H. (1934). Chromosome constitution and behavior in ever-sporting and mottling in *Drosophila melanogaster*. *Genetics* **19**, 189–126.
- Grell, R. F. (1958). The effect of X chromosome loss on variegation. *Dros. Inform. Serv.* **32**, 124.
- Grell, R. F. (1970). The time of initiation of segregational pairing between nonhomologues in *Drosophila melanogaster*. A reexamination of w^{m4} . *Genetics* **64**, 337–365.
- Grewal, S. I., and Elgin, S. C. (2002). Heterochromatin: New possibilities for the inheritance of structure. *Curr. Opin. Genet. Dev.* **12**, 178–187.
- Grewal, S. I., and Rice, J. C. (2004). Regulation of heterochromatin by histone methylation and small RNAs. *Curr. Opin. Cell Biol.* **16**, 230–238.
- Grienberger, A., Miotto, B., Sagnier, T., Cavalli, G., Schrake, V., Gelli, V., Mariol, M.-C., Berenger, Y., Graba, Y., and Pradel, J. (2002). The MYST domain acetyltransferase Chameau functions in epigenetic mechanisms of transcriptional repression. *Curr. Biol.* **12**, 762–766.
- Grigliatti, T. (1991). Position-effect variegation—an assay for nonhistone chromosomal proteins and chromatin assembly and modifying factors. *Methods Cell Biol.* **35**, 588–625.
- Gruenbaum, Y., Margalit, A., Goldman, R. D., Shumaker, D. K., and Wilson, K. L. (2005). The nuclear lamina comes of age. *Nat. Rev. Mol. Cell Biol.* **6**, 21–31.
- Hackstein, J. H. P., and Hochstenbach, R. (1995). The elusive fertility genes of *Drosophila*: The ultimate haven for selfish genetic elements. *Trends Genet.* **11**, 195–200.
- Hannah, A. H. (1951). Localization and function of heterochromatin in *Drosophila melanogaster*. *Adv. Genet.* **113**, 191–203.
- Hari, K. L., Cook, D. R., and Karpen, G. H. (2001). The *Drosophila* *Su(var)2-10* locus regulates chromosome structure and function and encodes a member of the PIAS family. *Genes Dev.* **15**, 1334–1348.
- Harmon, B., and Sedat, J. (2005). Cell-by-cell dissection of gene expression and chromosomal interactions reveals consequences of nuclear reorganization. *PLoS Biol.* **3**, e67.
- Hartman-Goldstein, I. J. (1967). On the relationship between heterochromatin and variegation in *Drosophila*, with special references to temperature sensitive periods. *Genet. Res.* **10**, 143–159.
- Hazelrigg, T., and Petersen, S. (1992). An unusual genomic position effect on *Drosophila white* gene expression: pairing dependence, interactions with *zeste*, and molecular analysis of revertants. *Genetics* **130**, 125–138.
- Hazelrigg, T., Levis, R., and Rubin, G. M. (1984). Transformation of *white* locus DNA in *Drosophila*: Dosage compensation, *zeste* interaction and position effects. *Cell* **36**, 469–481.
- Heitz, E. (1928). Das Heterochromatin der Moose. *Jahrb Wiss Botanik* **69**, 762–818.
- Henderson, D. S., Banga, S. S., Grigliatti, T. A., and Boyd, J. B. (1994). Mutagen sensitivity and suppression of position-effect variegation result from mutations in *mus209*, the *Drosophila* gene encoding PCNA. *EMBO J.* **13**, 1450–1459.
- Henikoff, S. (1979). Position effect and variegation enhancers in an autosomal region of *Drosophila melanogaster*. *Genetics* **93**, 105–115.
- Henikoff, S. (1994). A reconsideration of the mechanism of position effect. *Genetics* **138**, 1–5.
- Henikoff, S., and Dreesen, T. D. (1989). Trans-inactivation of the *Drosophila brown* gene: Evidence for transcriptional repression and somatic pairing dependence. *Proc. Natl. Acad. Sci. USA* **86**, 6704–6708.

- Hessler, A. Y. (1958). V-type position effect at the *light* locus in *Drosophila melanogaster*. *Genetics* **43**, 395–403.
- Hilliker, A. J. (1976). Genetic analysis of the centromeric heterochromatin of chromosome 2 of *Drosophila melanogaster*: Deficiency mapping of EMS-induced lethal complementation groups. *Genetics* **83**, 765–782.
- Hinton, T. (1940). Report on *D. melanogaster* new mutants. *Dros. Inform. Serv.* **13**, 49.
- Hinton, T. (1942). A comparative study of certain heterochromatic regions in the mitotic and salivary gland chromosomes of *Drosophila melanogaster*. *Genetics* **27**, 119–127.
- Hinton, T. (1949). The role of heterochromatin in position effect. *Proc. 8th Int. Congr. Genet.* 595–596.
- Hinton, T. (1950). A correlation of phenotypic changes and chromosomal rearrangements at the two ends of an inversion. *Genetics* **35**, 188–205.
- Hinton, T., and Goodsmith, W. (1950). An analysis of phenotypic reverions at the *brown* locus in *Drosophila*. *J. Exp. Zool.* **114**, 103–114.
- Hirota, T., Lipp, J. J., Toh, B. H., and Peters, J. M. (2005). Histone H3 serine 10 phosphorylation by Aurora B causes HP1 dissociation from heterochromatin. *Nature* **438**, 1176–1180.
- Huiszinga, K. L., Brower-Toland, B., and Elgin, S. C. R. (2006). The contradictory definitions of heterochromatin: Transcription and silencing. *Chromosoma* **115**, 110–122.
- Hutchison, C. J. (2002). Lamins: Building blocks or regulators of gene expression? *Nat. Rev. Mol. Cell Biol.* **3**, 848–858.
- Ishii, K., Aribi, G., Lin, C., Van Houwe, G., and Laemmli, U. K. (2002). Chromatin boundaries in budding yeast: The nuclear pore connection. *Cell* **109**, 551–562.
- Ishizuka, A., Siomi, M. C., and Siomi, H. A. (2002). A *Drosophila* fragile X protein interacts with components of RNAi and ribosomal proteins. *Genes Dev.* **16**, 2497–2508.
- Jackson, D. A. (2003). The principles of nuclear structure. *Chromosome Res.* **11**, 387–401.
- Jacobs, S. A., Taverna, S. D., Zhang, Y., Briggs, S. D., Li, J., Eissenberg, J. C., Allis, C. D., and Khorasanizadeh, S. (2001). Specificity of the HP1 chromo domain for the methylated N-terminus of histone H3. *EMBO J.* **20**, 5232–5241.
- James, T. C., and Elgin, S. C. R. (1986). Identification of a nonhistone chromosomal protein associated with heterochromatin in *Drosophila melanogaster* and its gene. *Mol. Cell Biol.* **6**, 3862–3872.
- James, T. C., Eissenberg, J. C., Craig, C., Dietrich, V., Hobson, A., and Elgin, S. C. (1989). Distribution patterns of HP1, a heterochromatin-associated nonhistone chromosomal protein of *Drosophila*. *Eur. J. Cell Biol.* **50**, 170–180.
- Jin, Y., Wang, Y., Walker, D. L., Dong, H., Conley, C., Johansen, J., and Johansen, K. M. (1999). JIL-1: A novel chromosomal tandem kinase implicated in transcriptional regulation in *Drosophila*. *Mol. Cell* **4**, 129–135.
- Johansen, K. M., and Johansen, J. (2006). Regulation of chromatin structure by histone H3S10 phosphorylation. *Chromosome Res.* **14**, 393–404.
- Judd, B. (1955). Direct proof of a variegated-type position effect at the *white* locus in *Drosophila melanogaster*. *Genetics* **40**, 739–744.
- Karpfen, G. H., and Spradling, A. C. (1992). Analysis of subtelocentric heterochromatin in the *Drosophila* minichromosome *Dp1187* by single P element insertional mutagenesis. *Genetics* **132**, 737–753.
- Kaufmann, B. P. (1942). Reversion from *roughest* to wild type in *Drosophila melanogaster*. *Genetics* **27**, 537–549.
- Kaufmann, B. P. (1946). Organization of the chromosome. I. Break distribution and chromosome recombination in *Drosophila melanogaster*. *J. Exp. Zool.* **102**, 293–320.

- Kellum, R., and Schedl, P. (1992). A group of scs elements function as domain boundaries in an enhancer-blocking assay. *Mol. Cell Biol.* **12**, 2424–2431.
- Kouzarides, T. (2007). Chromatin modifications and their function. *Cell* **128**, 693–705.
- Kuhfittig, S., Szabad, J., Schotta, G., Hoffmann, J., Mathe, E., and Reuter, G. (2001). *pitkin^D*, a novel gain-of-function enhancer of position-effect variegation, affects chromatin regulation during oogenesis and early embryogenesis in *Drosophila*. *Genetics* **157**, 1227–1244.
- Labrador, M., and Corces, V. G. (2002). Setting the boundaries of chromatin domains and nuclear organization. *Cell* **111**, 151–154.
- Lachner, M., O'Carroll, D., Rea, S., Mechtler, K., and Jenuwein, T. (2001). Methylation of histone H3 lysine 9 creates a binding site for HP1 proteins. *Nature* **410**, 116–120.
- Lachner, M., O'Sullivan, R. J., and Jenuwein, T. (2003). An epigenetic road map for histone lysine methylation. *J. Cell Sci.* **116**, 2117–2124.
- Lakhotia, S. C., and Jacob, J. (1974). EM autoradiographic studies on polytene nuclei of *Drosophila melanogaster*. *Exp. Cell Res.* **86**, 253–263.
- Lanctôt, C., Cheutin, T., Cremer, M., Cavalli, G., and Cremer, T. (2007). Dynamic genome architecture in the nuclear space: Regulation of gene expression in three dimensions. *Nat. Rev. Genet.* **8**, 104–115.
- Lawrence, R. J., and Pikaard, C. S. (2004). Chromatin turn ons and turn offs of ribosomal RNA genes. *Cell Cycle* **3**, 880–883.
- Lei, E. P., and Corces, V. G. (2006). RNA interference machinery influences the nuclear organization of a chromatin insulator. *Nat. Genet.* **38**, 936–941.
- Lerach, S., Zhang, W., Bao, X., Deng, H., Girton, J., Johansen, J., and Johansen, K. M. (2006). Loss-of-function alleles of the JIL-1 kinase are strong suppressors of position effect variegation of the *w^{m4}* allele in *Drosophila*. *Genetics* **173**, 2403–2406.
- Lewis, E. B. (1950). The phenomenon of position effect. *Adv. Genet.* **3**, 73–115.
- Lewis, M. S., and Pikaard, C. S. (2001). Restricted chromosomal silencing in nucleolar dominance. *Proc. Natl. Acad. Sci. USA* **98**, 14536–14540.
- Lindsley, D. L., and Zimm, G. G. (1992). "The Genome of *Drosophila Melanogaster*." Academic Press, San Diego.
- Lindsley, D. L., Edington, C., and von Halle, E. S. (1960). Sex-linked recessive lethals in *Drosophila* whose expression is suppressed by the Y chromosome. *Genetics* **45**, 1649–1670.
- Locke, J., Kotarski, M. A., and Tartof, K. D. (1988). Dosage dependent modifiers of position effect variegation in *Drosophila* and a mass action model that explains their effects. *Genetics* **120**, 181–198.
- Lohe, A. R., Hilliker, A. J., and Roberts, P. A. (1993). Mapping simple repeated DNA sequences in heterochromatin of *Drosophila melanogaster*. *Genetics* **134**, 1149–1174.
- Luger, K. (2006). Dynamic nucleosomes. *Chromosome Res.* **14**, 5–16.
- Mattout, A., Goldberg, M., Tzur, Y., Margalit, A., and Gruenbaum, Y. (2006). Specific and conserved sequences in *D. melanogaster* and *C. elegans* lamins and histone H2A mediate the attachment of lamins to chromosomes. *J. Cell. Sci.* **120**, 77–85.
- Matzke, M. A., and Birchler, J. A. (2005). RNAi-mediated pathways in the nucleus. *Nat. Rev. Genet.* **6**, 24–35.
- Mendjan, S., Taipale, M., Kind, J., Holz, H., Gebhardt, P., Schelder, M., Vermeulen, M., Buscaino, K., Duncan, K., Mueller, J., Wilm, M., Stunnenberg, H. G., et al. (2006). Nuclear pore components are involved in the transcriptional regulation of dosage compensation in *Drosophila*. *Mol. Cell* **21**, 811–823.
- Mis, J., Ner, S. S., and Grigliatti, T. A. (2006). Identification of three histone methyltransferases in *Drosophila*: dG9a is a suppressor of PEV and is required for gene silencing. *Mol. Genet. Genomics* **275**, 513–526.

- Misteli, T. (2007). Beyond the sequence: Cellular organization of genome function. *Cell* **128**, 787–800.
- Mito, Y., Henikoff, J. G., and Henikoff, S. (2005). Genome-scale profiling of histone H3.3 replacement patterns. *Nat. Genet.* **37**, 1090–1097.
- Morgan, T. H., Bridges, C. B., and Sturtevant, A. H. (1925). The genetics of *Drosophila melanogaster*. *Biolphilia Genet.* **2**, 262.
- Morgan, T. H., Schultz, J., and Curry, V. (1941). Investigation on the constitution of the germinal material in relation to heredity. *Carnegie Inst. Yearbook* **40**, 282–287.
- Mottus, R., Sobels, R. E., and Grigliatti, T. A. (2000). Mutational analysis of a histone deacetylase in *Drosophila melanogaster*: Missense mutations suppress gene silencing associated with position effect variegation. *Genetics* **154**, 657–668.
- Muller, H. J. (1930). Types of visible variations induced by X-rays in *Drosophila*. *J. Genet.* **22**, 299–334.
- Muller, H. J. (1932). Further studies on the nature and causes of gene mutations. *Proc. Sixth Int. Congr. Genet.* **1**, 213–255.
- Nakayama, T., Nishioka, K., Dong, Y.-X., Shimojima, T., and Hirose, S. (2007). *Drosophila* GAGA factor directs histone H3.3 replacement that prevents the heterochromatin spreading. *Genes Dev.* **21**, 552–561.
- Neely, K. E., and Workman, J. L. (2002). Histone acetylation and chromatin remodeling: Which comes first? *Mol. Genet. Metab.* **76**, 1–5.
- Neuhaus, M. J. (1939). Genetic study of the Y chromosome in *Drosophila melanogster*. *J. Genet.* **37**, 229–254.
- Nishioka, K., Rice, J. C., Sarma, K., Erdjument-Bromage, H., Werner, J., Wang, Y., Sergei Chuikov, P., Valenzuela, P., Tempst, P., Steward, R., Lis, J. T., Allis, C. D., et al. (2002). PR-set7 is a nucleosome-specific methyltransferase that modifies lysine 20 of histone H4 and is associated with silent chromatin. *Mol. Cell* **9**, 1201–1213.
- Pai, C. Y., Lei, E. P., Ghosh, D., and Corces, V. G. (2004). The centrosomal protein CP190 is a component of the gypsy chromatin insulator. *Mol. Cell* **16**, 737–748.
- Pak, D. T., Pflumm, M., Chesnokov, I., Huang, D. W., Kellum, R., Marr, J., Romanowski, P., and Botchan, M. R. (1997). Association of the origin recognition complex with heterochromatin and HP1 in higher eukaryotes. *Cell* **91**, 311–323.
- Pal-Bhadra, M., Leibovitch, B. A., Gandhi, S. G., Rao, M., Bhadra, U., Birchler, J. A., and Elgin, S. C. (2004). Heterochromatic silencing and HP1 localization in *Drosophila* are dependent on the RNAi machinery. *Science* **303**, 669–672.
- Panshin, I. B. (1938). Cytogenic nature of position effect of genes *white* (mottled) and *cubitus interruptus*. *Biol. Zh.* **7**, 837–868.
- Pardue, M. L., and DeBaryshe, P. G. (1999). Telomeres and telomerase: More than the end of the line. *Chromosoma* **108**, 73–82.
- Parkhurst, S. M., Harrison, D. A., Remington, M. P., Spana, C., Kelley, R. L., Coyne, R. S., and Corces, V. G. (1988). The *Drosophila su(hw)* gene, which controls the phenotypic effect of the gypsy transposable element, encodes a putative DNA-binding protein. *Genes Dev.* **2**, 1205–1215.
- Parnell, T. J., Kuhn, E. J., Gilmore, B. L., Helou, C., Wold, M. S., and Geyer, P. K. (2006). Identification of genomic sites that bind the *Drosophila* suppressor of Hairy-wing insulator protein. *Mol. Cell Biol.* **26**, 5983–5993.
- Peng, J. C., and Karpen, G. H. (2006). H3K9 methylation and RNA interference regulate nucleolar organization and repeated DNA stability. *Nat. Cell Biol.* **9**, 25–35.
- Perrin, L., Demakova, O., Fanti, L., Kallenbach, S., Saingery, S., Macl'ceva, N. I., Pimpinelli, S., Zhitulev, I., and Pradel, J. (1998). Dynamics of the sub-nuclear distribution of Modulo and the regulation of position-effect variegation. *J. Cell Sci.* **111**, 2753–2761.

- Peterson, C. L. (2003). Transcriptional activation: Getting a grip on condensed chromatin. *Curr. Biol.* **13**, R195–R197.
- Pickersgill, H., Kalverda, B., de Wit, E., Talhout, W., Fornerod, M., and van Steensel, B. (2006). Characterization of the *Drosophila melanogaster* genome at the nuclear lamina. *Nat. Genet.* **38**, 1005–1014.
- Pimpinelli, S., Berloco, M., Fanti, L., Dimitri, P., Bonaccorsi, S., Marchetti, E., Caizzi, R., Aggese, C. C., and Gatti, M. (1995). Transposable elements are stable structural components of *Drosophila melanogaster* heterochromatin. *Proc. Natl. Acad. Sci. USA* **92**, 3804–3808.
- Pokholok, D. K., Harbison, C. T., Levine, S., Cole, M., Hannett, N. M., Lee, T. I., Bell, G. W., Walker, P. A., Rolfe, P. A., Herbolsheimer, E., Zeitlinger, J., Lewitter, F., et al. (2005). Genome-wide map of nucleosome acetylation and methylation in yeast. *Cell* **122**, 517–527.
- Puckett, L. D., and Snyder, L. A. (1973). Analysis of ribosomal RNA synthesis in two X chromosome inversions in *Drosophila melanogaster*. *Genetics* **74**, S221.
- Qi, D., Jin, H., Lilja, T., and Mannervik, M. (2006). *Drosophila* Reptin and other TIP60 complex components promote generation of silent chromatin. *Genetics* **174**, 241–251.
- Ramos, E., Ghosh, D., Baxter, E., and Corces, V. G. (2006). Genomic organization of *gypsy* chromatin insulators in *Drosophila melanogaster*. *Genetics* **172**, 2337–2349.
- Rea, S., Eisenhaber, F., O'Carroll, D., Strahl, B. D., Sun, Z. W., Schmid, M., Opravil, S., Mechteder, C. P., Ponting, C. P., Allis, C. D., and Jenuwein, T. (2000). Regulation of chromatin structure by site-specific histone H3 methyltransferases. *Nature* **406**, 593–599.
- Reugels, A. M., Kurek, R., Lammermann, U., and Bunemann, H. (2000). Mega-introns in the dynein gene DhDhc7(Y) on the heterochromatic Y chromosome give rise to the giant thread loops in primary spermatocytes of *Drosophila hydei*. *Genetics* **154**, 759–769.
- Reuter, G., and Spierer, P. (1992). Position effect variegation and chromatin proteins. *Bioessays* **14**, 605–612.
- Reuter, G., and Wolff, I. (1981). Isolation of dominant suppressor mutations for position-effect variegation in *Drosophila melanogaster*. *Mol. Gen. Genet.* **182**, 516–519.
- Reuter, G., Werner, W., and Hoffmann, H. J. (1982). Mutants affecting position-effect heterochromatization in *Drosophila melanogaster*. *Chromosoma* **85**, 539–551.
- Reuter, G., Wolff, I., and Friede, B. (1985). Functional properties of the heterochromatic sequences inducing *w^{m4}* position-effect variegation in *Drosophila melanogaster*. *Chromosoma* **93**, 132–139.
- Reuter, G., Dorn, R., Wustmann, G., Friede, B., and Rauh, G. (1986). Third chromosome suppressor of position-effect variegation in *Drosophila melanogaster*. *Mol. Gen. Genet.* **202**, 481–487.
- Reuter, G., Giarre, M., Farah, J., Gausz, J., Spierer, A., and Spierer, P. (1990). Dependence of position-effect variegation in *Drosophila* on the dose of a gene encoding an unusual zinc-finger protein. *Nature* **344**, 219–223.
- Risinger, C., Deiteher, D. L., Lundell, I., Schwartz, T. L., and Larhammar, D. (1997). Complex gene organization of synaptic protein SNAP-25 in *Drosophila melanogaster*. *Gene* **194**, 169–177.
- Rizki, T. M. (1961). Intracellular localization of kynurenine in the fat body of *Drosophila*. *J. Biophys. Biochem. Cytol.* **9**, 567–572.
- Roseman, R. R., Pirrotta, V., and Geyer, P. K. (1993). The su(Hw) protein insulates expression of the *Drosophila melanogaster white* gene from chromosomal position-effects. *EMBO J.* **12**, 435–442.
- Rubin, G. M., and Spradling, A. C. (1982). Genetic transformation of *Drosophila* with transposable element vectors. *Science* **218**, 348–353.
- Rzepecki, R., Bogachev, S. S., Kokozza, E., Stuurman, N., and Fisher, P. A. (1998). *In vivo* association of lamins with nucleic acids in *Drosophila melanogaster*. *J. Cell Sci.* **111**, 121–129.
- Sage, B. T., and Csink, A. K. (2003). Heterochromatic self-association, a determinant of nuclear organization, does not require sequence homology in *Drosophila*. *Genetics* **165**, 1183–1193.

- Sass, G. L., and Henikoff, S. (1999). Pairing-dependent mislocalization of a *Drosophila brown* gene reporter to a heterochromatic environment. *Genetics* **152**, 596–604.
- Schalet, A. (1969). Three Y suppressed phenotypes associated with X chromosomes carrying *In(1)B^{M1}*. *Dros. Inf. Serv.* **44**, 87.
- Schotta, G., Ebert, A., Krauss, V., Fischer, A., Hoffmann, J., Rea, S., Jenuwein, T., Dorn, R., and Reuter, G. (2002). Central role of *Drosophila SU(VAR)3-9* in histone H3-K9 methylation and heterochromatic gene silencing. *EMBO J.* **21**, 1121–1131.
- Schotta, G., Ebert, A., Dorn, R., and Reuter, G. (2003a). Position effect variegation and the genetic dissection of chromatin regulation in *Drosophila*. *Semin. Cell Dev. Biol.* **14**, 67–75.
- Schotta, G., Ebert, A., and Reuter, G. (2003b). *SU(VAR)3-9* is a conserved key function in heterochromatic gene silencing. *Genetica* **117**, 149–158.
- Schotta, G., Lachner, M., Sarma, K., Ebert, A., Sengupta, R., Reuter, G., Reinberg, D., and Jenuwein, T. (2004). A silencing pathway to induce H3-K9 and H4-K20 trimethylation at constitutive heterochromatin. *Genes Dev.* **18**, 1251–1262.
- Schubeler, D., MacAlpine, D. M., Scalzo, D., Wirbelauer, C., Kooperberg, C., van Leeuwen, F., Gottschling, D. E., O'Neill, L. P., Turner, B. M., Delrow, J., Bell, S. P., and Groudine, M. (2004). The histone modification pattern of active genes revealed through genome-wide chromatin analysis of a higher eukaryote. *Genes Dev.* **18**, 1263–1271.
- Schultz, J. (1936). Variegation in *Drosophila* and the inert chromosome regions. *Proc. Natl. Acad. Sci. USA* **22**, 27–33.
- Schultz, J. (1939). The function of heterochromatin. *Proc. Natl. Acad. Sci. USA* **22**, 27–33.
- Schultz, J. (1941). The function of heterochromatin. *Proc. 7th Intern. Congr. Genet.* **25**, 7–262.
- Schultz, J., and Dobzhansky, T. (1934). The relation of a dominant eye color in *Drosophila melanogaster* to the associated chromosome rearrangement. *Genetics* **19**, 344–364.
- Schulze, R. S., and Wallrath, L. L. (2007). Gene regulation by chromatin structure: Paradigms established in *Drosophila melanogaster*. *Annu. Rev. Entomol.* **52**, 171–192.
- Schulze, S. R., Sinclair, D. A., Fitzpatrick, K. A., and Honda, B. M. (2005). A genetic and molecular characterization of two proximal heterochromatic genes on chromosome 3 of *Drosophila melanogaster*. *Genetics* **169**, 2165–2177.
- Schwartz, B. E., and Ahmad, K. (2005). Transcriptional activation triggers deposition and removal of the histone variant H3.3. *Genes Dev.* **19**, 804–814.
- Seum, C., Spierer, A., Pauli, D., Szidonya, J., Reuter, G., and Spierer, P. (1996). Position-effect variegation in *Drosophila* depends on the dose of the gene encoding the E2F transcriptional activator and cell cycle regulator. *Development* **122**, 1949–1956.
- Shaffer, C. D., Stephens, G. E., Thompson, B. A., Funches, L., Bernat, J. A., Craig, C. A., and Elgin, S. C. R. (2002). Heterochromatin protein 2 (HP2), a partner of HP1 in *Drosophila* heterochromatin. *Proc. Natl. Acad. Sci. USA* **99**, 14332–14337.
- Shareef, M. M., King, C., Damaj, M., Badagu, R., Huang, D. W., and Kellum, R. (2001). *Drosophila* heterochromatin protein 1 (HP1)/Origin recognition complex (ORC) protein is associated with HP1 and ORC and functions in heterochromatin-induced silencing. *Mol. Biol. Cell* **12**, 1671–1685.
- Shogren-Knaak, M., Ishii, H., Sun, J.-M., Pazin, M. J., Davie, J. R., and Peterson, C. L. (2006). Histone H4-K16 acetylation controls chromatin structure and protein interactions. *Science* **311**, 844–847.
- Sinclair, D. A. R., Mottus, R. C., and Grigliatti, T. A. (1983). Genes which suppress position-effect variegation in *Drosophila melanogaster* are clustered. *Mol. Gen. Genet.* **191**, 326–333.
- Sinclair, D. A. R., Schulze, S., Silva, E., Fitzpatrick, K. A., and Honda, B. M. (2000). Essential genes in autosomal heterochromatin of *Drosophila melanogaster*. *Genetica* **109**, 9–18.
- Slatis, H. M. (1955). Position effects at the *brown* locus in *Drosophila melanogaster*. *Genetics* **40**, 5–23.

- Somech, R., Shaklai, S., Geller, O., Amariglio, N., Simon, A. J., Rechavi, G., and Gal-Yam, E. N. (2005). The nuclear-envelope protein and transcriptional repressor LAP2 β interacts with HDAC3 at the nuclear periphery, and induces histone H4 deacetylation. *J. Cell Sci.* **118**, 4017–4025.
- Spofford, J. B. (1967). Single-locus modification of position-effect variegation in *Drosophila melanogaster*. I. White variegation. *Genetics* **57**, 751–766.
- Spofford, J. B. (1973). Variegation for *dm* in *Dp(1;3)^{N264-58}*. *Dros. Inf. Serv.* **50**, 98.
- Spofford, J. B. (1976). Position-effect variegation in *Drosophila*. In “The Genetics and Biology of *Drosophila*” (M. Ashburner and E. Novitski, eds.), Vol. 1c, pp. 955–1018. Academic Press, London.
- Spofford, J. B., and DeSalle, R. (1991). Nucleolus organizer-suppressed position-effect variegation in *Drosophila melanogaster*. *Genet. Res.* **57**, 245–255.
- Spradling, A. C., and Karpen, G. H. (1990). Sixty years of mystery. *Genetics* **126**, 779–784.
- Spradling, A. C., and Rubin, G. M. (1982). Transposition of cloned P elements into *Drosophila* germ line chromosomes. *Science* **218**, 341–347.
- Steller, H., and Pirrotta, V. (1985). Expression of the *Drosophila white* gene under the control of the hsp-70 heat shock promoter. *EMBO J.* **4**, 3765–3772.
- Strahl, B. D., and Allis, C. D. (2000). The language of covalent histone modifications. *Nature* **403**, 1–45.
- Sturtevant, A. H. (1925). The effect of unequal crossing over at the *Bar* locus in *Drosophila*. *Genetics* **10**, 117–147.
- Sun, F. L., and Elgin, S. C. (1999). Putting boundaries on silence. *Cell* **99**, 459–462.
- Sun, X., Wahlstrom, J., and Karpen, G. (1997). Molecular structure of a functional *Drosophila* centromere. *Cell* **91**, 1007–1019.
- Swaminathan, J., Baxter, E. M., and Corces, V. G. (2005). The role of histone H2Av variant replacement and histone H4 acetylation in the establishment of *Drosophila* heterochromatin. *Genes Dev.* **19**, 65–76.
- Taddei, A., and Gasser, S. M. (2004). Multiple pathways for telomere tethering: Functional implications of subnuclear position for heterochromatin formation. *Biochim. Biophys. Acta* **1677**, 120–128.
- Talbert, P. B., and Henikoff, S. (2000). A reexamination of spreading of position-effect variegation in the *white-roughest* region of *Drosophila melanogaster*. *Genetics* **154**, 259–272.
- Talbert, P. B., LeCiel, C. D. S., and Henikoff, S. (1994). Modifications of the *Drosophila* heterochromatic mutation *brown Dominant* by linkage alterations. *Genetics* **136**, 559–571.
- Tartof, K. D., Hobbs, C., and Jones, M. (1984). A structural basis for variegating position effects. *Cell* **37**, 869–878.
- Teixeira, M. T., Dujon, B., and Fabre, E. (2002). Genome-wide nuclear morphology screen identifies novel genes involved in nuclear architecture and gene-silencing in *Saccharomyces cerevisiae*. *J. Mol. Biol.* **321**, 551–561.
- Thakar, R., and Csink, A. K. (2005). Changing chromatin dynamics and nuclear organization during differentiation in *Drosophila* larval tissue. *J. Cell Sci.* **118**, 951–960.
- Tobler, J., Bowman, J. T., and Simmons, J. B. (1971). Gene modulation in *Drosophila*: Dosage compensation and relocated v⁺ genes. *Biochem. Genet.* **5**, 11–117.
- Tschiersch, B., Hofmann, A., Krauss, V., Dorn, R., Korge, G., and Reuter, G. (1994). The protein encoded by the *Drosophila* position-effect variegation suppressor gene *Su(var)3-9* combines domains of antagonistic regulators of homeotic gene complexes. *EMBO J.* **13**, 3822–3831.
- Tumbar, T., Sudlow, G., and Belmont, A. S. (1999). Large-scale chromatin unfolding and remodeling induced by VP16 acidic activation domain. *J. Cell Biol.* **145**, 1341–1354.
- Turner, B. M. (2000). Histone acetylation and an epigenetic code. *Bioessays* **22**, 836–845.
- Turner, B. M. (2002). Cellular memory and the histone code. *Cell* **111**, 285–291.

- Turner, B. M., Birley, A. J., and Lavender, J. (1992). Histone H4 isoforms acetylated at specific lysine residues define individual chromosomes and chromatin domains in *Drosophila* polytene nuclei. *Cell* **69**, 375–384.
- van Driel, R., Fransz, P. F., and Verschure, P. J. (2003). The eukaryotic genome: A system regulated at different hierarchical levels. *J. Cell Sci.* **116**, 4067–4075.
- Verdel, A., and Moazed, D. (2005). RNAi-directed assembly of heterochromatin in fission yeast. *FEBS Lett.* **579**, 5872–5878.
- Verdel, A., Jia, S., Gerber, S., Sugiyama, T., Gygi, S., Grewal, S. I., and Moazed, D. (2004). RNAi-mediated targeting of heterochromatin by the RITS complex. *Science* **303**, 672–676.
- Verni, F., Gandhi, R., Goldberg, M. L., and Gatti, M. (2000). Genetic and molecular analysis of *wings apart-like* (*wapl*), a gene controlling heterochromatin organization in *Drosophila melanogaster*. *Genetics* **154**, 1693–1710.
- Wade, P. A., Pruss, D., and Wolffe, A. P. (1997). Histone acetylation: Chromatin in action. *Trends Biochem. Sci.* **22**, 128–132.
- Wakimoto, B., and Hearn, M. G. (1990). The effect of chromosome rearrangements on the expression of heterochromatic genes in chromosome 2L of *Drosophila melanogaster*. *Genetics* **125**, 141–154.
- Wallrath, L. L. (2000). *Drosophila* telomeric transgenes provide insights on mechanisms of gene silencing. *Genetica* **109**, 25–33.
- Wallrath, L. L., and Elgin, S. C. R. (1995). Position effect variegation in *Drosophila* is associated with an altered chromatin structure. *Genes Dev.* **9**, 1263–1277.
- Wallrath, L. L., Guntur, V. P., Rosman, L. E., and Elgin, S. C. (1996). DNA representation of variegating heterochromatic P-element inserts in diploid and polytene tissues of *Drosophila melanogaster*. *Chromosoma* **104**, 519–527.
- Walter, M. F., Jang, C., Kasravi, B., Donath, J., Mechler, J. M., and Biessmann, H. (1995). DNA organization and polymorphism of a wild-type *Drosophila* telomere region. *Chromosoma* **104**, 229–241.
- Wang, Y., Zhang, W., Jin, Y., Johansen, J., and Johansen, K. M. (2001). The JIL-1 tandem kinase mediates histone H3 phosphorylation and is required for maintenance of chromatin structure in *Drosophila*. *Cell* **105**, 433–443.
- Wang, L., Haeusler, R. A., Good, P. D., Thompson, M., Nagar, S., and Engelke, D. R. (2005). Silencing near rRNA genes requires nucleolar localization. *J. Biol. Chem.* **280**, 8637–8639.
- Wargent, J. M. (1972). Position-effect variegation in the *Revolute of Bridges* strain of *Drosophila melanogaster*. *Dros. Inform. Serv.* **49**, 50–51.
- Wei, G. H., Liu, E. P., and Liang, C. C. (2005). Chromatin domain boundaries: Insulators and beyond. *Cell Res.* **15**, 292–300.
- Weiler, K. S., and Wakimoto, B. T. (1995). Heterochromatin and gene expression in *Drosophila*. *Annu. Rev. Genet.* **29**, 577–605.
- West, A. G., Gaszner, M., and Felsenfeld, G. (2002). Insulators: Many functions, many mechanisms. *Genes Dev.* **16**, 271–288.
- Workman, J. L., and Kingston, R. E. (1998). Alteration of nucleosome structure as a mechanism of transcriptional regulation. *Annu. Rev. Biochem.* **67**, 545–579.
- Wustmann, G., Szidonya, J., Taubert, H., and Reuter, G. (1989). The genetics of position-effect modifying loci in *Drosophila melanogaster*. *Mol. Gen. Genet.* **217**, 520–527.
- Yamaguchi, M., Yoshida, H., Hirose, F., Inoue, Y. H., Hayashi, Y., Yamagishi, M., Nishi, Y., Tamai, K., Sakaguchi, K., and Matsukage, A. (2001). Ectopic expression of BEAF32A in the *Drosophila* eye imaginal disc inhibits differentiation of photoreceptor cells and induces apoptosis. *Chromosoma* **110**, 313–321.

- Yamamoto, K., and Sonoda, M. (2003). Self-interaction of heterochromatin protein 1 is required for direct binding to histone methyltransferase, SUV39H1. *Biochem. Biophys. Res. Commun.* **301**, 287–292.
- Yamamoto, Y., Girard, F., Bello, B., Affolter, M., and Gehring, W. J. (1997). The cramped gene of *Drosophila* is a member of the Polycomb-group, and interacts with *mus209*, the gene encoding Proliferating Cell Nuclear Antigen. *Development* **124**, 3385–3394.
- Yang, X. J. (2004). Lysine acetylation and the bromodomain: A new partnership for signaling. *Bioessays* **26**, 1076–1087.
- Yasufuji, T. M., Tagami, H., Nakatani, Y., and Felsenfeld, G. (2004). CTCF tethers an insulator to subnuclear sites, suggesting shared insulator mechanisms across species. *Mol. Cell* **13**, 291–298.
- Zaratiegui, M., Irvine, D. V., and Martienssen, R. A. (2007). Noncoding RNAs and gene silencing. *Cell* **128**, 763–776.
- Zhang, Y., and Reinberg, D. (2001). Transcription regulation by histone methylation: Interplay between different covalent modifications of the core histone tails. *Genes Dev.* **15**, 2343–2360.
- Zhang, W., Jin, Y., Ji, Y., Girton, J., Johansen, J., and Johansen, K. M. (2003). Genetic and phenotypic analysis of alleles of the *Drosophila* chromosomal JIL-1 kinase reveals a functional requirement at multiple developmental stages. *Genetics* **165**, 1341–1354.
- Zhang, W., Deng, H., Bao, X., Lerach, S., Girton, J., Johansen, J., and Johansen, K. M. (2006). The JIL-1 histone H3S10 kinase regulates dimethyl histone H3K9 modifications and heterochromatic spreading in *Drosophila*. *Development* **133**, 229–235.
- Zhao, K., Hart, C. M., and Laemmli, U. K. (1995). Visualization of chromosomal domains with boundary element-associated factor BEAF-32. *Cell* **81**, 879–889.
- Zhimulev, I. F. (1998). Polytene chromosomes, heterochromatin, and position effect variegation. *Adv. Genet.* **37**, 1–566.
- Zhimulev, I. F., and Belyaeva, E. S. (2003). Intercalary heterochromatin and genetic silencing. *BioEssays* **25**, 1040–1051.
- Zuckerkandl, E. (1974). A possible role of “inert” heterochromatin in cell differentiation. *Biochemie* **56**, 937–954.