

# CROWN ROT

*A Serious Disease of Hosta and Other Ornamentals*



**H**ostas are the top-selling herbaceous perennials nationwide thanks to their attractive foliage, endless diversity of shape and size, tolerance of shady areas, and minimal maintenance needs.

Few pests plague hostas in Iowa. Since the mid-1990s, however, a disease called crown rot has damaged many hosta plantings. This fungal disease, caused by *Sclerotium rolfsii*, can severely damage established hostas within a week, and is difficult to eradicate. Formerly restricted to warm, humid states in the southern United States, crown rot is now common in many Midwest gardens.

This bulletin explains how *S. rolfsii* managed to spread into the Midwest, how it causes crown rot in hostas, and ways to manage the disease.







## Symptoms

Symptoms begin to appear on hosta after prolonged hot, humid weather. The lower leaves begin to turn yellow, then brown, and wilt from the margins back toward the base (Figures 1, 2, and 3). The upper leaves may soon collapse, too. The wilted leaves can be easily pulled from the crown, because they have been attacked at the base of the petiole. The bases of these damaged petioles show a brown discoloration and mushy texture (Figure 4). Plants with less succulent stems, such as peony, are girdled at the base of stems, causing discolored and wilted leaves, but the stems may not collapse (Figure 5). Fluffy white threads (mycelium) of the crown rot fungus typically are present on the rotted tissue and surrounding soil (Figure 6). A closer look shows small spheres, about the size of mustard seeds, sprinkled on the soil (Figure 7). These tiny spheres, called sclerotia, allow the fungus to survive cold winters and other unfavorable conditions. As sclerotia mature, their color changes from white to a light tan or reddish brown.



**FIGURE 1**  
*A hosta bed showing the leaf yellowing typical of crown rot.*



**FIGURE 4**  
*Brown, softened tissue at the base of hosta petioles. Tiny, orange spheres on the soil surface below the petioles (see arrow) are sclerotia of the fungus *Sclerotium rolfsii*. Note penny for scale.*





# C R O W N   R O T



**FIGURE 2**  
*Close-up of a hosta with marginal yellowing and browning caused by crown rot.*



**FIGURE 3**  
*Collapse and death of lower leaves of hosta, caused by crown rot attack at the bases of the leaf petioles.*



**FIGURE 5**  
*A wilted, collapsed peony (center) whose fronds were attacked at the base by *Sclerotium rolfsii*. The source of the fungus was a nearby, infected hosta.*



**FIGURE 6**

White mycelium (fungal strands) of *Sclerotium rolfsii* on the base of a hosta petiole severed by the fungus.

**FIGURE 7**

Sclerotia of *Sclerotium rolfsii*. Sclerotia are approximately 1 mm in diameter (about the size of mustard seeds) and vary in color from white to brick red.




### How *S. Rolfsii* Causes Crown Rot

The crown rot life cycle begins with the germination of sclerotia (Figure 8). Mycelium fans out in all directions from the sclerotia, slowly growing across the surface of the soil in warm, moist weather. When a host plant is nearby,

the fungus exudes droplets containing oxalic acid and tissue-destroying enzymes. Oxalic acid poisons plants, causing their cell walls to break down.

# C R O W N   R O T



**FIGURE 8**  
Disease cycle of crown rot on hosta.

When *S. rolf sii*'s mycelium comes into contact with hosta, the oxalic acid kills the petioles at the base. With their support gone, the leaves begin to collapse. The stem bases, and the soil for several inches around the crown, can be covered with fluffy white mycelium. Mature mycelium begins to produce sclerotia. There is no required dormancy period, so newly formed sclerotia can produce new mycelium at once or remain dormant in the soil for years.

Sclerotia are hard-shelled survival structures. It was once thought that the cold winters of the Upper Midwest

would kill *S. rolf sii*, but gardeners have seen crown rot spread through their plantings year after year. It now appears that *S. rolf sii* sclerotia can survive temperatures well below freezing, either in plant debris or at the soil surface.

On the positive side, *S. rolf sii* seldom produces microscopic spores like most other fungi. Without spores, it cannot spread by moving on air currents, but only as sclerotia in soil or plants, or by mycelium that grows slowly across the ground from plant to plant.



TABLE 1

**Genera of herbaceous perennials known to be hosts for *Sclerotium rolfsii*. The plants on this list grow well in part sun to full shade in the Upper Midwest.**

**Names****Scientific–Common**

*Ajuga*–bugleweed  
*Ageratum*– chick weed, bluemink  
*Anemone*–windflower  
*Aquilegia*–columbine  
*Arrhenatherum*–oat grass  
*Asplenium*–dragontail fern  
*Begonia* –begonia  
*Caladium*–fancy-leaved caladium  
*Campanula*–bellflower  
*Chrysogonum*–goldenstar  
*Convallaria*–lily-of-the-valley  
*Dicentra*–bleeding heart  
*Digitalis*–foxglove  
*Dryopteris*–wood fern  
*Forsythia*–forsythia  
*Hemerocallis*–daylily  
*Hosta*–plantain lily, hosta  
*Hydrangea*– hydrangea  
*Lysimachia*–loosestrife  
*Petunia* –petunia  
*Phlox*–phlox  
*Ranunculus*–buttercup  
*Tagetes* –marigold,  
*Viburnum*– viburnum  
*Vinca*–periwinkle  
*Viola*–violet, pansy  
*Zinnia* –zinnia.

For a more complete listing of host genera, consult the following reference: Farr, D.F., & Rossman, A.Y. Fungal Databases, Systematic Mycology and Microbiology Laboratory, ARS, USDA from <http://nt.ars-grin.gov/fungaldatabases/>

## Spread of Disease

What are the circumstances behind crown rot's introduction into northern landscapes? The answer is unknown, but one possibility is that *S. rolfsii* has been spread on contaminated nursery plants (Figure 8). Others point to the nonstop exchange of hostas among gardeners throughout the United States. Once crown rot has entered a planting it is easily spread by sclerotia clinging to soil on shoes, tools, and plant material.

*S. rolfsii* can affect more than 500 different plant genera, including ornamentals, fruits, and vegetables. Table 1 lists genera of ornamental plants susceptible to crown rot and, like hosta, adapted to partial to full shade environments. Many susceptible ornamentals, such as daylily (*Hemerocallis*), impatiens, ajuga, peony, and vinca, are often planted in the same beds as hosta, so the disease can move from hosta to neighboring plants.

## Management

Effective management of crown rot requires combining several strategies. The most effective management practices are a combination of cultural tactics such as careful inspection before planting, sanitation to control spread, and choosing plants believed to be less susceptible (see Table 2). These methods will not cure infected hostas but can slow or stop the spread of disease.

### Cultural Management of Crown Rot in the Landscape

Avoidance is one of the best crown rot management tactics. Simply put, no fungus means no disease. But how do you keep *S. rolfsii* away from your hostas?

TABLE 2

**Genera of herbaceous perennials not known to be hosts for *Sclerotium rolfsii*. The plants on this list grow well in part sun to full shade in the Upper Midwest.**

*Acanthus*–bear's breech  
*Aconitum*–monkshood  
*Actaea*–baneberry  
*Adiantum*–maidenhair fern  
*Alchemilla*–lady's mantle  
*Anaphilis*–pearly everlasting  
*Arunacus*–goatsbeard  
*Astilbe*–false spirea  
*Cimicifuga*–bugbane  
*Corydalis*–corydalis  
*Disporum*–fairy-bells  
*Epimedium*–barrenwort  
*Erythronium*–trout lily  
*Filipendula*–meadowsweet  
*Gaura*–gaura  
*Gentiana*–gentian  
*Geranium*–cranesbill  
*Hakonechloa*–hakone grass  
*Helleborus*–hellebore  
*Houttuynia*–chameleon plant  
*Lamiastrum*–yellow archangel  
*Lamium*–dead nettle  
*Mertensia*–bluebells  
*Myosotis*–forget-me-not  
*Osmunda*–royal fern  
*Papaver*–poppy  
*Polemonium*–Jacob's ladder  
*Polystichum*–shield fern  
*Primula*–primrose  
*Pulmonaria*–lungwort  
*Pulsatilla*–pasque flower  
*Sanguinaria*–bloodroot  
*Stachys*–lamb's ear  
*Thalictrum*–meadow rue  
*Tiarella*–foamflower  
*Trillium*–wake-robין  
*Trollius*–globeflower



One key to avoiding crown rot is careful inspection. Before purchasing or transplanting, examine plants for yellow, wilted lower leaves, sclerotia on the soil surface, softening and browning at the bases of petioles, and white mycelium around damaged tissue. Checking for these symptoms will reduce the risk of *S. rolf sii* entering the garden.

If the problem is already present in landscape plantings, the contaminated area containing the infected hostas and soil should be quarantined, and care should be used to avoid spreading any soil or plant material outside of this zone. You can excavate soil in the contaminated area to a depth of at least 8 inches, discard the soil, and replace it with uncontaminated soil. But the time, labor, and expense of this tactic is discouraging, especially when the contaminated area is more than a few square feet in size. Do not transplant from the contaminated area, because sclerotia clinging to the roots and crown can start a new infection cycle in another part of the garden. All tools and implements should be washed thoroughly, and the wash water should go back into the contaminated bed. Some recommendations call for dipping tools in a 10 percent bleach solution for a few minutes to kill *S. rolf sii*, but bleach is corrosive to tools and messy to work with. More important than bleaching is to scrub tools clean of all adhering soil. If you decide to use a bleach dip, clean your tools thoroughly beforehand. Other non-corrosive disinfectants are available for tool disinfection, including products with hydrogen peroxide, quaternary ammonium or ethanol 70%.

Many growers mulch their hosta beds to help provide the consistently moist soil conditions that hostas prefer. Since the mycelium of *S. rolf sii* can grow rampantly through moist mulch in warm weather, it may be helpful to maintain a mulch-free zone several inches wide around hosta crowns if crown rot has appeared in the planting. Some hosta growers also recommend planting crowns as high as

possible, with soil barely covering the roots, to reduce the odds that the vulnerable leaf petioles will come into contact with *S. rolf sii* from the soil.

Switching to plants that are not susceptible to *S. rolf sii* is another option. The problem is that so many types of plants are affected by crown rot that you need to choose carefully. Some ornamental plants not known to succumb to *S. rolf sii*, but adapted to partial-shade to full-shade environments in the Upper Midwest, are listed in Table 2. Many of the plants in Table 2, although not confirmed to be hosts of *S. rolf sii*, have not been tested for susceptibility to the fungus. Therefore, it is advisable to try them out on a limited basis before committing large numbers of plants to replace crown rot-killed hostas.

Many people have observed that some types of hostas seem to fare better than others against crown rot. However, crown rot resistance ratings are currently unavailable for hosta species and cultivars.

### ***S. Rolf sii* Problems in Propagation and Production**

A key to avoiding crown rot during plant propagation and production is careful inspection. Regularly examine plants for telltale leaf yellowing and collapse, sclerotia, softening and browning at the bases of petioles, and white mycelium around damaged tissue. Experienced hosta growers know that leaves can turn yellow for other reasons, such as heat stress or a fungal disease called *Rhizoctonia* root rot, but only *S. rolf sii* crown rot will show the other symptoms as well. The Plant Insect Diagnostic Clinic at Iowa State University can help you confirm a diagnosis. A small fee is assessed for this service. Visit the clinic website for current services, submission forms and fees at [clinic.ipm.iastate.edu](http://clinic.ipm.iastate.edu). Once you are convinced that the problem is crown rot, discard the symptomatic plants as well as the planting media and the pot.



# C R O W N R O T



Pre-plant fumigation of production fields used to be an alternative to minimize problems with *S. rolfsii*. However, fumigation is expensive, and the products containing methyl bromide have been phased out due to human and environmental toxicity concerns. Fumigation is not recommended for the home garden because these materials are dangerous to humans.

Fungicides are sometimes used to suppress *S. rolfsii* in hosta as well as other ornamental and food crops. Fungicides containing pentachloronitrobenzene (also known as quintozone or PCNB) typically are applied preventively to soil or growing media. Trade names of PCNB products labeled for control of crown rot on hosta include Terraclor<sup>®</sup>.

Note that the label of this product do not specifically mention hosta, but they are legal to use on hosta because hosta is included within the broad designation “ornamentals” on the label. Since phytotoxic reactions to PCNB can occur, it is advisable to treat a small bed area or a small number of pots first, then check the hosta’s reaction, before attempting larger-scale treatments. To our knowledge, PCNB products labeled for use on ornamentals are available only through commercial pesticide dealers rather than at garden centers or other retail outlets. Sanitizing agents such as chlorine bleach have been used against crown rot, but these products are not legally labeled for this use.

---

*Prepared by* Brooke Edmunds, research assistant, and Mark Gleason, Extension plant pathologist and professor, Department of Plant Pathology; and Ursula Schuch, assistant professor, Department of Horticulture, Iowa State University. Revised by Lina Rodriguez Salamanca, extension plant pathologist and diagnostician with the Plant and Insect Diagnostic Clinic, Iowa State University.

*Designed by* Mary Sailer, Spring Valley Studio.

*Photo Credits:* Figures 1–7: Extension Plant Pathology, Iowa State University. Figure 8: Nita Upchurch.

## IOWA STATE UNIVERSITY Extension and Outreach

This institution is an equal opportunity provider. For the full non-discrimination statement or accommodation inquiries, go to [www.extension.iastate.edu/diversity/ext](http://www.extension.iastate.edu/diversity/ext).