

FINITE TOPOLOGY SELF-TRANSLATING SURFACES FOR THE MEAN CURVATURE FLOW IN \mathbb{R}^3

JUAN DÁVILA, MANUEL DEL PINO, AND XUAN HIEN NGUYEN

ABSTRACT. Finite topology self translating surfaces to mean curvature flow of surfaces constitute a key element for the analysis of Type II singularities from a compact surface, since they arise in a limit after suitable blow-up scalings around the singularity. We find in \mathbb{R}^3 a surface M orientable, embedded and complete with finite topology (and large genus) with three ends asymptotically paraboloidal, such that the moving surface $\Sigma(t) = M + te_z$ evolves by mean curvature flow. This amounts to the equation $H_M = \nu \cdot e_z$ where H_M denotes mean curvature, ν is a choice of unit normal to M , and e_z is a unit vector along the z -axis. The surface M is in correspondence with the classical 3-end Costa-Hoffmann-Meeks minimal surface with large genus, which has two asymptotically catenoidal ends and one planar end, and a long array of small tunnels in the intersection region resembling a periodic Scherk surface. This example is the first non-trivial one of its kind, and it suggests a strong connection between this problem and the theory of embedded, complete minimal surfaces with finite total curvature.

1. INTRODUCTION

We say that a family of orientable, embedded hypersurfaces $\Sigma(t)$ in \mathbb{R}^{n+1} evolves by mean curvature if each point of $\Sigma(t)$ moves in the normal direction with a velocity proportional to its mean curvature at that point. More precisely, there is a smooth family of diffeomorphisms $Y(\cdot, t) : \Sigma(0) \rightarrow \Sigma(t)$, $t > 0$, determined by the mean curvature flow equation

$$(1.1) \quad \frac{\partial Y}{\partial t} = H_{\Sigma(t)}(Y)\nu(Y)$$

where $H_{\Sigma(t)}(Y)$ designates the mean curvature of the surface $\Sigma(t)$ at the point $Y(y, t)$, $y \in \Sigma(0)$, namely the trace of its second fundamental form, ν is a choice of unit normal vector.

The mean curvature flow is one of the most important examples of parabolic geometric evolution of manifolds. Relatively simple in form, it generates a wealth of interesting phenomena which is so far only partly understood. Extensive, deep studies on the properties of this equation have been performed in the last 25 years or so. We refer for instance the reader to the surveys [3] and [30].

A classical, global-in-time definition of a weak solution to mean curvature flow is due to Brakke. These solutions typically develop finite time singularities. When they arise, the evolving manifold loses smoothness, and a change of topology of the surface may occur as the singular time is crossed.

The basic issue of the theory for the mean curvature flow, is to understand the way singularities appear and to achieve an accurate description of the topology of the surface obtained after blowing-up the manifold around the singularity.

Singularities are usually classified as types I and II. If T is a time when a singularity appears, type I roughly means that the curvatures grow no faster than $(T - t)^{-\frac{1}{2}}$. In such a case, a blowing-up procedure, involving a time dependent scaling and translation leads in the limit to a “self shrinking” ancient solution, as established by Huisken in [15]. The appearance of these singularities turns out to be generic under suitable assumptions, see Colding and Minicozzi [4].

Instead, if the singularity is not of type I, it is called type II. In that case, a suitable normalization leads in the limit to an *eternal solution* to MC flow. See Colding and Minicozzi [5], Huisken and Sinestrari [16, 17]. An eternal solution to (1.1) is one that is defined at all times $t \in (-\infty, \infty)$.

The simplest type of eternal solutions are the *self-translating solutions*, surfaces that solve (1.1), do not change shape and travel at constant speed in some specific direction. A self-translating solution of the mean curvature flow (1.1), with speed $c > 0$ and direction $\mathbf{e} \in S^{N-1}$ is a hypersurface of the form

$$(1.2) \quad \Sigma(t) = cte + \Sigma(0),$$

that satisfies (1.1). Equivalently, such that

$$(1.3) \quad H_{\Sigma(0)} = c\mathbf{e} \cdot \nu.$$

This problem is nothing but the minimal surface equation in case $c = 0$. A result by Hamilton [11] states that in the case of a compact convex surface, the limiting scaled singularity does indeed take place in the form of a self-translating solution. This fact makes apparent the importance of eternal self-translating solutions in the understanding of singularity formation. On the other hand, the result in [11] is not known without some convexity assumptions. An open, challenging issue is to understand whether or not a given “self-translator” (convex or non-convex) can arise as a limit of a type II singularity for (1.1).

A situation in which strong insight has been obtained is the mean convex scenario (namely, surfaces with non-negative mean curvature, a property that is preserved under the flow). In fact under quite general assumptions, mean convexity in the singular limit becomes full convexity for the blown-up surface, as it has been established by B. White [34, 35], and by Huisken and Sinestrari [16, 17].

In spite of their importance in the theory for the mean curvature flow, relatively few examples of self-translating solutions are known, and a theory for their understanding, even in special classes is still far from achieved. Since for $c = 0$, equation (1.3) reduces to the minimal surface equation, it is natural to look for analogies with minimal surface theory in order to obtain new nontrivial examples. On the other hand, it has been proven by Ilmanen [21, 22], that the genus of a surface is nonincreasing along the mean curvature flow. Therefore, self-translators originated from a singularity in the flow of a compact surface must have finite genus, or **finite topology**.

The purpose of this paper is to construct new examples of self-translating surfaces to the mean curvature flow with the finite topology in \mathbb{R}^3 . More precisely, we are

interested in tracing a parallel between the theory of embedded, complete minimal surfaces in with finite total Gauss curvature (which are precisely those with finite topology) and self-translators with positive speed. Before stating our main result, we recall next some classical examples of self-translators.

If $\Sigma(t) = \Sigma(0) + ct e_{n+1}$ is a **travelling graph**, namely

$$\Sigma(0) = \{(x, x_{n+1}) / x_{n+1} = F(x), x \in \Omega \subset \mathbb{R}^n\}$$

then equation (1.3) reduces to the elliptic PDE for F ,

$$(1.4) \quad \nabla \cdot \left(\frac{\nabla F}{\sqrt{1 + |\nabla F|^2}} \right) = \frac{c}{\sqrt{1 + |\nabla F|^2}} \quad \text{in } \Omega \subset \mathbb{R}^n.$$

For instance for $n = 1$ and $c = 1$, Grayson [10] found the explicit solution, so-called the *grim reaper* curve \mathcal{G} , given by the graph

$$(1.5) \quad x_2 = F(x_1) = -\log(\cos x_1), \quad x_1 \in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right).$$

In other words, $\Sigma(t) = \mathcal{G} + te_2$ solves (1.1).

For dimensions $n \geq 2$, there exist entire convex solutions to Equation (1.4). Altschuler and Wu [1] found a radially symmetric convex solution $F(|x|)$ to (1.4) by blowing-up a type II singularity of mean curvature flow. This solution can be explicitly found by solving the radial PDE (1.4) which becomes simply

$$(1.6) \quad \frac{F''}{1 + (F')^2} + (n-1) \frac{F'}{r} = c.$$

See [2] and [8]. The resulting surface is asymptotically a paraboloid: at main order, when $c = 1$, it has the behavior

$$(1.7) \quad F(r) = \frac{r^2}{2(n-1)} - \log r + O(r^{-1}) \quad \text{as } r \rightarrow +\infty.$$

We shall denote by \mathcal{P} the graph of this entire graphical self-translator (which is unique up to an additive constant) which we shall refer to as the **travelling paraboloid**. Of course this means that $\Sigma(t) = \mathcal{P} + te_{n+1}$ solves (1.1).

Xu-Jia Wang [33] proved that for $n = 2$, solutions of (1.4) are necessarily radially symmetric about some point, thus in particular they are convex. Surprisingly, for dimensions $n \geq 3$, Wang was able to construct nonradial convex solutions of (1.4).

In dimension $n + 1$, $n \geq 2$, Angenent and Velázquez [2] constructed an axially symmetric solution to (1.1) that develops a type II singularity with a tip that blows-up precisely into the paraboloid \mathcal{P} . On the other hand, B. White proved that the convex surface in \mathbb{R}^{n+1} given by the $\mathcal{G} \times \mathbb{R}^{n-1}$ where \mathcal{G} is the grim reaper curve (1.5), cannot arise as a blow-up of a type II singularity for (1.1).

A non-graphical, two-end axially symmetric self translating solutions of (1.1) for $n \geq 2$ has been found by direct integration of the radial PDE (1.4) by Clutterbuck, Schnurer and Schulze [8]. It can be described as as follows:

Given any number $R > 0$, there is a self-translating solution of (1.1)

$$\Sigma(t) = \mathcal{W} + te_{n+1}$$

where \mathcal{W} is a two-end smooth surface of revolution of the form

$$\mathcal{W} = \mathcal{W}^+ \cup \mathcal{W}^-, \quad \mathcal{W}^\pm = \{(x, x_{n+1}) / x_{n+1} = F^\pm(|x|)\}$$

where the functions $F^\pm(r)$ solve (1.6) for $c = 1$ and $r > R$, with $F^-(r) < F^+(r)$ and $F^+(R) = F^-(R)$. It is shown in [8] that the functions F^\pm have the asymptotic behavior (1.7) of \mathcal{P} up to an additive constant. See Figure 2. We call the two-end translating surface \mathcal{W} the **travelling catenoid**. The reason is natural: when $c = 0$ equation (1.6) is nothing but the minimal surface equation for an axially symmetric minimal surface around the x_{n+1} -axis. When $n = 3$ the equation leads (up to translations) to the plane $x_3 = 0$, or the standard catenoid $r = \cosh(x_3)$. The catenoid is exactly the parallel to \mathcal{W} . The plane is actually in correspondence with the paraboloid \mathcal{P} .

These simple, however important examples, are the only ones available with finite topology. On the other hand, the third author has constructed self translating surfaces with infinite topology, periodic in one direction in [25, 26, 27].

Embedded minimal surfaces of finite total curvature in \mathbb{R}^3 . The theory of embedded, minimal surfaces of finite total curvature in \mathbb{R}^3 has reached a spectacular development in the last 30 years or so. For about two centuries, only two examples of such surfaces were known: the plane and the catenoid. The first nontrivial example was found in 1981 by C. Costa [6, 7]. The Costa surface is a genus one minimal surface, complete and properly embedded, which outside a large ball has exactly three components (its ends), two of which are asymptotically catenoids with the same axis and opposite directions, the third one asymptotic to a plane perpendicular to that axis. Hoffman and Meeks [12, 13, 14] built a class of three-end, embedded minimal surfaces, with the same look as Costa's far away, but with an array of tunnels that provides arbitrary genus k . These are known as the Costa-Hoffman-Meeks surfaces, see Figure 1. Many other examples of multiple-end embedded minimal surfaces have been found since.

All surfaces of this kind are constituted, away from a compact region, by the disjoint union of ends ordered along one coordinate axis, which are asymptotic to planes or to catenoids with parallel symmetry axes, as established by Osserman [28], Schoen [29] and Jorge and Meeks [23]. The topology of such a surface is thus characterized by the genus of a compact region and the number of ends, having therefore finite topology.

Main result: the travelling CHM surface of large genus. In what follows we restrict ourselves to the the case $n + 1 = 3$.

Our purpose is to construct new complete and embedded surfaces in \mathbb{R}^3 which are self translating under mean curvature flow. After a rotation and dilation we can assume that $c = 1$ and that the travelling direction is the that of the positive x_3 -axis. Thus we look for orientable, embedded complete surfaces M in \mathbb{R}^3 satisfying the equation

$$(1.8) \quad H_M = e_z \cdot \nu.$$

where $e_z = e_3$. In other words, the moving surface $\Sigma(t) = M + te_z$ satisfies equation (1.1). A major difficulty to extend the theory of finite total curvature minimal surfaces in Euclidean 3d space to equation (1.8) is that much of the theory developed relies in the powerful tool given by the Weierstrass representation formula, which is not available in our setting. Unlike the static case, the travelling catenoid for instance is not asymptotically flat and does not have total finite total Gauss curvature.




FIGURE 1. Costa-Hoffman-Meeks surface (from www.indiana.edu/~minimal/)

What we establish in our main result is the existence of a 3-end surface M that solves (1.8), homeomorphic to a Costa-Hoffmann-Meeks surface with large genus, whose ends behave like those of a travelling catenoid and a paraboloid.

More precisely, let us consider the union of a travelling paraboloid \mathcal{P} and a travelling catenoid \mathcal{W} , which intersect transversally on a circle C_ρ for some $\rho > 0$. See Figure 2.

Our surface looks outside a compact set like $\mathcal{P} \cup \mathcal{W}$ in Figure 2, while near the circle C_ρ the look is that of the static CHM surface in Figure 1.

Theorem 1.1. *Let \mathcal{P} and \mathcal{W} be respectively a travelling paraboloid and catenoid, which intersect transversally. Then for all $\varepsilon > 0$ small, there is a complete embedded 3-end surface M_ε satisfying equation (1.8), which lies within an ε -neighborhood of $\mathcal{P} \cup \mathcal{W}$. Besides we have that*

$$\text{genus}(M_\varepsilon) \sim \frac{1}{\varepsilon}.$$

The construction provides much finer properties of the surface M_ε . Let us point out that the CHM with large genus was found in [13] to approach in the multiple-tunnel zone, a Scherk singly periodic minimal surface. See Figure 1.




FIGURE 2. Travelling paraboloid and catenoid

Kapouleas [18, 19, 20], Traizet [31, 32] and Hauswirth and Pacard [12] established a method for the reverse operation, namely, starting with a union of intersecting catenoids and planes, they desingularize them using Scherk surfaces to produce smooth minimal surfaces (complete and embedded). A key element in those constructions is a fine knowledge of the Jacobi operator of the Scherk surface and along the asymptotically flat ends. This approach was used by the third author to construct translating surfaces in \mathbb{R}^3 built from a 2d picture of intersecting parallel grim reapers and vertical lines, trivially extended in an additional direction, desingularized in that direction by infinite Scherk surfaces, see [25, 26, 27]. We shall use a similar scheme in our construction. The context here is considerably more delicate, since no periodicity is involved (the ultimate reason why the topology resulting is finite), and the fine interplay between the slowly vanishing curvatures and the Jacobi operators of the different pieces requires new ideas. Our method extends to the construction of more general surfaces built upon desingularization of intersection of multiple travelling catenoids and travelling paraboloids, but for simplicity in the exposition we shall restrict ourselves to the basic context of Theorem 1.1. Before proceeding into the detailed proof, we sketch below the core ingredients of it.

1.1. Sketch of the proof of Theorem 1.1. After a change of scale of $1/\varepsilon$, the problem is equivalent to finding a complete embedded surface $M \subset \mathbb{R}^3$ that satisfies

$$(1.9) \quad H_M = \varepsilon \nu \cdot e_z.$$

The first step is to construct a surface \mathcal{M} that is close to being a solution to this equation. This is accomplished by desingularizing the union of \mathcal{P}/ε and $\mathcal{W}_R/\varepsilon$ using singly periodic Scherk surfaces. At a large distance from C_ρ/ε , the approximation \mathcal{M} is $\mathcal{P}/\varepsilon \cup \mathcal{W}_R/\varepsilon$ and in some neighborhood of C_ρ/ε , it is a slightly bent singly periodic Scherk surface. We call the core of \mathcal{M} the region where the desingularization is made. The actual approximation \mathcal{M} will depend on four real parameters: $\beta_1, \beta_4, \tau_1, \tau_4$, which are going to be small, of order ε .

Let ν denote a choice of unit normal of \mathcal{M} . We search for a solution of (1.9) in the form of the normal graph over \mathcal{M} of a function $\phi : \mathcal{M} \rightarrow \mathbb{R}$, that is, of the form

$$\mathcal{M}_\phi = \{x + \phi\nu(x) : x \in \mathcal{M}\}.$$

Let H_ϕ and ν_ϕ denote the mean curvature and normal vector of \mathcal{M}_ϕ , respectively, while H and ν denote those of \mathcal{M} . Then

$$(1.10) \quad H_\phi = H + \Delta\phi + |A|^2\phi + Q_1,$$

$$\nu_\phi = \nu - \nabla\phi + Q_2,$$

where Δ is the Laplace-Beltrami operator on \mathcal{M} , ∇ is the tangential component of the gradient, and Q_1, Q_2 are quadratic functions in $\phi, \nabla\phi, D^2\phi$. This allows us to write equation (1.9) as

$$(1.11) \quad \Delta\phi + |A|^2\phi + \varepsilon\nabla\phi \cdot e_z + H - \varepsilon\nu \cdot e_z + Q(x, \phi, \nabla\phi, D^2\phi) = 0 \quad \text{in } \mathcal{M}.$$

To solve (1.11), we linearize around $\phi = 0$, and the following linear operator becomes relevant:

$$\mathcal{L}_\varepsilon(\phi) = \Delta\phi + |A|^2\phi + \varepsilon\nabla\phi \cdot e_z.$$

We work with the following norms for functions ϕ, h defined on \mathcal{M} , where $0 < \gamma < 1$, $0 < \alpha < 1$ are fixed:

$$(1.12) \quad \|\phi\|_* = \sup_{s(x) \leq \delta_s/\varepsilon} e^{\gamma s(x)} \|\phi\|_{C^{2,\alpha}(\overline{B}_1(x))} + \varepsilon^2 \sup_{s(x) > \delta_s/\varepsilon} e^{\gamma\delta_s/\varepsilon + \varepsilon\gamma s(x)} \|\phi\|_{C^{2,\alpha}(\overline{B}_1(x))}$$

and

$$(1.13) \quad \|h\|_{**} = \sup_{s(x) \leq \delta_s/\varepsilon} e^{\gamma s(x)} \|h\|_{C^\alpha(\overline{B}_1(x))} + \sup_{s(x) > \delta_s/\varepsilon} e^{\gamma\delta_s/\varepsilon + \varepsilon\gamma s(x)} \|h\|_{C^\alpha(\overline{B}_1(x))}.$$

Here $\delta_s > 0$ is a small fixed parameter. The function $s : \mathcal{M} \rightarrow \mathbb{R}$ measures geodesic distance to the core of \mathcal{M} and will be defined precisely later on, and $B_1(x)$ is the geodesic ball centered at x with radius 1.

The term in (1.11) that does not depend on ϕ is

$$E = H - \varepsilon\nu \cdot e_z.$$

We have the following approximation for it.

Proposition 1.2. *E can be decomposed as*

$$E = E_0 + E_d$$

with

$$\|E_0\|_{**} \leq C\varepsilon$$

and

$$E_d = \tau_1 w_1 + \tau_4 w_2 + \beta_1 w'_1 + \beta_4 w'_2 + O\left(\sum_{i=1,4} \beta_i^2 + \tau_i^2\right).$$

The functions w_i, w'_i are defined later in (2.8), (2.9), but they and the ones appearing in $O(\sum_{i=1,4} \beta_i^2 + \tau_i^2)$ are smooth with compact support, in particular have finite $\|\cdot\|_{**}$ norm.

The following claim illustrates the invertibility of the linear operator \mathcal{L}_{ve} , although it will not be used directly. Let us fix \mathcal{M} by fixing the parameters $\beta_1, \beta_4, \tau_1, \tau_4$ sufficiently small and consider the problem

$$(1.14) \quad \Delta\phi + |A|^2\phi + \varepsilon\partial_z\phi = h + \sum_{i=1,4} \tilde{\beta}_i w'_i + \tilde{\tau}_i w_i \quad \text{in } \mathcal{M}.$$

Then, for $\varepsilon > 0$ small, there is a linear operator $h \mapsto \phi, \tilde{\beta}_i, \tilde{\tau}_i$ that produces for $\|h\|_{**} < \infty$ a solution of (1.14) with

$$\|\phi\|_* + |\tilde{\beta}_1| + |\tilde{\beta}_4| + |\tilde{\tau}_1| + |\tilde{\tau}_4| \leq C\|h\|_{**},$$

where C is independent of ε .

Finally, the next result shows that the quadratic term Q in (1.11) is well adapted to the norms (1.12) and (1.13).

Proposition 1.3. *Assume $\phi_i \in C^{2,\alpha}(\mathcal{M})$ ($i = 1, 2$) and $\|\phi_i\|_* \leq 1$. Then, for $\varepsilon > 0$ small,*

$$\|Q(\cdot, \phi_1, \nabla\phi_1, D^2\phi_1) - Q(\cdot, \phi_2, \nabla\phi_2, D^2\phi_2)\|_{**} \leq C(\|\phi_1\|_* + \|\phi_2\|_*)\|\phi_1 - \phi_2\|_*,$$

with C independent of ϕ_i and ε .

These results can be used to prove Theorem 1.1 by the contraction mapping principle, which is done in Section 6. The preparatory steps are the construction of an initial approximate solution in Section 2 and some geometric computations in Section 3, which lead to the estimate of E in Proposition 1.2 and the estimate of Q in Proposition 1.3. In Section 4, we analyze the Jacobi equation for the Scherk surface and in Section 5.1, we study the Jacobi operator on the ends, which are the regions far from the desingularization.

2. CONSTRUCTION OF AN INITIAL APPROXIMATION

The purpose of this section is to construct a surface \mathcal{M} that will serve as an initial approximation to (1.8).

Let F_0 the unique radially symmetric solution of

$$(2.1) \quad \frac{F''}{1 + (F')^2} + \frac{F'}{r} = 1, \quad F(0) = 0$$

and let \mathcal{P} be the corresponding surface $z = F_0(r)$. Let \mathcal{W} be a catenoidal self-translating solution of MCF, which can be written as $\mathcal{W} = \mathcal{W}^+ \cup \mathcal{W}^-$ where \mathcal{W}_R^\pm is given by $z = F^\pm(r)$ and F_R^\pm satisfies (2.1) for $r > R$, with $F^+(R) = F^-(R)$, $\lim_{r \rightarrow R^+} (F^+)'(r) = \infty$, $\lim_{r \rightarrow R^+} (F^-)'(r) = -\infty$.

We assume that \mathcal{P} and \mathcal{W} intersect transversally at a unique circle C_ρ of radius $\rho > 0$. To quantify the transversality, we fix a small constant $\delta_\alpha > 0$ so that all the intersection angles are greater than $4\delta_\alpha$. In this section, we are going to replace $\mathcal{P} \cup \mathcal{W}$ in a neighborhood of C_ρ with an appropriately bent Scherk surface. The number of periods used, and thus the number of handles, is of order ε^{-1} . Two of the three ends of the resulting approximate solution will differ slightly from the original ends.

2.1. Self-translating rotationally symmetric surfaces. We briefly recall some properties of self-translating rotationally symmetric surfaces. Let $\varepsilon > 0$ be a small constant, let $\gamma(s) = (\gamma_1(s), \gamma_3(s))$, $s \in [0, \infty)$ be a smooth planar curve parametrized by arc length and let S and S_ε be the surfaces of revolution parametrized by

$$(2.2) \quad \begin{cases} (s, \theta) \mapsto X(s, \theta) := (\gamma_1(ps) \cos(\theta), \gamma_1(ps) \sin(\theta), \gamma_3(ps)) \\ (s, \theta) \mapsto X_\varepsilon(s, \theta) := \varepsilon^{-1}(\gamma_1(\varepsilon ps) \cos(\varepsilon\theta), \gamma_1(\varepsilon ps) \sin(\varepsilon\theta), \gamma_3(\varepsilon ps)), \end{cases}$$

where $p = \gamma_1(0)$, and $s \in [0, \infty)$, $\theta \in [0, 2\pi]$. (The reason for introducing p in (2.2) is to make the parametrization conformal at $s = 0$.)

The surface S (S_ε respectively) is a self-translating surface under mean curvature flow with velocity e_z (εe_z respectively) if and only if γ , parametrized by arc length, satisfies the differential equation

$$(2.3) \quad -\gamma_1''\gamma_3' + \gamma_1'\gamma_3'' + \frac{\gamma_3'}{\gamma_1} - \gamma_1' = 0.$$

Another way to represent an axially symmetric self-translating solution is through the graph of a radial function, $z = F(r)$, where F satisfies (2.1) on some interval (R, ∞) . Then $\varphi = F'$ satisfies

$$(2.4) \quad \varphi' = (1 + \varphi^2) \left(1 - \frac{\varphi}{r}\right).$$

Given $R > 0$ and an initial condition $\varphi(R) = \varphi_0 \in \mathbb{R}$, equation (2.4) has unique solution, which is defined for all $r \geq R$, see [8]. All solutions have the common asymptotic behavior

$$(2.5) \quad \varphi(r) = r - \frac{1}{r} - \frac{2}{r^3} + O\left(\frac{1}{r^5}\right), \quad \varphi'(r) = 1 + \frac{1}{r^2} + O\left(\frac{1}{r^4}\right),$$

as $r \rightarrow \infty$, see [2, 8] (actually an expansion to arbitrary order is possible).

Using $\gamma_1(s) = r(s)$, $\gamma_3(s) = F(r)$, with $s(r) = \int_0^r \sqrt{1 + \varphi(t)^2} dt$ and the asymptotic behavior (2.5), we can deduce the following estimates.

Lemma 2.1. *For a smooth planar curve $\gamma(s) = (\gamma_1(s), \gamma_3(s))$, $s \in [0, \infty)$ parametrized by arc length with γ_1 and γ_3 satisfying (2.3), we have*

$$\begin{aligned} \gamma_1(s) &= \sqrt{2s} + \frac{1}{2} + o(1) & \gamma_3(s) &= s + O(\sqrt{s}) \\ \gamma_1'(s) &= \frac{1}{\sqrt{2s}} + o(s^{-1/2}) & \gamma_3'(s) &= 1 + O(s^{-1/2}) \\ \gamma_1''(s) &= O(s^{-3/2}) & \gamma_3''(s) &= O(s^{-2}) \end{aligned}$$

as s tends to infinity.

2.2. The Scherk surfaces. Let x, y, z be Euclidean coordinates in \mathbb{R}^3 and consider the one parameter family of minimal surfaces $\{\Sigma(\alpha)\}_{\alpha \in (0, \pi/2)}$ given by the equation

$$(2.6) \quad \cos^2(\alpha) \cosh\left(\frac{x}{\cos \alpha}\right) - \sin^2(\alpha) \cosh\left(\frac{y}{\sin \alpha}\right) - \cos(z) = 0.$$

Outside of a large cylinder around the z -axis, $\Sigma(\alpha)$ has four connected components. We call these components the wings of $\Sigma(\alpha)$ and number them according to the quadrant where they lie. Each wing of $\Sigma(\alpha)$ is asymptotic to a half-plane forming an angle α with the xz -plane (note that the asymptotic half-planes do not contain the z -axis unless $\alpha = \pi/4$). Here, we will restrict the parameter α to $[\delta_\alpha, \pi/2 - \delta_\alpha]$

so that the geometry on all the $\Sigma(\alpha)$'s can be uniformly bounded as stated in the following lemma.

Let H^+ be the half-plane $\{(s, z) : s > 0\}$. Note that the parameter s here is on a different scale than the one used in the previous section. We construct approximate solutions satisfying (1.9) here, while the rotationally symmetric surfaces in Section 2.1 satisfy (1.8).

Lemma 2.2. *$\Sigma(\alpha)$ is a singly periodic embedded complete minimal surface which depends smoothly on α . There is a constant $a = a(\delta_\alpha) > 0$ and smooth functions $f_\alpha : H^+ \rightarrow \mathbb{R}$ so that the wings of Σ_α can be expressed as the graph of f_α over half-planes. More precisely, the half-plane asymptotic to the first wing can be parametrized by $A_\alpha^1 : H^+ \rightarrow \mathbb{R}^3$, with*

$$A_\alpha^1(s, z) := (a + s)((\cos \alpha)e_x + (\sin \alpha)e_y) + ze_z + b_\alpha \nu_\alpha,$$

where $b_\alpha = \sin(2\alpha) \log |\cot \alpha|$ and $\nu_\alpha = -(\sin \alpha)e_x + (\cos \alpha)e_y$. The wing itself is parametrized by $F_\alpha^1 : H^+ \rightarrow \mathbb{R}^3$, which is defined by

$$F_\alpha^1(s, z) := A_\alpha^1(s, z) + f_\alpha(s, z)\nu_\alpha.$$

The functions f_α and F_α^1 depend smoothly on α . Moreover, we have

$$\left\| e^s \frac{d^i f_\alpha}{d\alpha^i} \right\|_{C^k(\overline{H^+})} \leq C_{k,i} e^{-a}$$

for any $k, i \in \mathbb{N}$.

The function $f_\alpha(s, z)$ satisfies the minimal surface equation

$$(2.7) \quad \partial_s \left(\frac{\partial_s f}{\sqrt{1 + (\partial_s f)^2 + (\partial_z f)^2}} \right) + \partial_z \left(\frac{\partial_z f}{\sqrt{1 + (\partial_s f)^2 + (\partial_z f)^2}} \right) = 0,$$

for $s > 0, z \in \mathbb{R}$.


Definition 2.3. *Let us denote by \mathcal{R}_{yz} the reflection across the yz -plane and \mathcal{R}_{xz} the reflection across the xz -plane. The parametrizations of the second, third, and fourth wings are given by*

$$F_\alpha^2 = \mathcal{R}_{yz} \circ F_\alpha^1, \quad F_\alpha^3 = \mathcal{R}_{xz} \circ \mathcal{R}_{yz} \circ F_\alpha^1, \quad F_\alpha^4 = \mathcal{R}_{xz} \circ F_\alpha^1.$$

The i th wing of $\Sigma(\alpha)$ is given by $F_\alpha^i(H^+)$ and is denoted by $W^i(\alpha)$. The parametrizations of the corresponding asymptotic half-planes are obtained by replacing F_α^1 by A_α^1 in the above formulas. We use A_α^i to denote the parametrization of the i th asymptotic half-plane as well as its image, $A_\alpha^i(H^+)$. The **inner core** of $\Sigma(\alpha)$ is the surface without its four wings.

Each half-plane $A_\alpha^i(H^+)$ starts close to the boundary of the corresponding wing W^i and intersects neither the xz -plane nor the yz -plane. Each wing and each asymptotic half-plane inherit the coordinates (s, z) from their descriptions in Lemma 2.2 and Definition 2.3.

2.3. Dislocation of the Scherk surfaces. We now perform dislocations on the first and fourth wings of $\Sigma(\alpha)$. These perturbations will help us to deal with the kernel of the linear operator \mathcal{L}_ε associated to normal perturbations of solutions to (1.9). Indeed, because translated solutions of (1.9) remain solutions, the functions $e_x \cdot \nu$, $e_y \cdot \nu$, and $e_z \cdot \nu$ are in the kernel of \mathcal{L}_ε . Here we have taken the normal component of the translations because we are considering normal perturbations.

FIGURE 3. Sections of the Scherk surface $\Sigma(\alpha)$.

The last function, $e_z \cdot \nu$ does not satisfy our imposed symmetries. Therefore, we can discard it from the kernel. The other two remain. In Section 4, we will show that the Dirichlet problem for the linear operator can be solved on a truncated piece of $\Sigma(\alpha)$, up to constants at the boundary. By adding a linear combination of the functions in the kernel, we can obtain a solution that vanishes on the boundary of two adjacent wings, say the second and third wings. To obtain a solution that vanishes on all the connected pieces of the boundary, we will artificially translate the first and fourth wing by constants τ_1 and τ_4 .

The linear operator \mathcal{L}_ε is close to linear operator $L := \Delta + |A|^2$ associated to the equation $H = 0$, so we have small eigenvalues due to changes of Scherk angle and rotation. Indeed, because there is a one parameter family of Scherk surfaces, we expect a function in the kernel of the Jacobi operator L , namely, the normal component of the motion associated to changing the angle α . One more dimension is generated by rotation of the Scherk surfaces around the z -axis. To summarize, besides the translations, we have two more dimensions in the kernel of L generated by linear functions along the wings. This is reasonable since the L is close to the Laplace operator along the wings. By adding a linear combination of these two linear eigenfunctions, we can force exponential decay along the second and third wings again. As before, we will generate linear functions on the first and fourth wings through rotations by angles β_1 and β_4 respectively.

Definition 2.4. For $\beta \in \mathbb{R}$, we define the map $Z_\beta : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ to be the rotation of angle β (counterclockwise in the xy -plane) around the z -axis:

$$Z_\beta(x, y, z) = (\cos(\beta)x - \sin(\beta)y, \sin(\beta)x + \cos(\beta)y, z).$$




FIGURE 4. Dislocations in wing 1

In what follows, we will confine β to $(-\delta_p, \delta_p)$, where $\delta_p > 0$ is a small fixed number.

We consider two constants $R_{rot} > 10$, $R_{tr} > R_{rot} + 10$, and a family of smooth transition functions $\eta_b : \mathbb{R} \rightarrow \mathbb{R}$ such that $0 \leq \eta_b(s) \leq 1$, $\eta_b(s) = 0$ for $s < b$, and $\eta_b(s) = 1$ for $s > b + 1$. The numbers R_{rot} , R_{tr} will be fixed later to be large.

Given $\alpha \in [\delta_\alpha, \pi/2 - \delta_\alpha]$, $\beta_1, \beta_4, \tau_1, \tau_4 \in (-\delta_p, \delta_p)$, we modify the first and fourth wings in the following way: the i th wing is shifted by τ_i at around $s = R_{tr}$, then it is rotated by an angle β_i at distance $s = R_{rot}$. The parametrization of the new i th wing, for $i = 1, 4$, is given by $F^i[\alpha, \beta_i, \tau_i] : H^+ \rightarrow \mathbb{R}^3$, where

$$\begin{aligned} F^1[\alpha, \beta_1, \tau_1](s, z) &= (1 - \eta_{R_{rot}}(s))F_\alpha^1(s, z) + \eta_{R_{rot}}(s)Z_{\beta_1}(F_\alpha^1(s, z) + \tau_1\eta_{R_{tr}}(s)\nu_\alpha) \\ F^4[\alpha, \beta_4, \tau_4](s, z) &= \mathcal{R}_{xz} \circ F^1[\alpha, \beta_4, \tau_4](s, z), \end{aligned}$$

and \mathcal{R}_{xz} is the reflection across the xz -plane. Note that the i th wing is moved away from the x -axis for positive constants β_i and τ_i . We denote the new wings by $W^i[\alpha, \beta_i, \tau_i] := F^i[\alpha, \beta_i, \tau_i](H^+)$, $i = 1, 4$ (see Figure 4).

The wings have natural coordinates (s, z) given by the parametrizations F^1 and F^4 . The surface $\Sigma'[\alpha, \beta_1, \beta_4, \tau_1, \tau_4]$ (or Σ' for short) is defined to be the union of the inner core of $\Sigma(\alpha)$ with the four wings $W^2(\alpha)$, $W^3(\alpha)$, $W^1[\alpha, \beta_1, \tau_1]$, and $W^4[\alpha, \beta_4, \tau_4]$. We will call the region of Σ' for which $s \in [0, R_{tr} + 10]$ the **outer core**.

Remark 2.5. The maps $F_\alpha^i \circ (F^i[\alpha, \beta_i, \tau_i])^{-1}$ and $F^i[\alpha, \beta_i, \tau_i] \circ (F_\alpha^i)^{-1}$, $i = 1, 4$ can be used to pullback tensors defined on $W^i(\alpha)$ to $W^i[\alpha, \beta_i, \tau_i]$ and vice versa: in the case of a function f defined on $W^i[\alpha, \beta_i, \tau_i]$, the composition $f \circ F^i[\alpha, \beta_i, \tau_i] \circ (F_\alpha^i)^{-1}$ is the corresponding pullback function on $W^i(\alpha)$. Taking each wing at a time, these maps transport functions and tensors between Σ' and $\Sigma(\alpha)$. This is very useful as it lets us work on a fixed surface, usually $\Sigma(\alpha)$. We will use the same notation for functions and tensors on Σ' or their pullback to $\Sigma(\alpha)$. For example, $H_{\Sigma'}$ could denote the mean curvature of Σ' as a function on Σ' or its pullback to $\Sigma(\alpha)$. The same notation convention applies to the unit normal vector ν , the metric g , and the second fundamental form A .

Let us define the following functions on $\Sigma(\alpha)$, which capture the contribution of the dislocations to the mean curvature:

$$(2.8) \quad w_1 := \frac{d}{d\tau_1} \Big|_{\beta_i=\tau_i=0} H_{\Sigma'}, \quad w_2 := \frac{d}{d\tau_4} \Big|_{\beta_i=\tau_i=0} H_{\Sigma'},$$

$$(2.9) \quad w'_1 := \frac{d}{d\beta_1} \Big|_{\beta_i=\tau_i=0} H_{\Sigma'}, \quad w'_2 := \frac{d}{d\beta_4} \Big|_{\beta_i=\tau_i=0} H_{\Sigma'}.$$

These functions are compactly supported because rotations and translations do not change the mean curvature. They will later help us solve the Dirichlet problem associated to the Jacobi operator $\Delta + |A|^2$ on the Scherk surfaces in Section 4.

Because the parameters β_i are associated to rotations, the functions w'_1 and w'_2 can be written explicitly as the Jacobi operator on the normal component of rotation at a point $(x, y, z) \in \Sigma(\alpha)$:

$$(2.10) \quad \begin{cases} w'_1(x, y, z) = (\Delta_{\Sigma(\alpha)} + |A_{\Sigma(\alpha)}|^2)(\eta_{rot,1}\nu_{\Sigma(\alpha)} \cdot (-y, x, 0)), \\ w'_2(x, y, z) = (\Delta_{\Sigma(\alpha)} + |A_{\Sigma(\alpha)}|^2)(\eta_{rot,4}\nu_{\Sigma(\alpha)} \cdot (-y, x, 0)), \end{cases}$$

where $\eta_{rot,1}(s)$ is defined as $\eta_{R_{rot}}(s)$ on wing 1 and zero elsewhere and similarly for $\eta_{rot,4}$. We also have,

$$(2.11) \quad \begin{cases} w_1 = (\Delta_{\Sigma(\alpha)} + |A_{\Sigma(\alpha)}|^2)(\eta_{tr,1}), \\ w_2 = (\Delta_{\Sigma(\alpha)} + |A_{\Sigma(\alpha)}|^2)(\eta_{tr,4}), \end{cases}$$

on $\Sigma(\alpha)$, where $\eta_{tr,1}(s) = \eta_{R_{tr}}$ on wing 1 and zero elsewhere, and similarly for $\eta_{tr,4}$.

2.4. Wrapping the dislocated Scherk surfaces around a circle. We first rotate our new surface Σ' so that its second and third wings match the directions of two chosen pieces of catenoid or paraboloid coming out of the intersection circle. The wrapping is performed by simply using a smooth map from a tubular neighborhood of the z -axis to a neighborhood of a large circle. The scaling factor is ε^{-1} so our target circle will have a radius of order ε^{-1} .

Definition 2.6. For $\varepsilon > 0$ and $\varrho > 0$, we define

$$B_{\varepsilon,\varrho}(x, y, z) = (\varepsilon^{-1}\varrho + x)(\cos(\varepsilon\varrho^{-1}z), \sin(\varepsilon\varrho^{-1}z), 0) + (0, 0, y).$$

This map takes a segment of length $2\pi\varepsilon^{-1}\varrho$ on the z -axis to the circle of radius $\varepsilon^{-1}\varrho$.

We can not wrap the whole surface Σ' , so we cut its four wings at $s = R_{tr} + 10$ and denote the new surface by $\bar{\Sigma}'$, with a “bar” on top to indicate that it has a boundary. Our desingularizing surface is a dislocated rotated wrapped Scherk surface

$$(2.12) \quad \bar{\Sigma} := B_{\varepsilon,\rho_\varepsilon} \circ Z_\beta(\bar{\Sigma}'),$$

where the angle β has yet to be fixed and ρ_ε is the closest number in $\varepsilon\mathbb{Z}$ to ρ (the radius associated to the original intersection C_ρ)

We wish to prolong the wings of the desingularizing surface $\bar{\Sigma}$ with pieces of self-translating catenoids or paraboloids. At this point, it will be useful to record the boundary, not of the surface $\bar{\Sigma}$ itself, but of the asymptotic plane underneath at $s = R_{tr} + 10$. We will extend the asymptotic pieces first, then construct the approximate surface by adding the graph of f_α .




FIGURE 5. Paraboloid and catenoid with transversal intersection at C_ρ .

2.5. Fitting the Scherk surface. It is now time to examine the initial configuration in detail. We will work with cross-sections in the xz -plane. Let C_ρ denote the intersection of the paraboloid and catenoid and let $\alpha_0 \in [\delta_\alpha, \pi/2 - \delta_\alpha]$ be half of the angle of intersection between the top \mathcal{W}_R and the inner part of \mathcal{P} (see Figure 5).

On the bounded part of the paraboloid, we take two points P and P' at distances $(a + b_{\alpha_0} \cot \alpha_0 + R_{tr} + 20)\varepsilon$ and $(a + b_{\alpha_0} \cot \alpha_0 + R_{tr} + 10)\varepsilon$ from C_ρ respectively and consider the half-line starting at P' tangent to the paraboloid and pointing to C_ρ . Recall that a was chosen in Lemma 2.2 and that the term $a + b_{\alpha_0} \cot \alpha_0$ is present because the distance from the intersection of the two (extended) asymptotic planes $A_\alpha^2(\mathbb{R}^2)$ and $A_\alpha^3(\mathbb{R}^2)$ to the line $A_\alpha^2(\{s = 0\})$ on is $a + b_\alpha \cot \alpha$ by Lemma 2.2. The new object $\tilde{\mathcal{P}}$ is formed by the paraboloid up to P , a smooth interpolating curve from P to P' , and the tangent half-line after P' (see Figure 6). We will also denote its corresponding surface of revolution by $\tilde{\mathcal{W}}$. We do a similar construction with the catenoid \mathcal{W}_R and denote the new object $\tilde{\mathcal{W}}$ (see Figure 6).

For $\varepsilon > 0$ small, the curves $\tilde{\mathcal{P}}$ and $\tilde{\mathcal{W}}$ intersect at a point $C_{\tilde{\rho}}$. We choose the angle α of the Scherk surface and the angle β of the rotation such that lines A_α^2 , A_α^3 are parallel to the segments $C_{\tilde{\rho}}Q'$ and $C_{\tilde{\rho}}P'$ respectively. Note that $\alpha = \alpha_0(1 + O(\varepsilon))$ and we did not dislocate the second and third wings of the Scherk surface, so α and β do not depend on β_1, β_2, τ_1 , or τ_2 .

Because we have approximated our original curves \mathcal{W}_R and \mathcal{P} up to first order, the new intersection point $C_{\tilde{\rho}}$ is at distance $O(\varepsilon^2)$ from C_ρ . By the same reasoning, the distance from $C_{\tilde{\rho}}$ to either P' or Q' is $(a + b_{\alpha_0} \cot \alpha_0 + R_{tr} + 10)\varepsilon + O(\varepsilon^2)$. Combining with the estimate on α , we have

$$(2.13) \quad |C_{\tilde{\rho}}Q'|, |C_{\tilde{\rho}}P'| = (a + b_\alpha \cot \alpha + R_{tr} + 10)\varepsilon + O(\varepsilon^2).$$

We have to adjust the scale so that the dislocated bent Scherk surface $\bar{\Sigma}$ given in (2.12) fits around the circle of radius $\tilde{\rho}$ and so that its second and third asymptotic half-planes contain part of the line segments of $\tilde{\mathcal{W}}$ and $\tilde{\mathcal{P}}$ respectively. Considering the image of the z -axis under $\varepsilon B_{\varepsilon, \rho_\varepsilon} \circ Z_\beta$ would be a mistake because in general,




FIGURE 6. Step in the construction.

the second and third asymptotic planes do not meet there. Instead, we look at the image under $\varepsilon B_{\varepsilon, \rho_\varepsilon} \circ Z_\beta$ of the line $(b_\alpha / \sin \alpha, 0, z)$ (see Figure 3) and obtain a circle of radius $\rho' = \rho_\varepsilon + \varepsilon \frac{b_\alpha \cos \beta}{\sin \alpha} = \rho_\varepsilon(1 + O(\varepsilon)) = \rho(1 + O(\varepsilon))$. This is the desired radius.

We had to wrap our Scherk surface around a circle of radius ρ_ε to get an embedded surface, so now we adjust the scale by defining $\lambda_\varepsilon = \frac{\rho}{\rho'} = 1 + O(\varepsilon^2)$. This function is not continuous in ε and the jumps occur when the number of periods of the desingularizing surface increases. We take $\lambda_\varepsilon \varepsilon \bar{\Sigma}$ and shift it vertically so that the asymptotic cone associated with the second wing matches the cone generated by the straight part of \tilde{W} on an open set. The cone associated to the third wing aligns automatically with the cone of \tilde{P} by our choice of α and β . We record the amount of vertical displacement with the constant d_ε and denote the shifted surface by $\lambda_\varepsilon \varepsilon \bar{\Sigma}^\dagger$.

We now work on the large scale of equation (1.9). The scaled surface $\lambda_\varepsilon \bar{\Sigma}^\dagger$ has a boundary at $s = R_{tr} + 10$. We wish extend the underlying asymptotic cones with pieces of catenoidal ends that will match the cones in a C^1 manner. We take the curve γ_2 to be just a parametrization of \mathcal{W}_R and γ_3 to be the curve generating the inner part of \mathcal{P} . Thanks to the estimate (2.13), the curves γ_2 and γ_3 , the C^1 matching will occur for some $s \in (R_{tr} + 9, R_{tr} + 11)$ if ε is small enough. For $i = 1$ and 4 , we consider the circle on the i th asymptotic cone corresponding to $s = R_{tr} + 10$ and the tangent unit vector to the cone perpendicular to this circle, pointing away from the core. This gives us an initial position and velocity for the unique curve $\gamma_i = (\gamma_{i,1}, \gamma_{i,3}) : [0, \infty) \rightarrow \mathbb{R}^2$, $i = 1, 4$, generating a rotationally self-translating surface. The curves γ_1 and γ_4 are perturbations of sections of the original paraboloidal and lower catenoidal ends. We can assume without loss of generality that the γ_i 's are parametrized by arclength.

The surface \mathcal{M} , which is an approximate solution to (1.9) is defined in the following way. In the region $s \geq R_{tr} + 20$ the i -th wing of \mathcal{M} is taken as a graph

over the rotationally symmetric surface generated by γ_i . More precisely, let

$$X_i(t, \theta) := \varepsilon^{-1}(\gamma_{i,1}(\varepsilon pt) \cos(\varepsilon\theta), \gamma_{i,1}(\varepsilon pt) \sin(\varepsilon\theta), \gamma_{i,3}(\varepsilon pt)),$$

for $t \geq 0$, $\theta \in [0, 2\pi/\varepsilon]$, where $p = \gamma_1(0)$. The factor p is to make the parametrization conformal at $t = 0$, which we will take to be $s = R_{tr} + 20$. The unit normal vector is

$$\nu(t, \theta) = (-\gamma'_{i,3}(\varepsilon pt) \cos(\varepsilon\theta), -\gamma'_{i,3}(\varepsilon pt) \sin(\varepsilon\theta), \gamma'_{i,1}(\varepsilon pt)).$$

We parametrize the i -th wing of \mathcal{M} in the region $R_{tr} + 20 \leq s \leq 5\delta_s/\varepsilon$ by

$$(s, \theta) \mapsto X_i(s - (R_{tr} + 20), \theta) + u(s, \theta)\nu(s - (R_{tr} + 20), \theta)$$

where the function u is given by

$$u(s, \theta) = pf_\alpha(s, \theta)\eta(\varepsilon s),$$

where f_α is as in Lemma 2.2 and η is a cut-off function satisfying $\eta(s) = 1$ for $s \leq 4\delta_s$ and $\eta(s) = 0$ for $s \geq 5\delta_s$. For $s \geq 5\delta_s/\varepsilon$ the surface \mathcal{M} is the union of the four pieces of rotationally symmetric self-translating surfaces generated by the graphs of γ_i 's.

In the region $R_{tr} + 9 \leq s \leq R_{tr} + 20$ we smoothly interpolate the previous parametrization for $s \geq R_{tr} + 20$ with the corresponding one for $s \leq R_{tr} + 9$, where the surface can be written as the graph of a function over a cone.

Lemma 2.7. *There exists a constant $\delta_p > 0$ depending only on δ_α so that the surface $\mathcal{M}[\varepsilon, \beta_1, \beta_4, \tau_1, \tau_4]$ is embedded for $\beta_1, \beta_4, \tau_1, \tau_4 \in (-\delta_p, \delta_p)$ and $\varepsilon \in (0, \delta_p)$. Moreover, \mathcal{M} depends smoothly on β_1, β_4, τ_1 , and τ_4 . It also depends smoothly on ε , except on a countable set.*

Proof. The only point that needs an argument is that the unbounded ends do not intersect for all $\beta_1, \beta_4, \tau_1, \tau_4$ small. This is a consequence of the following observation: consider two solutions $\varphi_i = \varphi_i(r)$, $i = 1, 2$ of (2.4) defined for all $r \geq R$ with initial conditions

$$\varphi_i(R) = \varphi_{i,0}, \quad \varphi_{1,0} > \varphi_{2,0}.$$

By uniqueness of solutions to ODE,

$$\varphi_1(r) > \varphi_2(r) \quad \forall r \geq R.$$

Let $F_i(r)$ be such that $F'_i = \varphi_i$ and $F_1(0) = F_2(0)$. It follows that

$$F_1(r) > F_2(r) + m \quad \forall r \geq R,$$

with $m > 0$ and m remains positive if $\varphi_{1,0} - \varphi_{2,0}$ remains positive. \square

2.6. Summary of notation and terminology. We start by recalling the roles of the different parameters:

- ε controls the overall scale and the error in the construction.
- ρ_ε is the closest number in $\varepsilon\mathbb{Z}$ to ρ , which is the radius of the intersection of the paraboloid and catenoid in the original scale.
- α is the angle associated to the original Scherk surface (see Figure 3).
- β_1 and β_4 are the angles of rotation of the first and fourth wings respectively.
- τ_1 and τ_4 are the amount by which the first and fourth wings are translated (along the normal of the asymptotic plane to the respective wing).

Then we have the different scaled and bent Scherk surfaces:

- $\Sigma(\alpha)$ is the original minimal Scherk surface given by (2.6). Its wings $W^i(\alpha)$ are asymptotic to the half-planes $A_\alpha^i(H^+)$.
- $\Sigma' = \Sigma'[\alpha, \beta_1, \beta_4, \tau_1, \tau_4]$ is $\Sigma(\alpha)$ with dislocations (see Figure 4).
- $\bar{\Sigma} := B_{\varepsilon, \rho_\varepsilon} \circ Z_\beta(\Sigma')$ is Σ' wrapped around a circle of radius $\rho_\varepsilon/\varepsilon$.
- $\lambda_\varepsilon \varepsilon \bar{\Sigma}^\dagger$ is the previous Scherk surface scaled by a factor $\lambda_\varepsilon = \varepsilon(1 + O(\varepsilon^2))$ and shifted by d_ε in the \vec{e}_z direction so that it fits the configuration in Figure 6.

Finally, let us recall the names of the different parts of the initial approximation $\mathcal{M} = \mathcal{M}[\varepsilon, \beta_1, \beta_4, \tau_1, \tau_4]$:

- The inner core is where the handles are.
- The middle core is where we perform all the dislocations. It corresponds to the region $0 \leq s \leq R_{tr} + 10$.
- The region $s \in (R_{tr} + 9, R_{tr} + 20)$ is a transition region called the outer core. Note that there is a change of scale in this region from $\lambda_\varepsilon \varepsilon$ to ε but this will not create too much error as the switch is contained in a bounded region with s small compared to ε^{-1} .
- The core is the union of the inner, middle, and outer core.
- In the region $\{s \in (R_{tr} + 20, 4\delta_s/\varepsilon)\}$, \mathcal{M} is the graph of f_α over the rotationally symmetric surface generated by γ_i , $i = 1, \dots, 4$.
- The region $\{s \in (4\delta_s/\varepsilon, 5\delta_s/\varepsilon)\}$ is a transition region where we cut-off the graph of f_α . We have to do it far enough so that the function f_α is small and the length of the interval has to be long enough so we don't create too much error.
- The wings are defined to be the region $\{s \in (0, 5\delta_s/\varepsilon)\}$.
- For $\{s \geq 5\delta_s/\varepsilon\}$, \mathcal{M} is just the rotationally symmetric surface generated by the curves γ_i 's.

3. GEOMETRIC COMPUTATIONS

In this section, we perform computations related to the ansatz \mathcal{M} . In particular, we prove Proposition 1.2, which gives the error. We also use these computations to prove Proposition 1.3 for the quadratic terms appearing in the expansion of the mean curvature and normal vector.

3.1. Perturbation by normal graphs. Consider a surface M immersed in \mathbb{R}^3 with local parametrization of class C^2 :

$$X : U \subset \mathbb{R}^2 \rightarrow M, \quad X = X(x_1, x_2).$$

We use the notation

$$e_i = \partial_i X = \partial_{x_i} X$$

for tangent vectors and we take the normal unit vector to be

$$\nu = \frac{e_1 \times e_2}{|e_1 \times e_2|},$$

where \times is the cross product in \mathbb{R}^3 . The metric of M is denoted by

$$g_{ij} = \langle e_i, e_j \rangle,$$

and its inverse by g^{ij} , where $\langle \cdot, \cdot \rangle$ is the standard inner product in \mathbb{R}^3 . We recall that

$$\partial_i \nu = -A_i^j e_j$$

where we use Einstein's convention of summation over repeated indices, and A_i^j is the second fundamental form, which can be computed as

$$A_i^j = A_{ik}g^{kj}, \quad A_{ij} = \langle X_{ij}, \nu \rangle = -\langle e_i, \partial_j \nu \rangle.$$

The mean curvature H of M can be computed in the following way

$$H = \text{trace of } A = A_1^1 + A_2^2 = g^{ij} \langle X_{ij}, \nu \rangle.$$

Consider a function $u \in C^2(M)$. We write

$$u_i = \partial_{x_i} u, \quad u_{ij} = \partial_{x_i x_j} u.$$

Let $\tilde{X} = X + \nu u$ be the graph of u over M , we then have

$$\tilde{e}_i = \partial_{x_i} \tilde{X} = e_i + u_i \nu - u A_i^k e_k$$

and

$$\tilde{g}_{ij} = \langle \tilde{e}_i, \tilde{e}_j \rangle = g_{ij} - u(A_j^l g_{il} + A_i^k g_{kj}) + u^2 A_i^k A_j^l g_{kl} + u_i u_j.$$

We compute the cross product

$$\begin{aligned} \tilde{e}_1 \times \tilde{e}_2 &= e_1 \times e_2 (1 - uH - u^2 G) + u_1 \nu \times e_2 + u_2 e_1 \times \nu \\ &\quad - uu_1 A_2^l \nu \times e_l - uu_2 A_1^k e_k \times \nu, \end{aligned}$$

where $G = A_1^1 A_2^2 - A_1^2 A_2^1$ is the Gauss curvature. We also compute the derivatives of \tilde{X} :

$$\begin{aligned} \tilde{X}_{ij} &= \partial_j (e_i + \nu u_i + \nu u_j) \\ &= e_{ij} + u_{ij} \nu - u_i A_j^l e_l - u_j A_i^k e_k - u(A_i^k)_j e_k - u A_i^k e_{kj} \end{aligned}$$

The mean curvature of M_u is given by

$$\tilde{H} = \tilde{g}^{ij} \langle \tilde{X}_{ij}, \frac{\tilde{e}_1 \times \tilde{e}_2}{|\tilde{e}_1 \times \tilde{e}_2|} \rangle$$

Explicitly, the scalar product is

$$\begin{aligned} &\langle \tilde{e}_1 \times \tilde{e}_2, \tilde{X}_{ij} \rangle \\ &= \langle e_1 \times e_2 (1 - uH - u^2 G) + u_1 \nu \times e_2 + u_2 e_1 \times \nu - uu_1 A_2^l \nu \times e_l - uu_2 A_1^k e_k \times \nu, \\ &\quad e_{ij} + u_{ij} \nu - u_i A_j^l e_l - u_j A_i^k e_k - u(A_i^k)_j e_k - u A_i^k e_{kj} \rangle \end{aligned}$$

with $\det \tilde{g} = |\tilde{e}_1 \times \tilde{e}_2|^2$ and

$$\tilde{g}^{-1} = \frac{1}{\det \tilde{g}} \begin{bmatrix} \tilde{g}_{22} & -\tilde{g}_{12} \\ -\tilde{g}_{12} & \tilde{g}_{11} \end{bmatrix}.$$

3.2. Norms of tensors. We work with the following norms, which are independent of coordinates.

Definition 3.1. *The pointwise norm of a tensor $T_{a_1 \dots a_r}^{b_1 \dots b_s}$ is given by*

$$|T|^2 := T_{a_1 \dots a_r}^{b_1 \dots b_s} T_{c_1 \dots c_r}^{d_1 \dots d_s} g^{a_1 c_1} \dots g^{a_r c_r} g_{b_1 d_1} \dots g_{b_s d_s},$$

with summation over repeated indices.

We use the notation ∇_i to denote the covariant derivative with respect to $\frac{\partial}{\partial x_i}$. In the case where the metric is diagonal, the norms of the gradient and Hessian of a function u are

$$|\nabla u|^2 = |u_i u_j g^{ij}| = \frac{(u_1)^2}{g_{11}} + \frac{(u_2)^2}{g_{22}},$$

$$|\nabla^2 u|^2 = (g^{11} \nabla_{11} u)^2 + (g^{22} \nabla_{22} u)^2 + 2g^{11} g^{22} (\nabla_{12} u)^2.$$

For the second fundamental form, we have

$$|A|^2 = (g^{11} A_{11})^2 + (g^{22} A_{22})^2 + 2(g^{11} g^{22} A_{12} A_{12}),$$

$$|\nabla A|^2 = \sum_{i,j,k=1}^2 (\nabla_i A_j^k)^2 g^{ii} g^{jj} g_{kk}.$$

3.3. Geometry of rotationally symmetric self-translating surfaces. We compute various geometric quantities attached to the parametrization X_ε given in (2.2). We use ∂_i or $(\)_i$ to denote regular differentiation with respect to the variables s ($i = 1$) or θ ($i = 2$). Let $\{e_1, e_2\}$ be the tangent vectors to S_ε given by

$$(3.1) \quad e_1 = \partial_1 X_\varepsilon = p(\gamma_1'(\varepsilon ps) \cos(\varepsilon\theta), \gamma_1'(\varepsilon ps) \sin(\varepsilon\theta), \gamma_3'(\varepsilon ps)),$$

$$(3.2) \quad e_2 = \partial_2 X_\varepsilon = (-\gamma_1(\varepsilon ps) \sin(\varepsilon\theta), \gamma_1(\varepsilon ps) \cos(\varepsilon\theta), 0),$$

We recall that $(\gamma_1(s), \gamma_3(s))$ is parametrized by arc length and that $p := \gamma_1(0) > 0$. The associated metric is then

$$(3.3) \quad g_{11} = p^2, \quad g_{12} = g_{21} = 0, \quad g_{22} = \gamma_1'^2.$$

The only nonzero Christoffel symbols are

$$(3.4) \quad \Gamma_{22,1} = -\varepsilon p \gamma_1 \gamma_1', \quad \Gamma_{12,2} = \Gamma_{21,2} = \varepsilon p \gamma_1 \gamma_1',$$

$$\Gamma_{22}^1 = -\varepsilon \frac{\gamma_1}{p} \gamma_1', \quad \Gamma_{12}^2 = \Gamma_{21}^2 = \varepsilon \frac{p}{\gamma_1} \gamma_1'.$$

Using $A_{ij} = \langle \partial_i e_j, \nu \rangle$, we obtain the coordinates of the second fundamental form

$$(3.5) \quad \begin{cases} A_{11} = -\varepsilon p^2 (-\gamma_1'' \gamma_3' + \gamma_1' \gamma_3''), & A_{12} = 0, & A_{22} = \varepsilon \gamma_1 \gamma_3' \\ A_1^1 = -\varepsilon (-\gamma_1'' \gamma_3' + \gamma_1' \gamma_3''), & A_1^2 = A_2^1 = 0 & A_2^2 = \varepsilon \gamma_1^{-1} \gamma_3' \end{cases}$$

where all the functions are taken at εps .

The following proposition is an immediate corollary of (3.5) and the growth of γ_1, γ_3 given in Lemma 2.1.

Proposition 3.2. *In the coordinates given by X_ε , the second fundamental form A and Christoffel symbols on S_ε satisfy*

$$(3.6) \quad \left| \frac{d^k}{ds^k} A_i^j(\varepsilon ps) \right| \leq C \varepsilon^{k+1},$$

$$(3.7) \quad |\Gamma_{ij}^k| \leq C \varepsilon.$$

Proposition 3.3. *We have*

$$(3.8) \quad |\nabla^k A| \leq C \varepsilon^{k+1}, \quad k = 0, 1, 2.$$

Proof. The fact that $|A|^2 \leq C\varepsilon^2$ is straightforward from (3.5), (3.3) and Proposition 3.2. For the first covariant derivative of A , we recall $\nabla_k A_i^j = \partial_k A_1^j - \Gamma_{ki}^l A_l^j + \Gamma_{kl}^j A_i^l$ with implied summation over $l = 1, 2$. Upon inspection, $\nabla_1 A_1^2$ vanishes. If $(i, j, k) \neq (1, 2, 1)$, the quantity $\sqrt{g^{kk} g^{ii} g_{jj}}$ is bounded, therefore

$$\begin{aligned} \sqrt{g^{kk} g^{ii} g_{jj}} |\nabla_k A_i^k| &\leq C(|\partial_k A_1^j| + |\Gamma_{ki}^l A_l^j| + |\Gamma_{kl}^j A_i^l|) \\ &\leq C\varepsilon^2 \end{aligned}$$

and the estimate for $|\nabla A|$ is proved.

For the second covariant derivative, we argue similarly. Recall that $\nabla_{lk}^2 A_i^j = \partial_l(\nabla_k A_i^j) - \Gamma_{lk}^m \nabla_m A_i^j - \Gamma_{li}^m \nabla_k A_m^j + \Gamma_{lm}^j \nabla_k A_i^m$. Note that $\nabla_{11}^2 A_1^2 = 0$. As before, if $(i, j, k, m) \neq (1, 2, 1, 1)$, the product $\sqrt{g^{kk} g^{ii} g_{jj} g^{ll}}$ is bounded and we prove $|\nabla^2 A| \leq C\varepsilon^3$ by combining (3.6), (3.7), and (3.8) for $k = 0, 1$. \square

3.4. Computation of the error.

Proof of Proposition 1.2. The initial approximation \mathcal{M} consists of three parts. The core of \mathcal{M} is a smooth perturbation of a compact piece of Scherk minimal surface, when we consider one period only. This introduces curvatures of order ε , together with some dislocations, so the statement of Proposition 1.2 follows directly for the error restricted to this part.

The region $s \geq 5\delta_s/\varepsilon$ of \mathcal{M} is a rotationally symmetric self-translating surface, so that E is zero here.

Where $R_{tr} + 20 \leq s \leq 5\delta_s/\varepsilon$, the surface \mathcal{M} is a graph over a self-translating rotationally symmetric surface S_ε . We parametrize S_ε with X_ε defined in (2.2), which we will write for convenience as X , so that

$$X(s, \theta) := \varepsilon^{-1}(\gamma_1(\varepsilon ps) \cos(\varepsilon\theta), \gamma_1(\varepsilon ps) \sin(\varepsilon\theta), \gamma_3(\varepsilon ps)),$$

where $p = \gamma_1(0)$, and $s \in [0, \infty)$, $\theta \in [0, 2\pi]$.

Then \mathcal{M} in this region is parametrized by

$$(3.9) \quad (s, \theta) \mapsto \tilde{X}(s, \theta) = X(s, \theta) + u(s, \theta)\nu.$$

where the function u is given by

$$(3.10) \quad u(s, \theta) = pf_\alpha(s, \theta)\eta(\varepsilon s),$$

with f_α given in Lemma 2.2 and η is a cut-off function satisfying $\eta(s) = 1$ for $s \leq 4\delta_s$ and $\eta(s) = 0$ for $s \geq 5\delta_s$. We observe that

$$(3.11) \quad |e^s \partial^k u(s, \theta)| \leq Ce^{-a}, \quad k = 0, \dots, 5,$$

and that $f_\alpha(s, \theta)$ satisfies the minimal surface equation (2.7).

In the rest of this proof, g, A, H, ν denote the metric, second fundamental form, mean curvature and Gauss map of the rotationally symmetric surface S_ε and $\tilde{g}, \tilde{A}, \tilde{H}, \tilde{\nu}$ the ones of \mathcal{M} given by the parametrization above.

In the rest of this section, we shall work in the region $s \leq 5\delta_s/\varepsilon$. Following the calculations of Section 3.1 with the parametrization (3.9) and using (3.8) we have

$$\begin{aligned} \tilde{e}_1 &= (1 - uA_1^1)e_1 - uA_1^2e_2 + u_1\nu = e_1 + \nu u_1 + O(\varepsilon e^{-s}), \\ \tilde{e}_2 &= -uA_2^1e_1 + (1 - uA_2^2)e_2 + u_2\nu = e_2 + \nu u_2 + O(\varepsilon e^{-s}), \end{aligned}$$

where $O(\varepsilon e^{-s})$ is in the C^2 sense on the region $s \leq 5\delta_s/\varepsilon$ as $\varepsilon \rightarrow 0$, and e_1, e_2 are given in (3.1), (3.2). We compute the metric \tilde{g}_{ij} :

$$(3.12) \quad \tilde{g}_{ij} = g_{ij} - 2uA_{ij} + u^2 A_i^l A_j^m g_{lm} + u_i u_j = g_{ij} + u_i u_j + O(\varepsilon e^{-s}).$$

Using (3.3), we have

$$\tilde{g}^{-1} = \frac{1}{\det(\tilde{g})} \begin{pmatrix} \gamma_1^2 + u_2^2 & -u_1 u_2 \\ -u_1 u_2 & p^2 + u_1^2 \end{pmatrix} + O(\varepsilon e^{-s}),$$

where γ_1 is evaluated at $p\varepsilon s$, and

$$(3.13) \quad \det(\tilde{g}) = p^2 \gamma_1^2 + \gamma_1^2 u_1^2 + p^2 u_2^2 + O(\varepsilon e^{-s}).$$

For the normal direction, we recall that e_1 and e_2 are orthogonal and obtain

$$(3.14) \quad \tilde{e}_1 \times \tilde{e}_2 = \det(g)^{1/2} \nu - u_1 e_1 - u_2 e_2 + O(\varepsilon e^{-s}).$$

Next, we compute

$$(3.15) \quad \tilde{X}_{ij} = e_{ij} + u_{ij} \nu + O(\varepsilon e^{-s})$$

and

$$\tilde{A}_{ij} = \langle \tilde{X}_{ij}, \tilde{\nu} \rangle = \frac{\det(g)^{1/2}}{\det(\tilde{g})^{1/2}} (A_{ij} + u_{ij}) + O(\varepsilon e^{-s}).$$

Since $\frac{\det(g)}{\det(\tilde{g})} = 1 + O(\varepsilon e^{-s})$ and A_{ij} are $O(\varepsilon)$, we get

$$(3.16) \quad \tilde{A}_{ij} = A_{ij} + u_{ij} + O(\varepsilon e^{-s}).$$

With this, the second fundamental form can be expressed as

$$\begin{aligned} \tilde{A}_1^1 &= \frac{1}{\det(\tilde{g})} [(A_{11} + u_{11})(\gamma_1^2 + u_2^2) - (A_{12} + u_{12})u_1 u_2] + O(\varepsilon e^{-s}) \\ \tilde{A}_2^2 &= \frac{1}{\det(\tilde{g})} [-(A_{21} + u_{21})u_1 u_2 + (A_{22} + u_{22})(p^2 + u_1^2)] + O(\varepsilon e^{-s}). \end{aligned}$$

Therefore

$$\begin{aligned} \tilde{H} &= \frac{1}{\det(\tilde{g})} [A_{11} \gamma_1^2 + A_{22} p^2 + u_{11}(\gamma_1^2 + u_2^2) - 2u_{12} u_1 u_2 + u_{22}(p^2 + u_1^2)] \\ &\quad + O(\varepsilon e^{-s}). \end{aligned}$$

Let $\bar{u} = \frac{1}{p} u = f_\alpha(s, \theta) \eta(\varepsilon s)$. We recall that $H = \frac{1}{\det(g)} [A_{11} \gamma_1^2 + A_{22} p^2]$ and expand $\gamma_1(p\varepsilon s) = p + O(\varepsilon s)$ in the region $s \leq 5\delta_s/\varepsilon$ to get

$$\tilde{H} = H + \frac{p^3}{\det(\tilde{g})} [\bar{u}_{11}(1 + \bar{u}_2^2) - 2\bar{u}_{12} \bar{u}_1 \bar{u}_2 + \bar{u}_{22}(1 + \bar{u}_1^2)] + O(\varepsilon s e^{-s})$$

Because $\eta(\varepsilon s) = 1$ for $s \leq 4\delta_s/\varepsilon$ and f_α satisfies (2.7), we actually have

$$\bar{u}_{11}(1 + \bar{u}_2^2) - 2\bar{u}_{12} \bar{u}_1 \bar{u}_2 + \bar{u}_{22}(1 + \bar{u}_1^2) = 0$$

in this region. Thus we obtain

$$\tilde{H} = H + O(\varepsilon s e^{-s}), \quad s \leq 4\delta_s/\varepsilon.$$

Also in this region, from (3.13), (3.14) and $|\tilde{e}_1 \times \tilde{e}_2| = \det(\tilde{g})^{1/2}$ we also have $\varepsilon \tilde{\nu} \cdot e_z = \varepsilon \nu \cdot e_z + O(\varepsilon e^{-s})$ and since S_ε is a self translating surface, we get

$$\tilde{H} - \varepsilon \tilde{\nu} \cdot e_z = O(\varepsilon e^{-\gamma s}).$$

The same estimate holds for derivatives of $\tilde{H} - \varepsilon\tilde{\nu} \cdot e_z$ (all the parametrizations here are smooth), which implies the corresponding estimate in the weighted C^α norm.

When $4\delta_s/\varepsilon \leq s \leq 5\delta_s/\varepsilon$, we have $|\partial^k u(s)| \leq Ce^{-s} \leq \varepsilon e^{-\gamma s}$ for ε small enough, so

$$\tilde{H} = H + O(\varepsilon e^{-\gamma s}), \quad \varepsilon\tilde{\nu} \cdot e_z = \varepsilon\nu \cdot e_z + O(\varepsilon e^{-\gamma s}),$$

and the desired estimate holds. \square

3.5. Estimate of the Jacobi operator. Here we use the following notation: The metric, Christoffel symbols, and second fundamental forms on a rotationally symmetric piece of self-translating surface are denoted by g , Γ_{ij}^k , and A , the corresponding quantities for the ansatz are $g_{\mathcal{M}}$, $\Gamma_{ij,\mathcal{M}}^k$, and $A_{\mathcal{M}}$ while the ones on the corresponding original Scherk surface are g_{Σ} , $\Gamma_{ij,\Sigma}^k$, and A_{Σ} . For short, we write $\Delta_{g_{\mathcal{M}}} = \Delta_{\mathcal{M}}$, $\Delta_{g_{\Sigma}} = \Delta_{\Sigma}$. In the following proposition, the operators on \mathcal{M} (the left-hand side) are pulled back to Σ using the transformations of Section 2.

Proposition 3.4. *For $s \leq 5\delta_s/\varepsilon$, we have*

$$\Delta_{\mathcal{M}} + |A_{\mathcal{M}}|^2 + \varepsilon\langle e_z, \nabla_{g_{\mathcal{M}}} \rangle = \Delta_{\Sigma} + |A_{\Sigma}|^2 + L'$$

where L' is a second order differential operator with coefficients with C^1 norm bounded by $O(\delta_s + \delta_p + \varepsilon)$.

Proof. We again divide into several regions. For $s \leq R_{tr} + 20$, \mathcal{M} is obtained from the Scherk surface by a bending, which introduces terms of order ε and dislocations of order δ_p , so the estimate for L' here follows.

When $R_{tr} + 20 \leq s \leq 5\delta_s/\varepsilon$, \mathcal{M} can be described by the parametrization (3.9). In this region, we express all geometric quantities of \mathcal{M} and Σ as functions of the coordinates s and θ .

We note that $\Delta_{\mathcal{M}} - \Delta_{\Sigma}$ is a second order operator with coefficients whose C^1 norm can be estimated from the C^2 norm of $g_{\mathcal{M}} - g_{\Sigma}$. The ansatz is the graph of u from (3.10), so to be consistent we take the Scherk surface parametrized by $(s, \theta) \mapsto (ps, p\theta, pf_{\alpha}(s, \theta))$. Then (3.12) gives

$$g_{ij,\Sigma} - g_{ij,\mathcal{M}} = p^2\delta_{ij} + p^2f_{\alpha,i}f_{\alpha,j} - (g_{ij} + u_i u_j) + O(\varepsilon e^{-s}).$$

We use (3.3), $\|u\|_{C^3} \leq C$, and the expansion $\gamma_1(p\varepsilon s) = p + O(\varepsilon s)$ to deduce

$$(3.17) \quad \|g_{ij,\Sigma} - \bar{g}_{ij,\mathcal{M}}\|_{C^2} \leq C(\delta_s + \varepsilon),$$

where the norm is computed over $R_{tr} + 20 \leq s \leq 5\delta_s/\varepsilon$. Because the metrics are uniformly equivalent, in order to bound $|A_{\mathcal{M}}|^2 - |A_{\Sigma}|^2$, it suffices to control $|A_{ij,\mathcal{M}}(s, \theta) - A_{ij,\Sigma}(s, \theta)|$. By (3.16),

$$A_{ij,\Sigma} - A_{ij,\mathcal{M}} = pf_{ij} - (u_{ij} + A_{ij}) + O(\varepsilon e^{-s}),$$

where $O(\varepsilon e^{-s})$ is in C^1 norm. The functions u and f are equal if $s \leq 4\delta_s/\varepsilon$ and we have $e^{-s} \leq C\varepsilon$ otherwise. Moreover, $A_{ij} = O(\varepsilon)$ when $s \leq 5\delta_s/\varepsilon$ by (3.5). Therefore,

$$(3.18) \quad \|A_{ij,\Sigma} - A_{ij,\mathcal{M}}\|_{C^1} \leq C\varepsilon,$$

where the norm is over $R_{tr} + 20 \leq s \leq 5\delta_s/\varepsilon$.

Finally, the term $\varepsilon\langle e_z, \nabla_{g_{\mathcal{M}}} \rangle$ has coefficients of order ε in C^1 norm over $R_{tr} + 20 \leq s \leq 5\delta_s/\varepsilon$. \square

3.6. Estimates of the quadratic terms. Here, we prove Proposition 1.3 for functions defined on the surface \mathcal{M} . We recall from (3.18) and Proposition 3.3 that each $|\nabla^i A|$ remains uniformly bounded on \mathcal{M} , for $i = 0, 1, 2$.

Let Q_1 be defined by (1.10) and assume $|uA| < 1$. For this computation it is convenient to work with coordinates that are normal at a certain point $x_0 \in \mathcal{M}$, which means

$$g_{ij}(x_0) = \delta_{ij} \quad \text{and} \quad \partial_k g_{ij}(x_0) = 0.$$

This implies

$$\langle X_{ij}, e_k \rangle = 0,$$

at x_0 . Moreover, by a further rotation $\langle X_{12}, \nu \rangle = 0$ at x_0 so that $A_1^2(x_0) = 0$. With these properties, following the computation in Section 3.1, we obtain at the point x_0 :

$$\tilde{g}^{-1} = \frac{1}{\det(\tilde{g})} \begin{bmatrix} 1 - 2uA_2^2 + u^2(A_2^2)^2 + u_2^2 & -u_1u_2 \\ -u_1u_2 & 1 - 2uA_1^1 + u^2(A_1^1)^2 + u_1^2 \end{bmatrix}$$

with

$$\det(\tilde{g}) = |\tilde{e}_1 \times \tilde{e}_2|^2 = (1 - uH - u^2G)^2 + u_1^2(1 - uA_2^2)^2 + u_2^2(1 - uA_1^1)^2,$$

where G is the Gaussian curvature.

We will use Q to denote different functions of u, u_i, u_{ij}, x with the properties:

$$(3.19) \quad \begin{cases} Q \text{ is } C^\infty \text{ in } u, u_i, u_{ij}, \\ Q(0, 0, 0, x) = 0, \quad D_u Q(0, 0, 0, x) = 0, \\ D_{u_i} Q(0, 0, 0, x) = 0, \quad D_{u_{ij}} Q(0, 0, 0, x) = 0, \\ \text{second derivatives with respect to } u, u_i, u_{ij} \text{ are bounded} \\ \text{by universal functions of } |A| \text{ and } |\nabla A| \text{ for } |uA| < 1/2, \\ Q \text{ is linear in } u_{ij}. \end{cases}$$

Then we can write

$$\tilde{g}^{-1} = \frac{1}{\det(\tilde{g})} \begin{bmatrix} 1 - 2uA_2^2 & 0 \\ 0 & 1 - 2uA_1^1 \end{bmatrix} + Q$$

and

$$\det(\tilde{g}) = 1 - 2uH + Q.$$

Similarly,

$$\langle \tilde{X}_{ij}, \tilde{e}_1 \times \tilde{e}_2 \rangle = (1 - uH)A_i^j + u_{ij} - u(A_i^j)^2 + Q,$$

therefore

$$\tilde{H} = H + u_{11} + u_{22} + ((A_1^1)^2 + (A_2^2)^2)u + Q_1,$$

where Q_1 satisfies the properties (3.19). Let u, v be $C^{2,\alpha}$ functions on \mathcal{M} with $|uA| < 1/2, |vA| < 1/2$. To simplify notation, let $U(x) = (u(x), \nabla u(x), \nabla^2 u(x))$. From the properties of Q_1 and Taylor's formula

$$|Q_1(U(x), x) - Q_1(V(x), x)| \leq C(|U(x)| + |V(x)|)|U(x) - V(x)|.$$

To estimate the Hölder norm of Q_1 , we note that the expression for $\tilde{H} - \varepsilon \tilde{\nu} \cdot e_z$ is linear in the second derivative of u and we have C^1 bounds on all the other terms. We have C^α bounds on $\nabla_g u$ and C^1 bounds on everything else ($u, \nabla_g u, A$, and $\nabla_g A$).

4. THE JACOBI EQUATION ON SCHERK SURFACES

Let $\Sigma = \Sigma(\alpha)$ be the singly periodic Scherk surface defined by (2.6). In this section, we want to solve the problem involving the Jacobi operator on Σ ,

$$\Delta\phi + |A|^2\phi = h \quad \text{in } \Sigma,$$

where Δ is the Laplace-Beltrami operator and A is the second fundamental form of Σ .

We let s and z denote the coordinates on the wings $W^i(\alpha)$, $i = 1, \dots, 4$, described in Lemma 2.2. In the rest of the section, we will work with right-hand sides h defined on Σ and such that $\|e^{\gamma s}h\|_{L^\infty(\Sigma)} < \infty$ for a fixed $\gamma \in (0, 1)$.

We will work with functions that are 2π periodic in z and even with respect to z , that is, ϕ and h satisfy

$$(4.1) \quad \phi(x, y, z) = \phi(x, y, z + 2\pi), \quad \phi(x, y, z) = \phi(x, y, -z) \quad \forall (x, y, z) \in \Sigma,$$

which is equivalent to symmetry with respect to the planes $z = k\pi$, $k \in \mathbb{Z}$.

We choose the unit normal vector to Σ such that

$$(4.2) \quad \nu \cdot e_y > 0 \text{ on wings 1 and 2, and } \nu \cdot e_y < 0 \text{ on wings 3 and 4.}$$

Because translating the surface Σ leaves its mean curvature unchanged, the functions $\nu \cdot e$ are in the kernel of $\Delta + |A|^2$ for any fixed $e \in \mathbb{R}^3$. Hence $\nu \cdot e_x$, $\nu \cdot e_y$, and $\nu \cdot e_z$ are in the kernel of the Jacobi operator. Of these functions, $\nu \cdot e_x$ and $\nu \cdot e_y$ satisfy the symmetries (4.1) and $\nu \cdot e_z$ does not because it is antisymmetric with respect to $z = 0$. We will write

$$(4.3) \quad z_1 = \nu \cdot e_x, \quad z_2 = \nu \cdot e_y.$$

The main results in this section are the following. First, we consider the problem of finding a bounded solutions ϕ of

$$(4.4) \quad \begin{cases} \Delta\phi + |A|^2\phi = \sum_{i=1}^2 c_i \eta_0 z_i + h & \text{in } \Sigma, \\ \phi \text{ satisfies the symmetries (4.1)} \end{cases}$$

for which

$$(4.5) \quad \int_{\Sigma} \phi \eta_0 z_i = 0, \quad i = 1, 2,$$

where $\eta_0 \in C^\infty(\Sigma)$ is a smooth function depending only on $x^2 + y^2$ such that

$$(4.6) \quad 0 \leq \eta_0 \leq 1, \quad \eta_0 = 1 \text{ on } x^2 + y^2 \leq R_0^2, \text{ and } \eta_0 = 0 \text{ on } x^2 + y^2 \geq (R_0 + 1)^2,$$

where $R_0 > 1$ is fixed.

Proposition 4.1. *Let $0 < \gamma < 1$. Let h be defined in Σ , satisfy the symmetries (4.1), and $\|e^{\gamma s}h\|_{L^\infty(\Sigma)} < \infty$. Then there are unique $c_1, c_2 \in \mathbb{R}$ and $\phi \in L^\infty(\Sigma)$ satisfying (4.4) and (4.5). Moreover*

$$(4.7) \quad |c_1| + |c_2| + \|\phi\|_{L^\infty(\Sigma)} \leq C \|e^{\gamma s}h\|_{L^\infty(\Sigma)}$$

and

$$(4.8) \quad |\nabla\phi| \leq C(\|e^{\gamma s}h\|_{L^\infty(\Sigma)} + \|\phi\|_{L^\infty(\Sigma)})e^{-\gamma s}.$$

If ϕ is a bounded solution of (4.4), by (4.8) ϕ has a limit on each wing, that is, $L_i = \lim_{s \rightarrow \infty} \phi(s, z)$ on all wings $i = 1, \dots, 4$. Moreover, these limits define linear functionals of h and we have the estimate $|L_i| \leq C \|e^{\gamma s} h\|_{L^\infty}$. For later consideration, it is desirable to find a solution to (4.4) with limit equal to zero on all wings. To achieve this, the right hand side has to satisfy four restrictions, or equivalently has to be projected onto a space of codimension 4. We do this by considering the main terms introduced by the dislocations. So we consider now the problem

$$(4.9) \quad \begin{cases} \Delta \phi + |A|^2 \phi = h + \sum_{i=1}^2 (\beta_i w'_i + \tau_i w_i) & \text{in } \Sigma, \\ \phi \text{ satisfies (4.1),} \end{cases}$$

where the functions w_i, w'_i are defined in (2.8), (2.9).

Proposition 4.2. *Let $0 < \gamma < 1$ and h be a function satisfying $\|e^{\gamma s} h\|_{L^\infty(\Sigma)} < \infty$ and the symmetries (4.1). Then if R_{rot} and R_{tr} , which are the parameters in the construction associated to the dislocations, are fixed large enough, there exist β_i, τ_i , $i = 1, 2$ and ϕ a bounded solution of (4.9) such that ϕ has limit equal to zero on all wings. Moreover ϕ, β_i, τ_i depend linearly on h and*

$$(4.10) \quad \|e^{\gamma s} \phi\|_{L^\infty} + \sum_{i=1}^2 (|\beta_i| + |\tau_i|) \leq C \|e^{\gamma s} h\|_{L^\infty}.$$

Using this proposition, we fix the parameters R_{tr}, R_{rot} of the construction of the initial approximation \mathcal{M} . The following non-degeneracy property of the Jacobi operator is crucial in the proof of the above results and was proved by Montiel and Ros [24].

Proposition 4.3. *Any bounded solution ϕ of*

$$\Delta \phi + |A|^2 \phi = 0 \quad \text{in } \Sigma$$

is a linear combination of $\nu \cdot e_x, \nu \cdot e_y$ and $\nu \cdot e_z$.

The rest of the section is devoted to prove Propositions 4.1 and 4.2. We start by considering the problem

$$(4.11) \quad \begin{cases} \Delta \phi + |A|^2 \phi = h & \text{in } \Sigma_R \\ \phi = 0 & \text{on } \partial \Sigma_R, \quad \phi \text{ satisfies (4.1)} \end{cases}$$

where $R > 0$ is large and

$$(4.12) \quad \Sigma_R \text{ is the union of the core of } \Sigma \text{ and } \cup_{i=1}^4 W^i(\alpha) \cap \{s \leq R\}.$$

In the sequel, we work with $R \gg R_0 + 1$.

Lemma 4.4. *Let $0 < \gamma < 1$. Let h be defined in Σ_R , satisfy the symmetries (4.1), and $\|e^{\gamma s} h\|_{L^\infty(\Sigma)} < \infty$. Then there are R_1, C such that for all $R \geq R_1$ and any solution ϕ of (4.11) such that*

$$(4.13) \quad \int_{\Sigma_R} \phi \eta_0 z_i = 0 \quad i = 1, 2,$$

we have

$$(4.14) \quad \|\phi\|_{L^\infty(\Sigma_R)} \leq C \|e^{\gamma s} h\|_{L^\infty(\Sigma_R)}.$$

Proof. We proceed by contradiction and assume that for any positive integer n , there are R_n, ϕ_n, h_n such that $R_n \rightarrow \infty$ as $n \rightarrow \infty$, (4.11), (4.13) hold and

$$(4.15) \quad \|\phi_n\|_{L^\infty(\Sigma_{R_n})} = 1, \quad \|e^{\gamma s} h_n\|_{L^\infty(\Sigma_{R_n})} \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

Let us show first that $\phi_n \rightarrow 0$ uniformly on compact sets of Σ_{R_n} . Otherwise, up to a subsequence and using standard local estimates for elliptic equations, $\phi_n \rightarrow \phi$ uniformly on compact sets of Σ , where ϕ is bounded, not identically zero, and satisfies

$$\Delta\phi + |A|^2\phi = 0 \quad \text{in } \Sigma.$$

By Proposition 4.3, ϕ is a linear combination of $\nu \cdot e_x, \nu \cdot e_y$ and $\nu \cdot e_z$. But $\nu \cdot e_z$ is not present in this linear combination by the imposed symmetry (4.1), so

$$\phi = c_1\nu \cdot e_x + c_2\nu \cdot e_y$$

for some c_1, c_2 . But passing to the limit in (4.13), we deduce that ϕ satisfies (4.13). This implies that $c_1 = c_2 = 0$.

We note that $\psi = 1 - e^{-\gamma s}$ satisfies

$$\Delta\psi + |A|^2\psi \leq -Ce^{-\gamma s}$$

for some $C > 0$ and $s \geq s_0$ where s_0 is large enough. Using these barriers on each wing, we obtain that $\|\phi_n\|_{L^\infty(\Sigma_{R_n})} \rightarrow 0$ as $n \rightarrow \infty$, which contradicts (4.15). \square

With almost the same argument we can prove the next result.

Lemma 4.5. *Let $0 < \gamma < 1$. Let h be defined in Σ , satisfy the symmetries (4.1), and $\|e^{\gamma s} h\|_{L^\infty(\Sigma)} < \infty$. Then there is a constant C such that for any bounded solution ϕ of*

$$(4.16) \quad \begin{cases} \Delta\phi + |A|^2\phi = h & \text{in } \Sigma \\ \phi \text{ satisfies (4.1)} \end{cases}$$

which also satisfies (4.5), we have

$$\|\phi\|_{L^\infty(\Sigma)} \leq C\|e^{\gamma s} h\|_{L^\infty(\Sigma)}.$$

Proof. The proof changes only in the last step, when we use maximum principle to prove that $\phi \leq C\psi$. We change slightly the barriers by

$$\psi + \delta Z,$$

where Z is an element in the kernel of the Jacobi operator that grows linearly, and then take $\delta \rightarrow 0$. \square

Lemma 4.6. *Let $0 < \gamma < 1$. Let h be defined in Σ , satisfy the symmetries (4.1), and $\|e^{\gamma s} h\|_{L^\infty(\Sigma)} < \infty$. Suppose ϕ is a bounded solution of (4.16). Then*

$$|\nabla\phi| \leq C(\|e^{\gamma s} h\|_{L^\infty(\Sigma)} + \|\phi\|_{L^\infty(\Sigma)})e^{-\gamma s}.$$

Proof. Changing variables, we rewrite the equation on a fixed wing as

$$(4.17) \quad \Delta\phi = a_0(s, z)\phi + a_1(s, z)\nabla\phi + a_2(s, z)D^2\phi + h \quad \text{in } S$$

with boundary conditions

$$\frac{\partial\phi}{\partial z} = 0 \quad \text{on } (s_0, \infty) \times \{0, \pi\},$$

where S is the strip $(s_0, \infty) \times (0, \pi)$. We write the variables in S as (s, z) , $s > s_0$, $z \in (0, \pi)$ and in (4.17), $\Delta = \partial_{ss} + \partial_{zz}$. The functions a_0, a_1, a_2 are smooth and have the decay

$$|a_i(s, z)| \leq C e^{-s}.$$

For $T > s_0$, we have

$$\|\phi\|_{L^2((T, T+5) \times (0, \pi))} \leq C \|\phi\|_{L^\infty(S)}.$$

Since the coefficients of a_i are small as $s \rightarrow \infty$, we have from standard estimates

$$\|\phi\|_{H^2((T, T+1) \times (0, \pi))} \leq C(\|\phi\|_{L^\infty(S)} + \|e^{\gamma s} h\|_{L^\infty(S)}).$$

Hence,

(4.18)

$$\|a_0\phi + a_1\nabla\phi + a_2D^2\phi + h\|_{L^2((T, \infty) \times (0, \pi))} \leq C(\|\phi\|_{L^\infty(S)} + \|e^{\gamma s} h\|_{L^\infty(S)})e^{-\gamma T}.$$

Let $g = a_0\phi + a_1\nabla\phi + a_2D^2\phi + h$ and write

$$\phi(s, z) = \sum_{n=0}^{\infty} \phi_n(s) \cos(nz), \quad g(s, z) = \sum_{n=0}^{\infty} g_n(s) \cos(nz),$$

where, for $n \geq 0$, $\phi_n(s) = \frac{2}{\pi} \int_0^\pi \phi(s, z) \cos(nz) dz$ and $g_n(s) = \frac{2}{\pi} \int_0^\pi g(s, z) \cos(nz) dz$. We can write

$$(4.19) \quad \phi_0(s) = b_0 + \int_s^\infty (t-s)g_0(t) dt,$$

where $b_0 = \lim_{s \rightarrow \infty} \phi_0(s)$. The claim is that

$$(4.20) \quad \|\phi - b_0\|_{L^2((T, \infty) \times (0, \pi))} \leq C(\|\phi\|_{L^\infty(S)} + \|e^{\gamma s} h\|_{L^\infty(S)})e^{-\gamma T}.$$

To prove this, let $\tilde{\phi}(s, z) = \sum_{n=1}^{\infty} \phi_n(s) \cos(nz)$. We claim that, for T large

$$(4.21) \quad \|\tilde{\phi}\|_{L^2((T, \infty) \times (0, \pi))} \leq C(\|\phi\|_{L^\infty(S)} + \|e^{\gamma s} h\|_{L^\infty(S)})e^{-\gamma T}.$$

Indeed, we have

$$\phi_n'' - n^2\phi_n = g_n \quad \text{for } s > s_0.$$

Note that $\phi_n(s)$ is bounded as $s \rightarrow \infty$, so that $\phi_n(s)$ must have the form (for $n \geq 1$)

$$\phi_n(s) = d_n e^{-n(s-s_0)} + \phi_{0,n}(s),$$

where

$$d_n = \phi_n(s_0), \quad \phi_{0,n}(s) = -e^{-nd} \int_{s_0}^s e^{2nt} \int_t^\infty g_n(\tau) e^{-n\tau} d\tau dt.$$

Let $\gamma < a < 1$ be fixed. By the Cauchy-Schwarz inequality, for $n \geq 1$, we have

$$(4.22) \quad |\phi_{0,n}(s)| \leq \frac{1}{(4n(n-a))^{1/2}} e^{-as} \left(\int_{s_0}^s e^{2at} \int_t^\infty |g_n(\tau)|^2 d\tau dt \right)^{1/2}.$$

Therefore

$$\|\tilde{\phi}\|_{L^2((T, \infty) \times (0, \pi))}^2 \leq C e^{-2T} \sum_{n=1}^{\infty} d_n^2 + C \sum_{n=1}^{\infty} \int_T^\infty |\phi_{0,n}(s)|^2 ds,$$

and using (4.18), (4.22), we deduce (4.21). The estimate above and a similar one for the integral in (4.19) give (4.20).

Note that $\bar{\phi} = \phi - b_0$ satisfies

$$\Delta \bar{\phi} = a_0(s, z)\bar{\phi} + a_1(s, z)\nabla\bar{\phi} + a_2(s, z)D^2\bar{\phi} + h + a_0(s, z)b_0 \quad \text{in } S.$$

From standard elliptic estimates, we get

$$\begin{aligned} \|\bar{\phi} + |\nabla\bar{\phi}|\|_{L^\infty((T+1, T+2) \times (0, \pi))} &\leq C(\|\bar{\phi}\|_{L^2((T, T+3) \times (0, \pi))} \\ &\quad + \|h + a_0 b_0\|_{L^\infty((T, T+3) \times (0, \pi))}) \\ &\leq C(\|\phi\|_{L^\infty(S)} + \|h\|_{L^\infty(S)})e^{-\gamma T}. \quad \square \end{aligned}$$

To prove existence, let Σ_R be defined by (4.12) and consider the truncated problem of finding ϕ and c_1, c_2 such that

$$(4.23) \quad \begin{cases} \Delta\phi + |A|^2\phi = \sum_{i=1}^2 c_i \eta_0 z_i + h & \text{in } \Sigma_R, \\ \phi = 0 & \text{on } \partial\Sigma_R, \\ \phi \text{ satisfies (4.1).} \end{cases}$$

Lemma 4.7. *Let $0 < \gamma < 1$. Let h be defined in Σ_R , satisfy the symmetries (4.1), and $\|e^{\gamma s} h\|_{L^\infty(\Sigma_R)} < \infty$. Then there are unique $c_1, c_2 \in \mathbb{R}$ and $\phi \in L^\infty(\Sigma_R)$ satisfying (4.23) and (4.5). Moreover,*

$$(4.24) \quad |c_1| + |c_2| + \|\phi\|_{L^\infty(\Sigma_R)} \leq C\|e^{\gamma s} h\|_{L^\infty(\Sigma_R)},$$

with C independent of R .

Proof. We prove first (4.24). Indeed, by Lemma 4.4,

$$\|\phi\|_{L^\infty} \leq C(|c_1| + |c_2| + \|e^{\gamma s} h\|_{L^\infty}).$$

Multiplying by z_j and integrating in Σ_R , we find

$$(4.25) \quad \int_{\partial\Sigma_R} \frac{\partial\phi}{\partial\nu} z_j = c_j \int_{\Sigma_R} \eta_0 z_j^2.$$

We claim that

$$(4.26) \quad \int_{\partial\Sigma_R} \frac{\partial\phi}{\partial\nu} z_j = o(1)(\|\phi\|_{L^\infty} + \|e^{\gamma s} h\|_{L^\infty})$$

where $o(1) \rightarrow 0$ as $R \rightarrow \infty$. Combining (4.26) and (4.25), we get (4.24). To prove (4.26), we rewrite the equation on the i th wing as

$$L\phi = \sum_{i=1}^2 c_i \eta_0 z_i + h \quad \text{in } S_R,$$

where $L\phi = \Delta\phi + a_0\phi + a_1\nabla\phi + a_2 D^2\phi$ and $S_R = \{(s, z) : 0 < s < R, 0 < z < \pi\}$. Here $\Delta = \partial_{ss} + \partial_{zz}$ and $a_i(s, z)$ are smooth with $|a_i(s, z)| \leq Ce^{-s}$. Let $R_1 \gg R_0$ and $R \gg R_1$. The function $\bar{\phi} = \left(\frac{R-s}{R-R_1}\right)^\mu$, where $0 < \mu < 1$ is fixed, satisfies

$$L\bar{\phi} \leq -c\left(\frac{R-s}{R-R_1}\right)^{\mu-2}$$

in $(R_1, R) \times (0, \pi)$ for some $c > 0$, if we take R_1 large. By the maximum principle, $|\phi| \leq C(\|\phi\|_{L^\infty} + \|e^{\gamma s} h\|_{L^\infty})\bar{\phi}$ in $(R_1, R) \times (0, \pi)$. It follows that

$$\left|\frac{\partial\phi}{\partial s}(R, z)\right| \leq \frac{C}{R-R_1}(\|\phi\|_{L^\infty} + \|e^{\gamma s} h\|_{L^\infty}),$$

and this proves (4.26).

For the existence of a solution of (4.24), let us define the Hilbert space

$$H = \left\{ \phi \in H^1(\Sigma_R \cap \{z \in (0, \pi)\}) : \phi = 0 \text{ on } s = R, \int_{\Sigma_R \cap \{z \in (0, \pi)\}} \phi \eta_0 z_i = 0, i = 1, 2 \right\}$$

with the inner product

$$\langle \varphi_1, \varphi_2 \rangle = \int_{\Sigma_R \cap \{z \in (0, \pi)\}} \nabla \varphi_1 \nabla \varphi_2$$

where η_0 is the cut-off function with properties (4.6). We consider the following weak form of the equation to find $\phi \in H$:

$$(4.27) \quad \int_{\Sigma_R \cap \{z \in (0, \pi)\}} \nabla \phi \cdot \nabla \varphi - |A|^2 \phi \varphi = - \int_{\Sigma_R \cap \{z \in (0, \pi)\}} h \varphi \quad \forall \varphi \in H.$$

Let $T : H \rightarrow H$, $T(\phi) = \psi$ be the linear operator defined by the Riesz theorem from the relation

$$\langle \psi, \varphi \rangle = \int_{\Sigma_R \cap \{z \in (0, \pi)\}} |A|^2 \phi \varphi \quad \forall \varphi \in H.$$

Then T is compact and we can formulate (4.27) as finding $\phi \in H$ such that

$$(4.28) \quad \phi = T(\phi) + L_h,$$

where $L_h(\varphi) = \int_{\Sigma_R \cap \{z \in (0, \pi)\}} h \varphi$. By the Fredholm theorem, this problem is uniquely solvable for any h provided the only solution of $\phi = T(\phi)$ in H is $\phi = 0$. This holds by (4.24). \square

We now turn our attention to the problem on the whole Scherk surface Σ .

Proof of Proposition 4.1. First, let us show that, for any bounded solution ϕ, c_1, c_2 of (4.4) satisfying (4.5), the estimate (4.7) is valid. Indeed, by Lemma 4.5,

$$(4.29) \quad \|\phi\|_{L^\infty} \leq C(|c_1| + |c_2| + \|e^{\gamma s} h\|_{L^\infty}).$$

For $R \gg 1$, let $\Sigma_R = \Sigma \cap \{(x, y, z) : x^2 + y^2 \leq R^2\}$. Multiplying (4.4) by z_j and integrating in $\Sigma_R \cap \{z \in (0, \pi)\}$ we have

$$(4.30) \quad \int_{\partial \Sigma_R \cap \{z \in (0, \pi)\}} \frac{\partial \phi}{\partial \nu} z_j = c_j \int_{\Sigma_R \cap \{z \in (0, \pi)\}} \eta_0 z_j^2 + \int_{\Sigma_R \cap \{z \in (0, \pi)\}} h z_j$$

because $\int_{\Sigma_R \cap \{z \in (0, \pi)\}} \eta_0 z_1 z_2 = 0$ by symmetry. By Lemma 4.6,

$$(4.31) \quad \int_{\partial \Sigma_R \cap \{z \in (0, \pi)\}} \left| \frac{\partial \phi}{\partial \nu} z_i \right| \leq C e^{-\gamma R} (|c_1| + |c_2| + \|e^{\gamma s} h\|_{L^\infty} + \|\phi\|_{L^\infty}).$$

Combining (4.29), (4.30), and (4.31), we deduce (4.7) and prove the uniqueness.

By Lemma 4.7, for R large, the problem (4.23) is uniquely solvable, yielding a solution to ϕ_R (and constants $c_{1,R}$ and $c_{2,R}$), which remains bounded as $R \rightarrow \infty$ by (4.24). By standard elliptic estimates, ϕ_R converges locally uniformly in Σ to a solution of (4.4). Estimate (4.8) follows from Lemma 4.6. \square

Proof of Proposition 4.2. Given h as stated, let ϕ, c_1, c_2 be the solution to (4.4) provided by Proposition 4.1. Let $L_i = \lim_{s \rightarrow \infty} \phi(s, z)$ be the limit of ϕ on wing i . By adding appropriate multiples of z_1 and z_2 , we can make two of these limits equal

to zero. More precisely, by the choice of orientation (4.2) on Σ and the definition of z_1 and z_2 in (4.3), we have

$$\lim_{s \rightarrow \infty} z_1 = \begin{cases} -\sin \alpha & \text{on wing 1,} \\ \sin \alpha & \text{on wing 2,} \\ \sin \alpha & \text{on wing 3,} \\ -\sin \alpha & \text{on wing 4,} \end{cases} \quad \lim_{s \rightarrow \infty} z_2 = \begin{cases} \cos \alpha & \text{on wing 1,} \\ \cos \alpha & \text{on wing 2,} \\ -\cos \alpha & \text{on wing 3,} \\ -\cos \alpha & \text{on wing 4.} \end{cases}$$

Let d_1, d_2 satisfy

$$d_1 \sin \alpha + d_2 \cos \alpha = L_2, \quad d_1 \sin \alpha - d_2 \cos \alpha = L_3.$$

Then $\tilde{\phi} = \phi - d_1 z_1 - d_2 z_2$ satisfies (4.4) and has limit equal to zero on wings 2 and 3.

We remark that we could also achieve limit equal to zero on any two adjacent wings, but not on opposite ones in general. Also note that if the original ϕ satisfies the orthogonality conditions (4.5), the new $\tilde{\phi}$ does not in general.

Let $\tilde{\eta}_i$, $i = 1$ or $i = 4$, be smooth cut-off functions on Σ such that:

$$\begin{aligned} \tilde{\eta}_i &= 1 \text{ on wing } i \text{ and for } s \geq R_c + 1, \\ \tilde{\eta}_i &= 0 \text{ on wing } i \text{ for } 0 \leq s \leq R_c, \\ \tilde{\eta}_i &= 0 \text{ on the core and the rest of the wings.} \end{aligned}$$

Here $R_c > 0$ is a large constant to be fixed later. Define

$$\tilde{z}_1 = \nu \cdot \nu_\alpha, \quad \tilde{z}_4 = \nu \cdot (-\sin \alpha, -\cos \alpha, 0),$$

where $\nu_\alpha = (-\sin \alpha, \cos \alpha, 0)$ is the normal vector to the asymptotic plane of wing 1. Note that \tilde{z}_1, \tilde{z}_4 are in the kernel of $\Delta + |A|^2$, $\tilde{z}_1 \rightarrow 1$ as $s_1 \rightarrow \infty$, and $\tilde{z}_4 \rightarrow 1$ as $s_4 \rightarrow \infty$ (they are convenient linear combinations of z_1, z_2). Define

$$\hat{\phi} = \tilde{\phi} - L_1 \tilde{\eta}_1 \tilde{z}_1 - L_4 \tilde{\eta}_4 \tilde{z}_1.$$

Then $\hat{\phi}$ satisfies

$$\Delta \hat{\phi} + |A|^2 \hat{\phi} = h + \sum_{i=1}^2 c_i \eta_0 z_i - \sum_{i=1,4} L_i (\Delta + |A|^2) (\tilde{\eta}_i \tilde{z}_i) \quad \text{in } \Sigma$$

and $\hat{\phi}$ has limit equal to zero on all the wings. Moreover $\hat{\phi}$ satisfies the symmetries (4.1).

Suppose that β_i, τ_i , $i = 1, 2$, are given and let us consider the function $\hat{\phi}$ constructed previously with h replaced by $h + \sum_{i=1}^2 \beta_i w'_i + \tau_i w_i$. This $\hat{\phi}$ satisfies

$$(4.32) \quad \Delta \hat{\phi} + |A|^2 \hat{\phi} = h + \sum_{i=1}^2 c_i \eta_0 z_i - \sum_{i=1,4} L_i (\Delta + |A|^2) (\tilde{\eta}_i \tilde{z}_i) + \sum_{i=1}^2 (\beta_i w'_i + \tau_i w_i)$$

in Σ , has the symmetries (4.1) and the limits on all wings are equal to zero. In this construction, h, β_i, τ_i , $i = 1, 2$ are data and $\hat{\phi}, c_i$ ($i = 1, 2$), L_i ($i = 1, 4$) are bounded linear functions of these data.

We claim that there is a unique choice of β_i, τ_i , $i = 1, 2$, such that $c_1 = c_2 = L_1 = L_4 = 0$. To prove this, we test (4.32) with functions, which are going to be linear combinations of z_1, z_2 and

$$z_3 = \nu \cdot (-y, x, 0), \quad z_4 = \frac{\partial_\alpha S}{|\nabla_{x,y,z} S|},$$

where

$$S(x, y, z, \alpha) = \cos^2(\alpha) \cosh\left(\frac{x}{\cos \alpha}\right) - \sin^2(\alpha) \cosh\left(\frac{y}{\sin \alpha}\right) - \cos(z)$$

is the function (2.6) defining the Scherk surface, and $\nabla_{x,y,z} S = (\partial_x S, \partial_y S, \partial_z S)$. Note that z_3 arises from a rotation about the z -axis and z_4 is generated by the motion in α of the Scherk surfaces $\Sigma(\alpha)$, so these functions are in the kernel of the Jacobi operator $\Delta + |A|^2$ and have the symmetries (4.1). Also, z_3 and z_4 have linear growth.

Since $\hat{\phi}$ has exponential decay,

$$\int_{\Sigma \cap \{z \in (0, \pi)\}} (\Delta \hat{\phi} + |A|^2 \hat{\phi}) z_i = \int_{\Sigma \cap \{z \in (0, \pi)\}} (\Delta z_i + |A|^2 z_i) \hat{\phi} = 0,$$

for all $1 \leq i \leq 4$, while $\int_{\Sigma \cap \{z \in (0, \pi)\}} (RHS) z_i$ becomes an affine function of the numbers $c_1, c_2, L_1, L_4, \beta_1, \beta_2, \tau_1, \tau_2$. More precisely, we obtain

$$0 = \begin{bmatrix} \int h z_1 \\ \int h z_2 \\ \int h z_3 \\ \int h z_4 \end{bmatrix} + M_1 \begin{bmatrix} c_1 \\ c_2 \\ L_1 \\ L_4 \end{bmatrix} + M_2 \begin{bmatrix} \beta_1 \\ \beta_2 \\ \tau_1 \\ \tau_2 \end{bmatrix}.$$

The claim is that we can choose $\beta_1, \beta_2, \tau_1, \tau_2$ to achieve $c_1 = c_2 = L_1 = L_4 = 0$. For this, we will verify that M_1, M_2 are invertible (choosing adequately the parameters).

Note that, by symmetry,

$$\int_{\Sigma \cap \{z \in (0, \pi)\}} \eta_0 z_i z_j = c \delta_{ij}, \quad \text{for } i, j = 1, 2,$$

where $c > 0$ is some constant. Also thanks to symmetry, we have for $i = 1, 2$,

$$\int_{\Sigma \cap \{z \in (0, \pi)\}} \eta_0 z_i z_3 = 0, \quad \int_{\Sigma \cap \{z \in (0, \pi)\}} \eta_0 z_i z_4 = 0.$$

Let us compute for $i = 1$ or $i = 4$, and $j = 1, 2$:

$$\begin{aligned} \int_{\Sigma \cap \{z \in (0, \pi)\}} (\Delta + |A|^2)(\tilde{\eta}_i \tilde{z}_i) z_j &= \int_{\Sigma \cap \{z \in (0, \pi)\}} (2 \nabla \tilde{\eta}_i \cdot \nabla \tilde{z}_i + \tilde{z}_i \Delta \tilde{\eta}_i) z_j \\ &= \int_{\Sigma \cap \{z \in (0, \pi)\}} \nabla \cdot (\tilde{z}_i^2 \nabla \tilde{\eta}_i) \frac{z_j}{\tilde{z}_i} \\ &= - \int_{\Sigma \cap \{z \in (0, \pi)\}} \tilde{z}_i^2 \nabla \tilde{\eta}_i \nabla \left(\frac{z_j}{\tilde{z}_i} \right) = O(e^{-Rc}), \end{aligned}$$

where the last equality holds because \tilde{z}_i, z_j approach constants at an exponential rate.

For $i = 1, 4$, by the same computation, and using that z_3 has linear growth (for example $z_3 = \sin(\alpha)y + \cos(\alpha)x + O(se^{-s})$ on wing 1, where $s = \sqrt{x^2 + y^2} + O(1)$), we obtain

$$\begin{aligned} \int_{\Sigma \cap \{z \in (0, \pi)\}} (\Delta + |A|^2)(\tilde{\eta}_i \tilde{z}_i) z_3 &= - \int_{\Sigma \cap \{z \in (0, \pi)\}} \tilde{z}_i^2 \nabla \tilde{\eta}_i \nabla \left(\frac{z_3}{\tilde{z}_i} \right) \\ &= -\bar{B}(-1)^{i-1} + o(1) \end{aligned}$$

where $\bar{B} > 0$ and $o(1) \rightarrow 0$ as $R_c \rightarrow \infty$. Similarly, for $i = 1, 4$,

$$\begin{aligned} \int_{\Sigma \cap \{z \in (0, \pi)\}} (\Delta + |A|^2)(\tilde{\eta}_i \tilde{z}_i) z_4 &= - \int_{\Sigma \cap \{z \in (0, \pi)\}} \tilde{z}_i^2 \nabla \tilde{\eta}_i \nabla \left(\frac{z_4}{\tilde{z}_i} \right) \\ &= B + o(1) \end{aligned}$$

where $B > 0$. This implies that the matrix M_1 is invertible if R_c is taken large (and fixed).

Let us now estimate the matrix M_2 . This means, we need to estimate the following integrals for $i = 1, 2, j = 1, \dots, 4$:

$$\int_{\Sigma \cap \{z \in (0, \pi)\}} w_i z_j, \quad \int_{\Sigma \cap \{z \in (0, \pi)\}} w'_i z_j.$$

Using (2.11), for $j = 1, 2$, we have

$$\int_{\Sigma \cap \{z \in (0, \pi)\}} w_1 z_j = \int_{\Sigma \cap \{z \in (0, \pi)\}} (\Delta + |A|^2)(\eta_{tr,1}) z_j = O(e^{-R_{tr}})$$

and similarly $\int w_2 z_j = O(e^{-R_{tr}})$. Next, we compute

$$\begin{aligned} \int_{\Sigma \cap \{z \in (0, \pi)\}} w_1 z_3 &= \int_{\Sigma \cap \{z \in (0, \pi)\}} (\Delta + |A|^2)(\eta_{tr,1}) z_3 \\ &= \int_{\Sigma \cap \{z \in (0, \pi)\}} z_3 \Delta \eta_{tr,1} + \int_{\Sigma \cap \{z \in (0, \pi)\}} |A|^2 \eta_{tr,1} z_3. \end{aligned}$$

The first integral is supported on $R_{tr} \leq s \leq R_{tr} + 1$, so integrating by parts gives

$$\begin{aligned} \int_{\Sigma \cap \{z \in (0, \pi)\}} z_3 \Delta \eta_{tr,1} &= - \int_{\{R_{tr} \leq s \leq R_{tr} + 1\}} \nabla z_3 \nabla \eta_{tr,1} \\ &= - \nabla z_3 \hat{n}|_{s=R_{tr}+1} + \int_{\{R_{tr} \leq s \leq R_{tr} + 1\}} \eta_{tr,1} \Delta z_3, \end{aligned}$$

where \hat{n} is tangent to Σ and perpendicular to the curve $s = R_{tr} + 1$. Therefore

$$\begin{aligned} \int_{\Sigma \cap \{z \in (0, \pi)\}} w_1 z_3 &= - \nabla z_3 \hat{n}|_{s=R_{tr}+1} + \int_{\{s \geq R_{tr} + 1\}} |A|^2 z_3 \\ &= -\pi + o(1), \end{aligned}$$

where $o(1) \rightarrow 0$ as $R_{tr} \rightarrow \infty$, thanks to the behavior $z_3 = s + O(1)$ as $s \rightarrow \infty$ and a corresponding estimate for the its derivative. Similarly,

$$\begin{aligned} \int_{\Sigma \cap \{z \in (0, \pi)\}} w_2 z_3 &= -\pi + o(1), \\ \int_{\Sigma \cap \{z \in (0, \pi)\}} w_1 z_4 &= -\pi + o(1), \quad \int_{\Sigma \cap \{z \in (0, \pi)\}} w_2 z_4 = \pi + o(1). \end{aligned}$$

Let us now compute

$$\int_{\Sigma \cap \{z \in (0, \pi)\}} w'_1 z_1 = \int_{\Sigma \cap \{z \in (0, \pi)\}} (\Delta + |A|^2)(\eta_{rot,1} z_3) z_1,$$

where we have used (2.10). We have

$$\begin{aligned} \int_{\Sigma \cap \{z \in (0, \pi)\}} (\Delta + |A|^2)(\eta_{rot,1} z_3) z_1 &= \int_{\Sigma \cap \{z \in (0, \pi)\}} (z_3 \Delta \eta_{rot,1} + 2 \nabla \eta_{rot,1} \nabla z_3) z_1 \\ &= \int_{\Sigma \cap \{z \in (0, \pi)\}} z_1 \nabla \eta_{rot,1} \nabla z_3 - z_3 \nabla \eta_{rot,1} \nabla z_1 \\ &= -\pi \sin(\alpha) + o(1), \end{aligned}$$

where $o(1) \rightarrow 0$ as $R_{rot} \rightarrow 0$. In a similar way,

$$\begin{aligned} \int_{\Sigma \cap \{z \in (0, \pi)\}} w'_1 z_2 &= \pi \cos(\alpha) + o(1), \\ \int_{\Sigma \cap \{z \in (0, \pi)\}} w'_2 z_1 &= \pi \sin(\alpha) + o(1), \quad \int_{\Sigma \cap \{z \in (0, \pi)\}} w'_2 z_2 = \pi \cos(\alpha) + o(1). \end{aligned}$$

We have also

$$\int_{\Sigma \cap \{z \in (0, \pi)\}} w'_1 z_3 = 0, \quad \int_{\Sigma \cap \{z \in (0, \pi)\}} w'_2 z_3 = 0.$$

Indeed, consider

$$\begin{aligned} \int_{\Sigma \cap \{z \in (0, \pi)\}} w'_1 z_3 &= \int_{\Sigma \cap \{z \in (0, \pi)\}} (\Delta + |A|^2)(\eta_{rot,1} z_3) z_3 \\ &= \int_{\Sigma \cap \{z \in (0, \pi)\}} (z_3 \Delta \eta_{rot,1} + 2 \nabla \eta_{rot,1} \nabla z_3) z_3 \\ &= \int_{\Sigma \cap \{z \in (0, \pi)\}} \nabla \cdot (z_3^2 \nabla \eta_{rot,1}) \\ &= 0. \end{aligned}$$

The integral $\int w'_2 z_3 = 0$ is computed similarly.

Finally, we observe that

$$\int_{\Sigma \cap \{z \in (0, \pi)\}} w'_1 z_4 = O(1), \quad \int_{\Sigma \cap \{z \in (0, \pi)\}} w'_2 z_4 = O(1),$$

as $R_{rot} \rightarrow \infty$. Then

$$M_2 = \begin{bmatrix} \int w_1 z_1 & \int w_1 z_2 & \int w_1 z_3 & \int w_1 z_4 \\ \int w_2 z_1 & \int w_2 z_2 & \int w_2 z_3 & \int w_2 z_4 \\ \int w'_1 z_1 & \int w'_1 z_2 & \int w'_1 z_3 & \int w'_1 z_4 \\ \int w'_2 z_1 & \int w'_2 z_2 & \int w'_2 z_3 & \int w'_2 z_4 \end{bmatrix} = \begin{bmatrix} 0 & 0 & -\pi & -\pi \\ 0 & 0 & -\pi & \pi \\ -\pi \sin(\alpha) & \pi \cos(\alpha) & 0 & O(1) \\ \pi \sin(\alpha) & \pi \cos(\alpha) & 0 & O(1) \end{bmatrix} + o(1)$$

This shows that M_2 is invertible if we fix both $R_{rot} + 10 < R_{tr}$ large, which finishes the proof. \square

5. LINEAR THEORY

Let $\mathcal{E} = \mathcal{M} \cap \{s \geq \frac{\delta_s}{3\varepsilon}\}$. In this section we construct a right inverse of the operator $\Delta + |A|^2 + \varepsilon \partial_z$ on \mathcal{E} . More precisely we want, given h defined on \mathcal{E} with some decay, a solution ϕ of

$$(5.1) \quad \Delta \phi + |A|^2 \phi + \varepsilon \partial_z \phi = h \quad \text{on } \mathcal{E}.$$

Given $\alpha, \gamma \in (0, 1)$, let us define the following norms:

$$(5.2) \quad \|\phi_2\|_{*,\mathcal{E}} = \varepsilon^2 \sup_{x \in \mathcal{E}} e^{\gamma\delta_s/\varepsilon + \gamma\varepsilon s(x)} \|\phi_2\|_{C^{2,\alpha}(\bar{B}_1(x))}$$

$$(5.3) \quad \|h_2\|_{**,\mathcal{E}} = \sup_{x \in \mathcal{E}} e^{\gamma\delta_s/\varepsilon + \gamma\varepsilon s(x)} \|h_2\|_{C^{2,\alpha}(\bar{B}_1(x))},$$

where $B_1(x)$ is the geodesic ball with center x and radius 1, and s is the function defined in the construction of \mathcal{M} , Section 2. The factor $e^{\gamma\delta_s/\varepsilon}$ in front of both norms is not immediately relevant; it will be useful later.

We have the following result.

Proposition 5.1. *Let $0 < \gamma < 1$. Then there is a linear operator that associates to a function h defined on \mathcal{E} with $\|h\|_{**,\mathcal{E}} < \infty$ a solution ϕ to (5.1). Moreover,*

$$\|\phi\|_{*,\mathcal{E}} \leq C \|h\|_{**,\mathcal{E}}.$$

For the proof, we scale to size one, that is, we work on $\tilde{\mathcal{E}} = \varepsilon\mathcal{E}$. Then (5.1) becomes equivalent to

$$(5.4) \quad \Delta\phi + |A|^2\phi + \partial_z\phi = \tilde{h} \quad \text{on } \tilde{\mathcal{E}},$$

with $\tilde{h}(x) = \varepsilon^{-2}h(x/\varepsilon)$.

We study the linear operator on the unbounded pieces in the following section, then in Section 5.2, we deal with the bounded piece. We point out that in the radially symmetric case a related linear theory for the Jacobi operator was developed on [9].

5.1. Linear theory on the ends. Let \mathcal{E}_u be any of the unbounded components of \mathcal{E} and $\tilde{\mathcal{E}}_u = \varepsilon\mathcal{E}_u$. We introduce coordinates on $\tilde{\mathcal{E}}_u$ as follows. Consider a curve

$$s \mapsto (\gamma_1(s), \gamma_3(s)),$$

parametrized by arc length, with $s \in [0, \infty)$ and $\gamma_1'(s) > 0$, that solves the ordinary differential equation (2.3), with initial conditions at $s = 0$ chosen to be compatible with the construction of the initial approximation \mathcal{M} in Section 2.5. These initial conditions are functions of the parameters of the construction, in particular, the parameters $\beta_1, \beta_4, \tau_1, \tau_4$ used in the dislocations are all in $[-\delta_p, \delta_p]$. We note that the s here differs from s in the construction of \mathcal{M} , Section 2, by a shift and a scaling.

Then

$$(5.5) \quad X_0 = \nu_0(s, \theta) = (\gamma_1(s) \cos(\theta), \gamma_1(s) \sin(\theta), \gamma_3(s))$$

$s \in [0, \infty)$, $\theta \in [0, 2\pi]$, parametrizes part of the catenoid \mathcal{W}_R or the paraboloid \mathcal{P} . Let

$$\nu_0(s, \theta) = (-\gamma_3'(s) \cos(\theta), -\gamma_3'(s) \sin(\theta), \gamma_1'(s))$$

be a unit normal vector. Then a parametrization of the unbounded end $\tilde{\mathcal{E}}_u$ is given by

$$(5.6) \quad X(s, \theta) = X_0(s, \theta) + \nu_0(s, \theta)f(s/\varepsilon, \theta/\varepsilon)$$

where f is essentially a cut-off function times f_α , the function that allows one to write the Scherk surface as a graph over a plane, see Lemma 2.2. The important properties of f are that $|\partial^k f(\tilde{s}, \tilde{\theta})| \leq C_k e^{-\delta_s/(10\varepsilon)}$ for some C_k and that it vanishes for $\tilde{s} \geq 10\delta_s/\varepsilon$.

Then we have the following expression for the operator $\Delta + |A|^2 + \partial_z$ in the coordinates s and θ :

$$(5.7) \quad \Delta\phi + |A|^2\phi + \partial_z\phi = \partial_{ss}\phi + \frac{1}{\gamma_1(s)^2}\partial_{\theta\theta}\phi + \left(\frac{\gamma_1'(s)}{\gamma_1(s)} + \gamma_3'(s)\right)\partial_s\phi + |A|^2\phi + \tilde{L}\phi$$

where \tilde{L} is a second order differential operator in ϕ with coefficients that are $o(1)$ as $\varepsilon \rightarrow 0$ and supported in $s \in [0, 10\delta_s]$. Using (2.5) and the fact that the principal curvatures of a surface of revolution $z = F(r)$ are given by

$$\kappa_1 = \frac{F''(r)}{(1 + F'(r)^2)^{3/2}}, \quad \kappa_2 = \frac{F'(r)}{r(1 + F'(r)^2)^{1/2}},$$

we can write

$$\Delta\phi + |A|^2\phi + \partial_z\phi = \partial_{ss}\phi + a(s)\partial_{\theta\theta}\phi + b(s)\partial_s\phi + |A|^2\phi + \tilde{L}\phi,$$

where we have the following properties of the coefficients:

$$(5.8) \quad \begin{cases} a(s) = \frac{1}{2s}(1 + O(s^{-1/2})) \\ b(s) = 1 + O(s^{-1/2}) \\ |A|^2 = \frac{1}{2s}(1 + O(s^{-1/2})) \end{cases}$$

as $s \rightarrow \infty$, and $a(s) > 0$ for all $s \geq 0$.

Let

$$L_0\phi = \partial_{ss}\phi + a(s)\partial_{\theta\theta}\phi + b(s)\partial_s\phi + |A|^2\phi,$$

and consider the equation

$$(5.9) \quad L_0\phi = h, \quad s \in (0, \infty), \quad \theta \in [0, 2\pi].$$

To prove Proposition 5.1, we will first construct an inverse operator for L_0 .

Proposition 5.2. *Let $0 < \gamma < 1$. Then there is a linear operator $h \mapsto \phi$ that associates to a function $h = h(s, \theta)$ defined for $(s, \theta) \in (0, \infty) \times [0, 2\pi]$, 2π -periodic in θ with $\|e^{\gamma s}h\|_{L^\infty} < \infty$ a solution ϕ of (5.9), which is 2π -periodic in θ and satisfies*

$$\|e^{\gamma s}\phi\|_{L^\infty} \leq C\|e^{\gamma s}h\|_{L^\infty}.$$

For the proof of this result, we will write h and ϕ in Fourier series

$$(5.10) \quad \phi(s, \theta) = \sum_{k \in \mathbb{Z}} \phi_k(s)e^{ik\theta}, \quad h(s, \theta) = \sum_{k \in \mathbb{Z}} h_k(s)e^{ik\theta}.$$

Then, if ϕ is smooth with exponential decay, equation (5.9) is equivalent to

$$(5.11) \quad \phi_k'' + b(s)\phi_k' + (|A|^2 - a(s)k^2)\phi_k = h_k, \quad \text{for all } s > 0,$$

for all $k \in \mathbb{Z}$. We need a couple of lemmas before starting the proof of Proposition 5.2. Combined, they allow us to deal with the low modes, i.e $|k| \leq k_0$ for some fixed k_0 , and focus our attention on the higher frequencies.

Lemma 5.3. *Let $0 < \gamma < 1$. If $|h_k(s)| \leq Ce^{-\gamma s}$, then (5.11) has a unique solution ϕ_k with*

$$\|e^{\gamma s}\phi_k\|_{L^\infty} \leq C_k\|e^{\gamma s}h_k\|_{L^\infty}.$$

Proof. Note that as $s \rightarrow \infty$, equation (5.11) with $h_k = 0$ is asymptotic to

$$(5.12) \quad \phi_k'' + \phi_k' + \frac{b_k}{s} \phi_k = 0,$$

where $b_k = \frac{1-k^2}{2}$. Note that $b_0 = \frac{1}{2}$, $b_1 = 0$, and $b_k < 0$ for $k > 1$. We see that the homogeneous equation (5.12) has two independent solutions with the behaviors $s^{-b_k}(1 + O(s^{-1}))$ and $s^{b_k}e^{-s}(1 + O(s^{-1}))$ as $s \rightarrow \infty$. Using these solutions, it is possible to construct, for each $k \geq 0$, two elements $z_{1,k}$ and $z_{2,k}$ in the kernel of the operator $\phi'' + b(s)\phi' + (|A|^2 - a(s)k^2)\phi$ such that

$$z_{1,k}(s) = s^{-b_k}(1 + O(s^{-\sigma})), \quad z_{2,k}(s) = s^{b_k}e^{-s}(1 + O(s^{-\sigma})),$$

as $s \rightarrow \infty$, where $\sigma \in (0, 1)$. Now we can construct the solution ϕ_k using the variation of parameters formula:

$$\phi_k = -z_{1,k} \int \frac{h_k z_{2,k}}{W_k} + z_{2,k} \int \frac{h_k z_{1,k}}{W_k},$$

where $W_k = z_{1,k}z_{2,k}' - z_{1,k}'z_{2,k} = e^{-s}(1 + O(s^{-\sigma}))$ and the integrals are chosen to have the desired decay. An alternative for the construction is to use the super solution $\bar{\phi} = e^{-\gamma s}$ and the calculation as in Lemma 5.4. \square

Let $k_0 \in \mathbb{N}$. Let ϕ be a bounded measurable function on $[0, \infty) \times [0, 2\pi]$. We will say that the Fourier coefficients of ϕ of order less than k_0 vanish if

$$\int_0^{2\pi} \phi(s, \theta) e^{-ik\theta} d\theta = 0, \quad \forall s > 0, \quad \forall |k| < k_0.$$

Lemma 5.4. *There is a k_0 with the following property: Suppose that ϕ is continuous, satisfies (5.9), with ϕ and h 2π -periodic in θ , that*

$$\begin{aligned} \phi(0, \theta) &= 0 \quad \forall \theta \in [0, 2\pi], \\ \|e^{\gamma s} \phi\|_{L^\infty} &< \infty, \quad \|e^{\gamma s} h\|_{L^\infty} < \infty, \end{aligned}$$

and that the Fourier coefficients of order less than k_0 of ϕ and h are zero, then there is a constant C independent of ϕ and h such that

$$\|e^{\gamma s} \phi\|_{L^\infty} \leq C \|e^{\gamma s} h\|_{L^\infty}.$$

Proof. We proceed by contradiction. Assume that the statement fails, so that there exist sequences ϕ_n, h_n such that ϕ_n, h_n are 2π -periodic in θ , with vanishing Fourier coefficients of order less than k_0 (k_0 will be fixed later), ϕ_n solves (5.9) with right hand side h_n , $\phi_n(0, \theta) = 0$ for $\theta \in [0, 2\pi]$, and

$$(5.13) \quad \|e^{\gamma s} \phi_n\|_{L^\infty} = 1, \quad \|e^{\gamma s} h_n\|_{L^\infty} \rightarrow 0, \quad \text{as } n \rightarrow \infty.$$

Consider $\bar{\phi}(s) = e^{-\gamma s}$. By (5.8) we see that

$$L_0 \bar{\phi} = (\gamma^2 - \gamma) e^{-\gamma s} (1 + O(s)) \leq \frac{\gamma^2 - \gamma}{2} e^{-\gamma s}$$

for $s \geq s_0$ (here we fix $s_0 > 0$ large). Using the maximum principle with $\bar{\phi} + \sigma e^{-\tilde{\gamma}s}$ with $0 < \tilde{\gamma} < \gamma$ and $\sigma > 0$, and then letting $\sigma \rightarrow 0$, we obtain

$$\|e^{\gamma s} \phi_n\|_{L^\infty((s_0, \infty) \times [0, 2\pi])} \leq C \|e^{\gamma s} h_n\|_{L^\infty} + C \|e^{\gamma s} \phi_n\|_{L^\infty((0, s_0) \times [0, 2\pi])}.$$

From this and (5.13), we deduce that for a subsequence (also denoted ϕ_n)

$$\|e^{\gamma s} \phi_n\|_{L^\infty((0, s_0) \times [0, 2\pi])} \geq c_0$$

for some constant $c_0 > 0$. By standard elliptic estimates, up to a new subsequence, $\phi_n \rightarrow \phi$ uniformly on compact subsets of $[0, \infty) \times [0, 2\pi]$ and by the previous remark, $\phi \not\equiv 0$. But because of (5.13), ϕ satisfies

$$L_0\phi = 0 \quad \text{in } (0, \infty) \times [0, 2\pi],$$

with $\phi(0, \theta) = 0$ and $|\phi(s)| \leq e^{-\gamma s}$. Let us write ϕ as a Fourier series (5.10). Due to the hypotheses, $\phi_k = 0$ for $|k| < k_0$. Note that ϕ_k satisfies (5.11) with right-hand side equal to 0 and $\phi_k(0) = 0$. The function $\bar{\phi}(s) = e^{-\gamma s}$ satisfies

$$\bar{\phi}'' + b(s)\bar{\phi}' + (|A|^2 - a(s)k^2)\bar{\phi} = (\gamma^2 - \gamma b(s) + |A|^2 - a(s)k^2)e^{-\gamma s}.$$

But from the fact that $a(s) > 0$ for all $s \geq 0$ and the estimates on the coefficients (5.8), we see that there is k_0 such that

$$(5.14) \quad \bar{\phi}'' + b(s)\bar{\phi}' + (|A|^2 - a(s)k^2)\bar{\phi} < -c_k e^{-\gamma s}, \quad \forall |k| \geq k_0, \forall s \geq 0,$$

where $c_k > 0$. From the maximum principle, we deduce that $\phi_k \equiv 0$ for all $|k| \geq k_0$. This is a contradiction. \square

Proof of Proposition 5.2. Let k_0 be as in Lemma 5.4. Using Lemma 5.3, for each $|k| < k_0$, we find a solution ϕ_k of (5.11) in $(0, \infty)$. Then we need to prove the lemma only under the assumption that the Fourier coefficients of order less than k_0 of h vanish.

For the moment let us assume in addition that h is C^2 and

$$(5.15) \quad |h_{\theta\theta}(s, \theta)| \leq C e^{-\gamma s} \quad \text{for all } (s, \theta) \in [0, \infty) \times [0, 2\pi].$$

Writing h in Fourier series as in (5.10) for $|k| \geq k_0$ we can find a solution ϕ_k of (5.11) in $(0, \infty)$ with right hand side h_k , satisfying $\phi_k(0) = 0$. This can be done using the supersolution $\bar{\phi}(s) = e^{-\gamma s}$ and (5.14). Alternatively, one can use the variation of parameters formula and elements in the kernel as in Lemma 5.3. For $m > k_0$, let

$$\phi_m(s, \theta) = \sum_{k_0 \leq |k| \leq m} e^{ik\theta} \phi_k(s)$$

and similarly define h_m . By Lemma 5.4,

$$\|e^{\gamma s} \phi_m\|_{L^\infty} \leq C \|e^{\gamma s} h_m\|_{L^\infty},$$

with C independent of m . Note that

$$|h_k(s)| = \frac{1}{2\pi} \left| \int_0^{2\pi} h(s, \theta) e^{ik\theta} d\theta \right| = \frac{1}{2\pi k^2} \left| \int_0^{2\pi} h_{\theta\theta}(s, \theta) e^{ik\theta} d\theta \right| \leq \frac{C}{k^2} e^{-\gamma s}.$$

Then

$$|h_m(s, \theta)| = \left| \sum_{k_0 \leq |k| \leq m} h_k(s) e^{ik\theta} \right| \leq \sum_{k_0 \leq |k| \leq m} |h_k(s)| \leq C e^{-\gamma s} \sum_{k_0 \leq |k| \leq m} k^{-2}.$$

Therefore

$$\|e^{\gamma s} \phi_m\|_{L^\infty} \leq C,$$

with C independent of m . By standard elliptic estimates, for a subsequence $m \rightarrow \infty$ we find $\phi_m \rightarrow \phi$ uniformly on compact subsets of $[0, \infty) \times [0, 2\pi]$ and ϕ is the desired solution.

Next, we lift the assumption (5.15). Indeed, assume only $\|e^{\gamma s} h\|_{L^\infty} < \infty$ and that the Fourier coefficients of h vanish for $|k| < k_0$. Let ρ_n be a sequence of mollifiers in \mathbb{R}^2 and

$$h_n = h * \rho_n$$

(extending h by 0 for $s \leq 0$). We have $h_n \rightarrow h$ a.e; the Fourier coefficients of order less than k_0 of h_n vanish and

$$|h_n(s, \theta)| \leq C e^{-\gamma s},$$

$$|\partial_{\theta\theta} h_n(s, \theta)| \leq C_n e^{-\gamma s},$$

Using the previous argument we find a solution ϕ_n with

$$\|e^{\gamma s} \phi_n\|_{L^\infty} \leq C \|e^{\gamma s} h_n\|_{L^\infty} \leq C,$$

with C independent of n and right-hand side h_n . Passing to a subsequence, we find the desired solution. \square

5.2. Linear theory on the bounded piece. Let us consider the bounded component \mathcal{E}_b of $\mathcal{E} = \mathcal{M} \setminus \{s \leq \frac{\delta_s}{3\varepsilon}\}$ and let $\tilde{\mathcal{E}}_b = \varepsilon \mathcal{E}_b$.

Lemma 5.5. *For $h \in C^\alpha(\overline{\tilde{\mathcal{E}}_b})$, there is a unique solution ϕ of*

$$(5.16) \quad \begin{cases} \Delta\phi + |A|^2\phi + \partial_z\phi = h & \text{in } \tilde{\mathcal{E}}_b \\ \phi = 0 & \text{on } \partial\tilde{\mathcal{E}}_b. \end{cases}$$

with

$$(5.17) \quad \|\phi\|_{L^\infty} \leq C \|h\|_{L^\infty},$$

where the constant C is independent of h and ϕ .

Proof. Let $L\phi$ be $\Delta\phi + |A|^2\phi + \partial_z\phi$, where the geometric quantities and Laplacian are the ones for $\tilde{\mathcal{E}}_b$. We let L_0 denote the corresponding operator for the paraboloid \mathcal{P} .

We can parametrize \mathcal{P} with polar coordinates

$$X_0(r, \theta) = (r \cos(\theta), r \sin(\theta), F(r)),$$

with $r \geq 0$, $\theta \in [0, 2\pi]$, where $F = F_0$ is the unique radially symmetric solution of (2.1) with $F(0) = 0$. Then, in the coordinates r and θ ,

$$L_0\phi = B_1(r)\phi_{rr} + B_2(r)\phi_r + \frac{1}{r^2}\phi_{\theta\theta} + A^2(r)\phi,$$

where

$$\begin{aligned} B_1(r) &= \frac{1}{1 + F'(r)^2}, \\ B_2(r) &= \frac{1}{r(1 + F'(r)^2)} - \frac{F'(r)F''(r)}{(1 + F'(r)^2)^2}, \\ A^2(r) &= \frac{F''(r)^2}{(1 + F'(r)^2)^3} + \frac{F'(r)^2}{r^2(1 + F'(r)^2)}. \end{aligned}$$

As before, we denote by ν_0 the unit normal vector to \mathcal{P} such that $\langle \nu_0, e_z \rangle > 0$. The surface $\tilde{\mathcal{E}}_b$ can then be parametrized by

$$X_0(r, \theta) + \nu_0(r, \theta)f(r/\varepsilon, \theta/\varepsilon)$$

for $r \in [0, R_1]$, $\theta \in [0, 2\pi]$, with some $R_1 > 0$, and where f has the property that $f(r/\varepsilon, \theta/\varepsilon)$ is supported where $r \in [R_1 - 10\delta_s, R_1]$ and f and its derivatives can be bounded by $e^{-\delta_s/(10\varepsilon)}$. This implies that

$$L\phi = L_0\phi + \tilde{L}\phi,$$

where \tilde{L} is a second order differential operator in ϕ with coefficients that are $o(1)$ as $\varepsilon \rightarrow 0$ and supported in $r \in [R_1 - 10\delta_s, R_1]$.

Let $v(r) = \langle \nu_0, e_z \rangle$. Then $L_0v = 0$ and $v(r) > 0$ for all $r \geq 0$. We define now

$$\bar{\phi}(r) = v(r) - \mu e^{-Kr},$$

where $\mu = \frac{1}{2} \inf_{r \in [0, R_1]} v(r) > 0$ and $K > 0$ is to be chosen. Then, we compute

$$L\bar{\phi} = \mu e^{-Ks} [-B_1(r)K^2 + KB_2(r) - A^2(r)] - \mu \tilde{L}(e^{-Ks}) + \tilde{L}v.$$

But $\tilde{L}v = O(\varepsilon)$, $\tilde{L}(e^{-Ks}) = o(1)K^2e^{-Ks}$ where in the last expression $o(1)$ is uniform in K as $\varepsilon \rightarrow 0$. Since $B_1(r)$ is positive in $[0, R_1]$ we can choose K large so that

$$L\bar{\phi} \leq -c, \quad \forall r \in [0, R_1],$$

for some $c > 0$ and all $\varepsilon > 0$ small. Then $\bar{\phi}$ is a super solution for the operator L and hence the bound (5.17) holds. The equation is solved then by super and subsolutions. \square

Proof of Proposition 5.1. Let $0 < \gamma < 1$.

Let us consider first one of the unbounded ends \mathcal{E}_u and $\tilde{\mathcal{E}}_u = \varepsilon\mathcal{E}_u$. Recall that $\tilde{\mathcal{E}}_u$ is parametrized by (5.6), in particular, $s \geq 0$, $\theta \in [0, 2\pi)$ are global coordinates on this surface. We write $s(x)$ for $x \in \tilde{\mathcal{E}}_u$. Using s, θ we may identify $\tilde{\mathcal{E}}_u$ with an unbounded piece of the paraboloid or catenoid. We write this piece as E_u .

Let $\alpha \in (0, 1)$ and define the norms for functions defined on E_u :

$$\|h\|_{k, \alpha} = \sup_{x \in E_u} e^{\gamma s(x)} \|h\|_{C^{k, \alpha}(\bar{B}_1(x))},$$

where $B_1(x)$ is the geodesic ball of center x and radius 1 in E_u .

If h is defined on E_u and $\|h\|_{0, \alpha} < \infty$, using Proposition 5.2, we obtain a solution $\phi = T_0(h)$ of (5.9), 2π -periodic in θ , and such that

$$\|e^{\gamma s} T_0(h)\|_{L^\infty} \leq C \|e^{\gamma s} h\|_{L^\infty} \leq C \|h\|_{0, \alpha}.$$

By considering ϕ and h as functions on E_u , equation (5.9) becomes

$$\Delta_{E_u} \phi + |A_{E_u}|^2 \phi + \partial_z \phi = h \quad \text{on } E_u.$$

Standard elliptic estimates, applied on geodesic balls of radius 1 of E_u , give that

$$\|\phi\|_{2, \alpha} \leq C(\|h\|_{0, \alpha} + \|\phi\|_{C^{2, \alpha}(\partial E_u)}).$$

Note that ∂E_u in the coordinates s, θ corresponds to $s = 0$. Consider the representations of ϕ and h as Fourier series as in (5.10). We recall that, by construction, if $\phi = T_0(h)$ then $\phi(0, \cdot)$ has Fourier modes of index $|k| \geq k_0$ equal to zero, where k_0 is a fixed integer. Hence

$$\|\phi\|_{C^{2, \alpha}(\partial E_u)} \leq C \sum_{|k| \leq k_0} |\phi_k(0)|.$$

The solutions ϕ_k are constructed in Lemma 5.3 and, in particular, we have $|\phi_k(0)| \leq C_k \|e^{\gamma s} h_k\|_{L^\infty}$. From this, we deduce

$$\|\phi\|_{C^{2, \alpha}(\partial E_u)} \leq C \|e^{\gamma s} h\|_{L^\infty},$$

and hence

$$(5.18) \quad \|T_0(h)\|_{2,\alpha} \leq C\|h\|_{0,\alpha}.$$

To solve equation (5.4), we rewrite it

$$(5.19) \quad \Delta_{E_u}\phi + |A_{E_u}|^2\phi + \partial_z\phi = h + \tilde{L}\phi \quad \text{on } E_u,$$

where \tilde{L} a second order elliptic equation with coefficients that are $o(1)$ as $\varepsilon \rightarrow 0$ and with compact support. This translates to

$$(5.20) \quad \|\tilde{L}\phi\|_{0,\alpha} \leq o(1)\|\phi\|_{2,\alpha}.$$

Using the operator T_0 , we can find a solution of (5.19) by solving the fixed point problem

$$\phi = T_0(h + \tilde{L}\phi)$$

in the Banach space $\{\phi \in C^{2,\alpha}(\bar{E}_u) : \|\phi\|_{2,\alpha} < \infty\}$. By (5.18), (5.20) and the contraction mapping principle, this fixed point problem has a unique solution. This yields a solution of (5.4). By scaling we obtain therefore a solution of (5.1) in any of the unbounded ends.

The proof for the bounded component \mathcal{E}_b of $\mathcal{E} = \mathcal{M} \setminus \{s \leq \frac{\delta_s}{3\varepsilon}\}$ is similar, if one uses Lemma 5.5, and the fact that the boundary condition is taken equal to 0. \square

6. PROOF OF THEOREM 1.1

To prove the theorem, it is sufficient to find a solution ϕ of (1.11), that is,

$$(6.1) \quad \Delta\phi + |A|^2\phi + \varepsilon\nabla\phi \cdot e_z + E + Q(x, \phi, \nabla\phi, D^2\phi, x) = 0 \quad \text{in } \mathcal{M},$$

where \mathcal{M} is the surface constructed in Section 2, which depends on $\beta_1, \beta_2, \tau_1, \tau_4 \in [-\delta_p, \delta_p]$, and $E = H - \varepsilon\nu \cdot e_z$. Later, we will verify that $\{x + \phi(x)\nu(x) : x \in \mathcal{M}\}$ is an embedded complete surface.

Thanks to Proposition 1.2, $E = E_0 + E_d$ with $\|E_0\|_{**} \leq C\varepsilon$ and

$$E_d = \tau_1 w_1 + \tau_4 w_2 + \beta_1 w'_1 + \beta_4 w'_2 + O\left(\sum \beta_i^2 + \tau_i^2\right),$$

where $O\left(\sum \beta_i^2 + \tau_i^2\right)$ are smooth functions with compact support, with $\|\cdot\|_{**}$ bounded by $\sum_{i=1,4} \beta_i^2 + \tau_i^2$. Thus (6.1) takes the form

$$(6.2) \quad \begin{aligned} \Delta\phi + |A|^2\phi + \varepsilon\nabla\phi \cdot e_z + \tilde{E} + Q(\phi, \nabla\phi, D^2\phi) + \tau_1 w_1 \\ + \tau_4 w_2 + \beta_1 w'_1 + \beta_4 w'_2 = 0, \quad \text{in } \mathcal{M}, \end{aligned}$$

where $\tilde{E} = E_0 + O\left(\sum_{i=1,4} \beta_i^2 + \tau_i^2\right)$. Hence,

$$\|\tilde{E}\|_{**} \leq C \left(\varepsilon + \sum_{i=1,4} (\beta_i^2 + \tau_i^2) \right).$$

Note that $\mathcal{M} = \mathcal{M}[\beta_1, \beta_2, \tau_1, \tau_2]$ and the unknowns are ϕ and $\beta_1, \beta_2, \tau_1, \tau_2$.

We look for a solution ϕ of (6.2) of the form

$$\phi = \eta_1 \phi_1 + \eta_2 \phi_2,$$

where ϕ_1, ϕ_2 are new unknown functions, which solve an appropriate system, and η_1, η_2 are smooth cut-off functions such that

$$\eta_1(s) = \begin{cases} 1 & \text{if } s \leq \frac{2\delta_s}{\varepsilon} - 1 \\ 0 & \text{if } s \geq \frac{2\delta_s}{\varepsilon} \end{cases}$$

$$\eta_2(s) = \begin{cases} 0 & \text{if } s \leq \frac{\delta_s}{3\varepsilon} \\ 1 & \text{if } s \geq \frac{\delta_s}{2\varepsilon} \end{cases},$$

where $s = s(x)$ measures geodesic distance from the core of \mathcal{M} .

We introduce next the following system for ϕ_1, ϕ_2 :

$$(6.3) \quad \Delta\phi_1 + |A|^2\phi_1 + \varepsilon\partial_z\phi_1 = -\varepsilon\phi_2\partial_z\eta_2 - 2\nabla\phi_2\nabla\eta_2 - \phi_2\Delta\eta_2 \\ - \tilde{\eta}_1(\tilde{E} + Q(x, \phi, \nabla\phi, D^2\phi)) - \tau_1w_1 + \tau_4w_2 + \beta_1w'_1 + \beta_4w'_2 \quad \text{in } \mathcal{C}$$

$$(6.4) \quad \Delta\phi_2 + |A|^2\phi_2 + \varepsilon\partial_z\phi_2 = -\varepsilon\phi_1\partial_z\eta_1 - 2\nabla\phi_1\nabla\eta_1 - \phi_1\Delta\eta_1 \\ - \tilde{\eta}_2((\tilde{E} + Q(x, \phi, \nabla\phi, D^2\phi))) \quad \text{in } \mathcal{E},$$

where $\mathcal{E} = \mathcal{M} \cap \{s \geq \frac{\delta_s}{3\varepsilon}\}$ is the union of the ends, $\mathcal{C} = \mathcal{M} \cap \{s \leq \frac{2\delta_s}{\varepsilon}\}$ is close to a Scherk surface, and $\tilde{\eta}_1, \tilde{\eta}_2$ are smooth cut-off functions such that:

$$\tilde{\eta}_1(s) = \begin{cases} 1 & \text{if } s \leq \frac{\delta_s}{\varepsilon} - 1 \\ 0 & \text{if } s \geq \frac{\delta_s}{\varepsilon} \end{cases}$$

$$\tilde{\eta}_2(s) = 1 - \tilde{\eta}_1(s).$$

In the term $Q(\phi, \nabla\phi, D^2\phi)$ of (6.3), (6.4), ϕ means $\phi = \phi_1\eta_1 + \phi_2\eta_2$. If ϕ_1, ϕ_2 solve (6.3), (6.4), then multiplying (6.3) by η_1 and (6.4) by η_2 we see that $\phi = \phi_1\eta_1 + \phi_2\eta_2$ is a solution of (6.2).

Using the change of variables introduced in the construction of \mathcal{M} in Section 2, we see that solving (6.3) is equivalent to finding a solution to

$$(6.5) \quad \Delta_\Sigma\phi_1 + |A_\Sigma|^2\phi_1 = L'\phi_1 - \varepsilon\partial_z\phi_1 - \varepsilon\phi_1\partial_z\eta_1 - \varepsilon\phi_2\partial_z\eta_2 - 2\nabla\phi_2\nabla\eta_2 - \phi_2\Delta\eta_2 \\ - \tilde{\eta}_1(\tilde{E} + Q(x, \phi, \nabla\phi, D^2\phi)) - \tau_1w_1 + \tau_4w_2 + \beta_1w'_1 + \beta_4w'_2 \quad \text{in } \Sigma,$$

where now all the functions are considered on the Scherk surface Σ ; Δ_Σ and A_Σ refer to the Laplace operator and second fundamental form of Σ . By Proposition 3.4, L' is a second order operator with coefficients supported on $s \leq 2\delta_s/\varepsilon$ and whose C^1 norms on the region $s \leq 5\delta_s/\varepsilon$ are bounded by $\delta_s + \delta_p + \varepsilon$. In principle, by the change of variables, we need to solve (6.5) on a subset of Σ , but finding a solution in all Σ is sufficient. This solution is multiplied later by the cut-off η_1 .

Similarly, we consider $\mathcal{E}_0 = \mathcal{M}_0 \cap \{s \geq \frac{\delta_s}{3\varepsilon}\}$, where \mathcal{M}_0 is the initial approximation corresponding to $\beta_1 = \beta_2 = \tau_1 = \tau_2 = 0$. For $|\beta_i| + |\tau_i| \leq \delta_p$ and $\delta_p > 0$ fixed small, \mathcal{E} is mapped onto \mathcal{E}_0 and this mapping allows us to write (6.4) as

$$(6.6) \quad \Delta_{\mathcal{E}_0}\phi_2 + |A_{\mathcal{E}_0}|^2\phi_2 + \varepsilon\partial_z\phi_2 = L''\phi_2 - 2\nabla\phi_1\nabla\eta_1 - \phi_1\Delta\eta_1 \\ - \tilde{\eta}_2((\tilde{E} + Q(x, \phi, \nabla\phi, D^2\phi))) \quad \text{in } \mathcal{E}_0,$$

where now all functions are considered on the Scherk surface \mathcal{E}_0 ; $\Delta_{\mathcal{E}_0}$ and $A_{\mathcal{E}_0}$ refer to the Laplace operator and second fundamental form of \mathcal{E}_0 . The operator L'' has coefficients whose C^1 norms are $o(\delta_p)$ as $\delta_p \rightarrow 0$. Indeed, one can see this using the

form of the operator on the ends in (5.7) and the continuous dependence of ODE on initial conditions, because the parameters β_i, τ_i determine the initial condition for the differential equation (2.3).

We solve (6.5) on the Scherk surface using Proposition 4.2 with norms

$$(6.7) \quad \begin{aligned} \|\phi_1\|_{*,\Sigma} &= \sup_{x \in \Sigma} e^{\gamma s(x)} \|\phi_1\|_{C^{2,\alpha}(\bar{B}_1(x))}, \\ \|h_1\|_{**,\Sigma} &= \sup_{x \in \Sigma} e^{\gamma s(x)} \|h_1\|_{C^\alpha(\bar{B}_1(x))}. \end{aligned}$$

We consider the system (6.4), (6.5) as a fixed point problem for $(\phi_1, \phi_2, \beta_1, \beta_2, \tau_1, \tau_2)$ belonging to the subset \mathcal{B} of $C^{2,\alpha}(\Sigma) \times C^{2,\alpha}(\mathcal{E}_0) \times \mathbb{R}^4$ defined by

$$\mathcal{B} = \{(\phi_1, \phi_2, \beta_1, \beta_2, \tau_1, \tau_2) \mid \max_{i=1,2} (\|\phi_i\|_{*,\Sigma}, |\beta_i|, |\tau_i|) \leq M\varepsilon\},$$

where $M > 0$ is a constant to be chosen later and the norms are defined in (5.2) and (6.7).

Consider $(\phi_1, \phi_2, \beta_1, \beta_2, \tau_1, \tau_2) \in \mathcal{B}$. We define $F(\phi_1, \phi_2, \beta_1, \beta_2, \tau_1, \tau_2)$ as follows. Using Proposition 4.2, we let $\bar{\phi}_1, \bar{\beta}_1, \bar{\beta}_2, \bar{\tau}_1, \bar{\tau}_2$ be the solution of

$$(6.8) \quad \begin{aligned} \Delta_\Sigma \bar{\phi}_1 + |A_\Sigma|^2 \bar{\phi}_1 &= L' \phi_1 - \varepsilon \partial_z \phi_1 - \varepsilon \phi_1 \partial_z \eta_1 - \varepsilon \phi_2 \partial_z \eta_2 - 2 \nabla \phi_2 \nabla \eta_2 - \phi_2 \Delta \eta_2 \\ &\quad - \tilde{\eta}_1 (\tilde{E} + Q(x, \phi, \nabla \phi, D^2 \phi)) - \bar{\tau}_1 w_1 + \bar{\tau}_4 w_2 + \bar{\beta}_1 w'_1 + \bar{\beta}_4 w'_2 \quad \text{in } \Sigma. \end{aligned}$$

In the right-hand side of this equation, all terms are well defined in spite of the fact that ϕ_1, ϕ_2 (after changing variables) are defined only on some subsets, because of the cut-off functions. As we will verify later, the right-hand side has finite $\|\cdot\|_{**, \Sigma}$ norm, thus Proposition 4.2 applies. Next, using Proposition 5.1, we find a solution of

$$(6.9) \quad \begin{aligned} \Delta_{\mathcal{E}_0} \bar{\phi}_2 + |A_{\mathcal{E}_0}|^2 \bar{\phi}_2 + \varepsilon \partial_z \phi_2 &= L'' \phi_2 - 2 \nabla \phi_1 \nabla \eta_1 - \phi_1 \Delta \eta_1 \\ &\quad - \tilde{\eta}_2 ((\tilde{E} + Q(x, \phi, \nabla \phi, D^2 \phi))) \quad \text{in } \mathcal{E}_0, \end{aligned}$$

where the right-hand side will be proven to have finite $\|\cdot\|_{**, \mathcal{E}_0}$ norm. We define

$$(\bar{\phi}_1, \bar{\phi}_2, \bar{\beta}_1, \bar{\beta}_2, \bar{\tau}_1, \bar{\tau}_2) = F(\phi_1, \phi_2, \beta_1, \beta_2, \tau_1, \tau_2).$$

Let us verify that F maps \mathcal{B} into \mathcal{B} and is a contraction. Consider $(\phi_1, \phi_2, \beta_1, \beta_2, \tau_1, \tau_2) \in \mathcal{B}$ and let $(\bar{\phi}_1, \bar{\phi}_2, \bar{\beta}_1, \bar{\beta}_2, \bar{\tau}_1, \bar{\tau}_2) = F(\phi_1, \phi_2, \beta_1, \beta_2, \tau_1, \tau_2)$. Using (4.10) and standard elliptic estimates, we deduce

$$(6.10) \quad \begin{aligned} \|\bar{\phi}_1\|_{*,\Sigma} + |\bar{\beta}_i| + |\bar{\tau}_i| \\ \leq C \|L' \phi_1 - \varepsilon \partial_z \phi_1 - \varepsilon \phi_1 \partial_z \eta_1 - \varepsilon \phi_2 \partial_z \eta_2 - 2 \nabla \phi_2 \nabla \eta_2 - \phi_2 \Delta \eta_2\|_{**, \Sigma} \\ \quad + \|\tilde{\eta}_1 (\tilde{E} + Q(x, \phi, \nabla \phi, D^2 \phi))\|_{**, \Sigma}. \end{aligned}$$

We first remark that by Proposition 1.2, we have

$$\|\tilde{\eta}_1 \tilde{E}\|_{**, \Sigma} \leq C_E \varepsilon.$$

If x lies in the support of η_2 , that is, $\frac{\delta_s}{3\varepsilon} \leq s(x) \leq \frac{\delta_s}{2\varepsilon}$, then

$$\|\Delta \eta_2\|_{C^\alpha(\bar{B}_1(x))} \leq C \varepsilon^2, \quad \|\phi_2\|_{C^\alpha(\bar{B}_1(x))} \leq \varepsilon^{-2} e^{-\gamma \delta_s / \varepsilon - \varepsilon \gamma s(x)} \|\phi_2\|_{*, \mathcal{E}_0}.$$

Then

$$\|\phi_2 \Delta \eta_2\|_{**, \Sigma} \leq C e^{-\frac{\delta_s}{2\varepsilon}} \|\phi_2\|_{*, \mathcal{E}_0}.$$

Similarly,

$$\|\nabla\phi_2\nabla\eta_2\|_{**,\Sigma} \leq C\varepsilon^{-1}e^{-\frac{\delta_s}{2\varepsilon}}\|\phi_2\|_{*,\mathcal{E}_0}, \quad \text{and} \quad \|\phi_2\partial_z\eta_2\|_{**,\Sigma} \leq C\varepsilon^{-1}e^{-\frac{\delta_s}{2\varepsilon}}\|\phi_2\|_{*,\mathcal{E}_0}.$$

Because the C^1 norm of the coefficients of L' in $s \leq 5\delta_s/\varepsilon$ are bounded by $C(\delta_p + \delta_s + \varepsilon)$, we have

$$\|L'\phi_1\|_{**,\Sigma} \leq C(\delta_p + \delta_s + \varepsilon)\|\phi_1\|_{*,\Sigma}.$$

Also,

$$\|\varepsilon\partial_z\phi_1\|_{**,\Sigma} \leq C\varepsilon\|\phi_1\|_{*,\Sigma}.$$

For the Q term, analogously to Proposition 1.3, we have

$$\|\tilde{\eta}_1 Q(x, \phi, \nabla\phi, D^2\phi)\|_{**,\Sigma} \leq C\|\phi_1\|_{*,\Sigma}^2 + C\varepsilon^{-4}e^{-\frac{\gamma\delta_s}{\varepsilon}}\|\phi_2\|_{*,\mathcal{E}_0}^2.$$

Therefore, using (6.10) and the previous inequalities, we obtain

$$\begin{aligned} \|\bar{\phi}_1\|_{*,\Sigma} &\leq CC_E\varepsilon + C\varepsilon^{-1}e^{-\frac{\delta_s}{2\varepsilon}}\|\phi_2\|_{*,\mathcal{E}_0} + C(\delta_p + \delta_s + \varepsilon)\|\phi_1\|_{*,\Sigma} \\ &\quad + C\|\phi_1\|_{*,\Sigma}^2 + C\varepsilon^{-4}e^{-\frac{\gamma\delta_s}{\varepsilon}}\|\phi_2\|_{*,\mathcal{E}_0}^2, \end{aligned}$$

which gives

$$(6.11) \quad \begin{aligned} \|\bar{\phi}_1\|_{*,\Sigma} &\leq CC_E\varepsilon + C\varepsilon^{-1}e^{-\frac{\delta_s}{2\varepsilon}}M\varepsilon + C(\delta_p + \delta_s + \varepsilon)M\varepsilon + CM^2\varepsilon^2 + C\varepsilon^{-4}e^{-\frac{\gamma\delta_s}{\varepsilon}}M^2\varepsilon. \end{aligned}$$

To estimate $\bar{\phi}_2$, we use Proposition 5.1 to obtain

$$(6.12) \quad \|\bar{\phi}_2\|_{*,\mathcal{E}_0} \leq C\|L''\phi_2 - 2\nabla\phi_1\nabla\eta_1 - \phi_1\Delta\eta_1 - \tilde{\eta}_2((\tilde{E} + Q(\phi, \nabla\phi, D^2\phi)))\|_{**,\mathcal{E}_0}.$$

We first remark that, by Proposition 1.2, we have

$$\|\tilde{\eta}_2\tilde{E}\|_{**,\mathcal{E}_0} \leq C_E\varepsilon.$$

We note that $\Delta\eta_1$ is supported on $\frac{2\delta_s}{\varepsilon} - 1 \leq s \leq \frac{2\delta_s}{\varepsilon}$. Using the smoothness of η_1 , we get

$$\|\phi_1\Delta\eta_1\|_{**,\mathcal{E}_0} + \|\nabla\phi_1\nabla\eta_1\|_{**,\mathcal{E}_0} \leq Ce^{-\frac{\gamma\delta_s}{\varepsilon}}\|\phi_1\|_{*,\Sigma}.$$

Again, as in Proposition 1.3, we have

$$\|\tilde{\eta}_2 Q(x, \phi, \nabla\phi, D^2\phi)\|_{**,\mathcal{E}_0} \leq C\|\phi_1\|_{*,\Sigma}^2 + C\varepsilon^{-4}e^{-\frac{\gamma\delta_s}{\varepsilon}}\|\phi_2\|_{*,\mathcal{E}_0}^2.$$

Combining (6.12) with the previous inequalities, we arrive at

$$(6.13) \quad \|\bar{\phi}_2\|_{*,\Sigma} \leq CC_E\varepsilon + o(\delta_p)M\varepsilon + Ce^{-\frac{\delta_s}{2\varepsilon}}M\varepsilon + CM^2\varepsilon^2 + C\varepsilon^{-4}e^{-\frac{\gamma\delta_s}{\varepsilon}}M^2\varepsilon.$$

The right-hand sides of (6.11) and (6.13) are less than $M\varepsilon$ provided M is fixed large (for example $2CC_E$), then we fix δ_p, δ_s small and work with small $\varepsilon > 0$. A similar estimate holds for $\beta_i, \bar{\tau}_i$ and this shows that F maps \mathcal{B} to itself.

Let us verify that F is a contraction in \mathcal{B} . For this, we first claim that

$$|F(\phi_1, \phi_2, \beta_1, \beta_2, \tau_1, \tau_2) - F(\psi_1, \psi_2, \beta_1, \beta_2, \tau_1, \tau_2)| \leq o(1)(\|\phi_1 - \psi_1\|_{*,\Sigma} + \|\phi_2 - \psi_2\|_{*,\mathcal{E}_0})$$

for $(\phi_1, \phi_2, \beta_1, \beta_2, \tau_1, \tau_2), (\psi_1, \psi_2, \beta_1, \beta_2, \tau_1, \tau_2) \in \mathcal{B}$, where $o(1)$ is small if we choose the parameters $\delta_p, \delta_s > 0$ small and then let ε be small. The estimate relies on the same computations as before for the terms that are linear in ϕ_1, ϕ_2 in the right hand side of equations (6.8) and (6.9). For the nonlinear terms, it is enough

to have the following inequalities, whose proof is similar to Proposition 1.3: For $\phi = \eta_1\phi_1 + \eta_2\phi_2$, $\psi = \eta_1\psi_1 + \eta_2\psi_2$,

$$\begin{aligned} & \|\tilde{\eta}_1 Q(x, \phi, \nabla\phi, D^2\phi) - \tilde{\eta}_1 Q(x, \psi, \nabla\psi, D^2\psi)\|_{**,\Sigma} \\ & \leq C(\|\phi_1\|_{*,\Sigma} + \|\phi_2\|_{*,\varepsilon_0} + \|\psi_1\|_{*,\Sigma} + \|\psi_2\|_{*,\varepsilon_0})(\|\phi_1 - \psi_1\|_{*,\Sigma} + \|\phi_2 - \psi_2\|_{*,\varepsilon_0}) \end{aligned}$$

and there is a similar estimate for $\tilde{\eta}_2 Q$. The Lipschitz dependence with small constant of F on β_i , τ_i is proved using that, in each term in the right-hand side of (6.8) and (6.9), either the dependence on the parameters is Lipschitz with small constant or is quadratic (this is the case of \tilde{E}).

By the contraction mapping principle, for $\varepsilon > 0$ small, F has a unique fixed point in \mathcal{B} . This gives the desired solution.

Acknowledgements. We would like to thank Sigurd Angenent, Nikolaos Kapouleas, Frank Pacard and Juncheng Wei for useful conversations. The first and second authors have been supported by grants Fondecyt 1130360 and 1150066, Fondo Basal CMM and by Núcleo Mienio CAPDE.

REFERENCES

- [1] S.J. Altschuler, L.F. Wu, *Translating surfaces of the non-parametric mean curvature flow with prescribed contact angle*. Calc. Var. Partial Differential Equations 2 (1994), no. 1, 101–111.
- [2] S.B. Angenent, J. J. L. Velázquez, *Degenerate neckpinches in mean curvature flow*. J. Reine Angew. Math. 482 (1997), 15–66.
- [3] T.H. Colding, W. P. Minicozzi, *Minimal surfaces and mean curvature flow*. Surveys in geometric analysis and relativity, 73143, Adv. Lect. Math. (ALM), 20, Int. Press, Somerville, MA, 2011.
- [4] T.H. Colding, W. P. Minicozzi, *Generic mean curvature flow I: generic singularities*. Ann. of Math. (2) 175 (2012), no. 2, 755–833.
- [5] T.H. Colding, W.P. Minicozzi, *Sharp estimates for mean curvature flow of graphs*. J. Reine Angew. Math. 574, (2004), 187–195.
- [6] C.J. Costa, *Imersões mínimas em \mathbf{R}^3 de gênero um e curvatura total finita*, PhD thesis, IMPA, Rio de Janeiro, Brasil (1982).
- [7] C.J. Costa, *Example of a complete minimal immersion in \mathbf{R}^3 of genus one and three embedded ends*. Bol. Soc. Brasil. Mat. 15 (1984), no. 1-2, 47–54.
- [8] J. Clutterbuck, O.C. Schnürer, F. Schulze, *Stability of translating solutions to mean curvature flow*. Calc. Var. Partial Differential Equations 29 (2007), no. 3, 281–293.
- [9] M. del Pino, M. Kowalczyk, J. Wei, *Traveling waves with multiple and nonconvex fronts for a bistable semilinear parabolic equation*. Comm. Pure Appl. Math. 66 (2013), no. 4, 481–547.
- [10] M. Grayson, *The heat equation shrinks embedded plane curves to round points*. J. Differential Geom. 26 (1987), no. 2, 285–314.
- [11] R. Hamilton, *Harnack estimates for the mean curvature flow*, J. Diff. Geom. 41 (1995), 215–226.
- [12] L. Hauswirth, F. Pacard, *Higher genus Riemann minimal surfaces*, Invent. Math. 169 (2007), 569–620.
- [13] D. Hoffman, W.H. Meeks, *Limits of minimal surfaces and Scherk’s fifth surface*. Arch. Rational Mech. Anal. 111 (1990), no. 2, 181–195.
- [14] D. Hoffman, W.H. Meeks III, *Embedded minimal surfaces of finite topology*, Ann. Math. 131 (1990), 1–34.
- [15] G. Huisken, *Asymptotic behavior for singularities of the mean curvature flow*. J. Differential Geom. 31 (1990), no. 1, 285–299.
- [16] G. Huisken and C. Sinestrari, *Mean curvature flow singularities for mean convex surfaces*. Cal. Var. PDE 8(1999), 1–14.
- [17] G. Huisken and C. Sinestrari, *Convexity estimates for mean curvature flow and singularities of mean convex surfaces*, Acta Math. 183 no.1, (1999), 45–70.

- [18] N. Kapouleas, *Complete constant mean curvature surfaces in Euclidean three-space*. Ann. of Math. (2) 131 (1990), no. 2, 239–330.
- [19] N. Kapouleas, *Compact constant mean curvature surfaces in Euclidean three-space*. J. Differential Geom. 33 (1991), no. 3, 683–715.
- [20] N. Kapouleas, *Complete embedded minimal surfaces of finite total curvature*. J. Differential Geom. 47 (1997), no. 1, 95–169.
- [21] T. Ilmanen *Lectures on Mean Curvature Flow and Related Equations* 1998, (Former title: “Some Techniques in Geometric Heat Flow.” ICTP, Trieste 1995).
- [22] T. Ilmanen, *Generalized flow of sets by mean curvature on a manifold*. Indiana Univ. Math. J. 41 (1992), no. 3, 671–705.
- [23] L.P. Jorge, W.H. Meeks III, *The topology of complete minimal surfaces of finite total Gaussian curvature*, Topology 22 (1983), no. 2, 203–221.
- [24] S. Montiel, A. Ros, *Schrödinger operators associated to a holomorphic map*. Global differential geometry and global analysis (Berlin, 1990), 147–174, Lecture Notes in Math., 1481, Springer, Berlin, 1991.
- [25] X.H. Nguyen, *Translating tridents*. Comm. Partial Differential Equations 34 (2009), no. 1-3, 257–280.
- [26] X.H. Nguyen, *Complete embedded self-translating surfaces under mean curvature flow*. J. Geom. Anal. 23 (2013), no. 3, 1379–1426.
- [27] X.H. Nguyen, *Doubly periodic self-translating surfaces for the mean curvature flow*. Geom. Dedicata, to appear.
- [28] R. Osserman, *A survey of minimal surfaces*, Math. Studies 25, Van Nostrand, New York 1969.
- [29] R. Schoen, *Uniqueness, symmetry, and embeddedness of minimal surfaces*, J. Diff. Geom. 18 (1983), 791809.
- [30] W. Sheng, Xu-Jia Wang, *Regularity and singularity in the mean curvature flow*. Trends in partial differential equations, 399–436, Adv. Lect. Math. (ALM), 10, Int. Press, Somerville, MA, 2010.
- [31] M. Traizet, *Construction de surfaces minimales en recollant des surfaces de Scherk*. Ann. Inst. Fourier (Grenoble) 46 (1996), no. 5, 1385–1442.
- [32] M. Traizet, *An embedded minimal surface with no symmetries*, J. Diff. Geom. 60 (2002) 103–153.
- [33] X. Wang, *Convex solutions to the mean curvature flow*. Ann. of Math. (2) 173 (2011), no. 3, 1185–1239.
- [34] B. White, *The size of the singular set in mean curvature flow of mean-convex sets*, J. Amer. Math. Soc. 13 (2000), 665–695.
- [35] B. White, *The nature of singularities in mean curvature flow of mean-convex sets*, J. Amer. Math. Soc. 16 (2003), 121–138.