

An inhomogeneous model of protein dielectric properties: Intrinsic polarizabilities of amino acids

Xueyu Song^{a)}

Department of Chemistry, Iowa State University, Ames, Iowa 50011

(Received 20 September 2001; accepted 8 March 2002)

A simple inhomogeneous model of protein dielectric properties is discussed. A protein in solution is modeled as a collection of polarizable dipoles in a cavity embedded inside a dielectric medium. The intrinsic polarizabilities of 20 amino acids are assumed to be portable to all proteins in nature. A reasonable set of these polarizability values has been obtained by comparing dielectric fluctuations from molecular dynamics simulations with model calculations. The results are consistent within a data set of three small proteins. © 2002 American Institute of Physics. [DOI: 10.1063/1.1474582]

Dielectric properties of proteins play important roles in their structural and functional characteristics.¹⁻³ A simple approach is to approximate the protein interior region by a dielectric continuum with a single dielectric constant, which is embedded in a cavity surrounded by another dielectric continuum representing the solvent. The inhomogeneous nature of the protein interior, important for example in the dielectric response on the active site or embedded chromophore,⁴ is neglected in this approach. Although atomic point charges have been introduced inside the cavity to refine the model,³ an appropriate value of the dielectric constant for a protein is not well defined in general.

Molecular dynamics (MD) simulations allow for a detailed prediction of the protein dielectric properties in solution.^{5,6} Despite considerable progress made recently, the computational cost involved in simulating large proteins in solution is still demanding.

In this report, a physically well-defined middle ground between the dielectric continuum approach and detailed atomistic models such as MD methods is presented. A protein molecule in solution is represented by a set of polarizable dipoles embedded in a dielectric medium of solvent molecules. The interface separating the cavity containing the dipoles from the solvent region can be approximated as a solvent-accessible molecular surface generated by a suitable algorithm from the native structure of the protein. Dielectric properties of the protein are approximated by replacing each residue of the protein by a polarizable dipole. The positions of the dipoles can be consistently assigned based on the native structure of the protein. A set of polarizability values appropriate for each of the residues would completely determine model parameters, allowing for calculations of the protein dielectric response function.

An *intrinsic* set of polarizabilities is introduced and obtained for each of the 20 naturally occurring amino acids. This set is furthermore assumed to be universal, such that once a set of reasonable values has been determined, they can be used for predicting dielectric properties of any large proteins. The intrinsic polarizability of a dipole would be

modified by the interactions with other dipoles and the effect of solvent polarizations, depending on the particular environments produced by the protein of interest. We note that Voges and Karshikoff⁷ have introduced a similar model for the electronic polarizabilities of residues employing the idea of local dielectric constants for proteins. A distinguishing feature of our model is the way by which the nuclear intrinsic polarizabilities are determined. Therefore a combination of our model and the model by Voges and Karshikoff may offer a realistic representation for the dielectric inhomogeneity of proteins.

Our strategy of obtaining a reasonable set of intrinsic polarizabilities is to perform MD simulations of small proteins, and to compare the dipole moment fluctuations generated from the dynamical trajectories with a predicted expression of the model. A critical test of the concept of intrinsic polarizabilities would be the consistency of values determined for one residue with those for different residues of the same amino acid type, both within a single protein and in different proteins. By simulating three small proteins, and analyzing the data in terms of the model, we have found a reasonable set of intrinsic polarizabilities of amino acids, which is consistent within a protein and between different proteins.

Before providing a detailed presentation of our model and calculations we would like to put our work in a proper context. The basic idea of using an atomistic force field to simulate properties of proteins has been well established by extensive studies.^{5,6} In this case a consistent set of atomic parameters are obtained from quantum chemistry calculations or by fitting experimental data, and this set of parameters is assumed to be universal. Indeed many studies have shown the universality of these parameters.⁸ Due to the long-range nature of the electrostatic interaction and the well-defined protein native structure it is reasonable to assume that the dielectric properties of a protein can be described by a collection of normal modes originated from the collective atomic motions around its native structure. Since a protein consists of 20 structurally universal units it seems natural to assume that the polarizabilities of the amino acids in proteins are universal. For the electronic part of the polarizability this

^{a)}Electronic mail: xsong@iastate.edu

universality is evident and it is not far from the electronic polarizability of a free residue. But the nuclear part of the polarizability is not necessarily universal due to the drastic different local environments even for the same amino acid, and it will be distinctly different from the nuclear polarizability obtained from free residue calculation because of the significant constraints in the native structure. Our work presented here shows that it is possible to have a consistent and universal set of nuclear polarizabilities if two major effects due to local environments are removed from the molecular dynamics simulation, which is the method used to extract these polarizabilities in this study. The first one is the local reaction field due to the outside dielectric medium and it is removed by subtracting the exact reaction field contribution calculated from corresponding MD trajectories [cf. Eq. (9)], which had not been done in previous studies.^{6,9} The second local environment effect is the cross interactions among various polarizable dipoles and this effect is removed in Eq. (8). The removal of these local environment effects is the crucial factor to render the model effective and it has not been done before.

The Hamiltonian of a collection of polarizable dipoles inside a dielectric cavity can be written as

$$H = H_p + H_s + H_{ps}. \quad (1)$$

In Eq. (1), H_p is the Hamiltonian for the interacting polarizable dipoles inside the cavity,

$$H_p = \frac{1}{2} \sum_i (\mathbf{m}_i - \bar{\mathbf{m}}_i)^2 / \alpha_i + \frac{1}{2} \sum_{i \neq j} \mathbf{m}_i \cdot \mathbf{T}(\mathbf{r}_i - \mathbf{r}_j) \cdot \mathbf{m}_j, \quad (2)$$

where \mathbf{m}_i is the dipole moment vector of the i th residue centered at the position \mathbf{r}_i with polarizability α_i , and $\mathbf{T}(\mathbf{r}_i - \mathbf{r}_j) = \nabla_i \nabla_j (1/|\mathbf{r}_i - \mathbf{r}_j|)$ is the dipole-dipole interaction tensor, where ∇_i and ∇_j are the gradients with respect to \mathbf{r}_i and \mathbf{r}_j , respectively. In Eq. (2), the polarizability α_i of a particular residue is defined in terms of the dipole moment fluctuations with respect to $\bar{\mathbf{m}}_i$, which is also assumed to be an intrinsic property of the corresponding amino acid. The specific values of these intrinsic dipole moments, however, will not affect predictions of the protein dielectric response, as will be shown in the following.

As has been shown previously,^{2,11,12} the solvent Hamiltonian H_s in Eq. (1) can be written in a quadratic form in terms of electric field of the solvent $\mathbf{E}(\mathbf{r})$:

$$H_s = \frac{1}{2} \int \int_{\text{out}} d\mathbf{r} d\mathbf{r}' \mathbf{E}(\mathbf{r}) \cdot \chi_m^{-1}(\mathbf{r}, \mathbf{r}') \cdot \mathbf{E}(\mathbf{r}'), \quad (3)$$

where “out” denotes that the integration is limited to the region outside the cavity and the modified solvent response function $\chi_m(\mathbf{r}, \mathbf{r}')$ can be obtained from the electric field fluctuations in the presence of the cavity.¹⁰⁻¹²

The interaction Hamiltonian H_{ps} in Eq. (1) is defined as

$$H_{ps} = \sum_i \mathbf{m}_i \cdot \mathbf{E}(\mathbf{r}_i) \quad (4)$$

$$\mathbf{E}(\mathbf{r}_i) = \int_{\text{out}} d\mathbf{r} q(\mathbf{r}) \nabla \frac{1}{|\mathbf{r}_i - \mathbf{r}|},$$

where $q(\mathbf{r})$ is the charge density of the solvent and $\mathbf{E}(\mathbf{r}_i)$ is the electrostatic field at position \mathbf{r}_i due to solvent molecules. The summation over i runs over all of the residues in the protein. Gaussian electric field of the solvent can be integrated out to yield an effective Hamiltonian of the protein,

$$H_{\text{eff}} = H_p + \frac{1}{2} \sum_{ij} \mathbf{m}_i \cdot \mathbf{Q}(\mathbf{r}_i, \mathbf{r}_j) \cdot \mathbf{m}_j. \quad (5)$$

The solvent-mediated interaction tensor between the dipoles $\mathbf{Q}(\mathbf{r}_i, \mathbf{r}_j)$ in Eq. (5) is given by

$$\mathbf{Q}(\mathbf{r}_i, \mathbf{r}_j) = -\beta \left\langle \int \int_{\text{out}} d\mathbf{r} d\mathbf{r}' q(\mathbf{r}) \nabla \frac{1}{|\mathbf{r}_i - \mathbf{r}|} \nabla' \frac{1}{|\mathbf{r}' - \mathbf{r}_j|} q(\mathbf{r}') \right\rangle, \quad (6)$$

where $\langle \rangle$ denotes the ensemble average in the presence of the solute, where $1/\beta = k_B T$ is the Boltzmann constant times temperature. Equation (6) determines the solvent-mediated interaction between dipoles due to the reaction field from solvent polarization. From MD trajectories the ensemble average can be done and the reaction field tensor can be obtained directly from MD simulations. It should be noted that since the MD trajectories are generated in the presence of the solute and the ensemble average performed will be the ensemble average of the solvent molecules in the presence of the solute.

Combining Eqs. (2) and (5), we obtain an implicit relation determining the mean dipole moment $\langle \mathbf{m}_i \rangle$ of the i th residue in terms of intrinsic dipole moment of the corresponding amino acid,

$$\bar{\mathbf{m}}_i = \langle \mathbf{m}_i \rangle + \alpha_i \sum_j [(1 - \delta_{ij}) \mathbf{T}(\mathbf{r}_i, \mathbf{r}_j) + \mathbf{Q}(\mathbf{r}_i, \mathbf{r}_j)] \cdot \langle \mathbf{m}_j \rangle, \quad (7)$$

where δ_{ij} is the Kronecker delta for the residue indices i and j . Mean dipole moment $\langle \mathbf{m}_i \rangle$ represents the dipole moment “modified” from $\bar{\mathbf{m}}_i$ due to the interaction with other dipoles and surrounding solvents. It would correspond to the equilibrium dipole moment of a residue of the protein in solution.

Dipole moment correlation tensor $\mathbf{M}(\mathbf{r}_i, \mathbf{r}_j) = \beta \langle \delta \mathbf{m}_i \delta \mathbf{m}_j \rangle$, associated with the fluctuations $\delta \mathbf{m}_i = \mathbf{m}_i - \langle \mathbf{m}_i \rangle$, is given by due to the quadratic nature of Hamiltonian equation (5),

$$\mathbf{M}^{-1}(\mathbf{r}_i, \mathbf{r}_j) = \delta_{ij} \mathbf{I} / \alpha_i + (1 - \delta_{ij}) \mathbf{T}(\mathbf{r}_i, \mathbf{r}_j) + \mathbf{Q}(\mathbf{r}_i, \mathbf{r}_j). \quad (8)$$

Equation (8) shows that the correlation of the dipole moment fluctuations, and therefore the dielectric response of the protein is independent of the mean dipole moment $\langle \mathbf{m}_i \rangle$ or the intrinsic dipole moment $\bar{\mathbf{m}}_i$.

With the solvent-mediated interaction tensor $\mathbf{Q}(\mathbf{r}_i, \mathbf{r}_j)$ obtained from Eq. (6), Eq. (8) completely determines the dielectric response of a protein within the framework of the model, given the coordinates of the dipoles $\{\mathbf{r}_i\}$ based on the equilibrium solution structure of the protein, and the set of values of the intrinsic polarizabilities $\{\alpha_i\}$. The crucial point of the model lies in the determination of the intrinsic amino acid polarizabilities. If available, experimental data corresponding to the left-hand side of Eq. (8) would provide use-

TABLE I. Simulation and molecular surface configurations for three proteins, bovine pancreatic trypsin inhibitor (BPTI), ribonuclease inhibitor (RNI), and lysozyme (LYZ).

Protein	Box size (Å)	No. of water
BPTI	46.7	2955
RNI	51.2	3871
LYZ	56.6	5014

ful guidance to these intrinsic polarizabilities. Alternatively, MD simulation results can be used instead, provided that the right-hand side of Eq. (8) is also calculated within the model with conditions corresponding to the simulation. We have adopted the latter strategy, with adjustments of the boundary element calculations to fit MD results with Ewald summation.

MD simulations of three small proteins, bovine pancreatic trypsin inhibitor (BPTI), ribonuclease inhibitor, and lysozyme with water molecules have been performed to obtain the dipole moment correlation tensor $\mathbf{M}(\mathbf{r}_i, \mathbf{r}_j)$. Initial coordinates of the proteins were taken as crystal structures from the Protein Data Bank.¹³ Periodic boundary conditions have been used with cubic boxes of different lengths filled with TIP3P¹⁴ water molecules of bulk density 1.00 g/cm³. A CHARMM19 force field was used,¹⁵ with the integration time step 2 fs. Covalent bonds between hydrogen and heavy atoms were fixed by SHAKE algorithm.¹⁶ The particle mesh Ewald method¹⁷ was used for electrostatic interactions. For each system, data were collected over 1.0 ns after initial 0.3 ns equilibration. Temperature was maintained near 300 K by periodically rescaling particle velocities every 1000 steps. Since dipole moments of charged residues are not well defined the partial charges of the charged residues are assigned based upon Lazaridis and Karplus scheme¹⁸ to render the residue neutral. Table I shows the box sizes and the number of water molecules in a unit cell used for the simulations.

To make comparisons with the model calculation of dipole moment correlation tensor based on the geometry of a single cavity inside an infinitely extended dielectric medium, MD simulations of a protein inside a continuous dielectric material of infinite size would have to be performed. For MD simulations with periodic boundary conditions an Ewald summation has to be performed to calculate the reaction field based on the same configuration. The polarization reaction field inside the cavity in this case can be written as

$$\mathbf{Q}(\mathbf{r}_i, \mathbf{r}_j) = -\beta \sum_{\mathbf{n}} \left\langle \int \int_{\text{out}} d\mathbf{r}' q(\mathbf{r}') \times \nabla \frac{1}{|\mathbf{r}_i - \mathbf{r} + \mathbf{n}|} \nabla' \frac{1}{|\mathbf{r}' - \mathbf{r}_j + \mathbf{n}|} q(\mathbf{r}') \right\rangle, \quad (9)$$

where \mathbf{n} spans the integer multiples of the lattice vector corresponding to the periodic box.¹⁹ The formula of the Ewald potential with a spherical shell summation scheme on conducting boundary condition was used.²⁰

With the reaction field tensor $\mathbf{Q}(\mathbf{r}_i, \mathbf{r}_j)$ and the dipole moment correlation tensor $\mathbf{M}(\mathbf{r}_i, \mathbf{r}_j)$, Eq. (8) can be represented as a $3N \times 3N$ matrix equation with the set of 20 unknowns $\{\alpha_i\}$, where N is the number of residues in the protein. As can be seen from Eq. (8), only diagonal elements of the inverse of dipole moment correlation tensor is affected by the values of polarizabilities within the framework of the quadratic Hamiltonian defined by Eq. (2). It is readily verified that to minimize the mean square difference between both sides of the diagonal part of Eq. (8), one should take the average of $1/\alpha_i$ values for each residue of a particular type of amino acid. Amino acid compositions of the three proteins simulated are shown in Table II.

For each of the 20 amino acids, the averages and standard deviations of $1/\alpha_i$ were determined from the diagonal part of Eq. (8) with the solvent-mediated contribution calculated by Eq. (9), within the cumulative database of the three proteins shown in the last row of Table II.

Inverse intrinsic polarizabilities of amino acids determined are plotted in Fig. 1, and the corresponding average polarizabilities are shown in Table III. Amino acids with relatively long, polar side chains have larger polarizabilities than those with short, nonpolar side chains. Polarizabilities range from $\alpha_i \sim 1 \text{ \AA}^3$ to $\alpha_i \sim 40 \text{ \AA}^3$, in contrast to the implicit assumption of the dielectric continuum theory, $\alpha_i = \text{const}$. Also shown in Table III are the electronic polarizabilities given by the model of Voges and Karshikoff.⁷ It is clear that the nuclear and electronic polarizabilities have different origin. For example, the nonpolar side chain of tryptophan gives a small nuclear polarizability (1.8 \AA^3), on the other hand the electronic polarizability (21 \AA^3) is rather large due to the electron rich aromatic ring. In the Voges and Karshikoff model the nuclear polarizability is determined from the Onsager model based on the permanent dipole moment of the amino acid. Their values are substantially larger (Table III) than ours given here. This is not unexpected since the dipole fluctuations of residues in a protein are severely constrained in contrast to the unconstrained assumption in

TABLE II. Amino acid composition of the proteins. N is the total number of residues for each protein.

Proteins	N	A	R	N	D	C	Q	E	G	H	I	L	K	M	F	P	S	T	W	Y	V
BPTI	58	6	6	3	2	6	1	2	6	...	2	2	4	1	4	4	1	3	...	4	1
RNI	89	6	3	3	4	2	6	11	5	1	6	12	6	...	2	2	5	4	3	3	5
LYZ	129	13	10	14	6	8	1	2	13	2	6	9	7	2	2	2	10	7	6	4	5
Total	276	25	19	20	12	16	8	15	24	3	14	23	17	3	8	8	16	14	9	11	11




FIG. 1. The inverse intrinsic polarizabilities of amino acids as a function of amino acids. Our abscissa axis is uniform and the ordinate axis is in logarithmic scale, therefore, the polarizability distribution will be close to exponential since all data fall on a straight line approximately. Points and the total lengths of the error bars represent the means and standard deviations of the inverse polarizabilities calculated from Eq. (8).

Onsager's model. If the molecular volume v of each amino acid is known we can define a dimensionless polarizability $y = 4\pi\alpha/3v$ and Clausius–Mossotti relation or other models yield an effective dielectric constant, which can be viewed as a local dielectric constant used in the Voges and Karshikoff model. In our model, once these intrinsic polarizabilities are known the dielectric properties of a protein can be calculated and there is no need to invoke the local dielectric constant concept.

TABLE III. Intrinsic polarizability(α) of amino acids in unit of \AA^3 obtained by taking the inverse of the averages shown in Fig. 1 and its standard deviation shown in parentheses. N_s is the number of residues sampled in the average for an amino acid type. Electronic polarizability (α_∞) of amino acids in unit of \AA^3 from Voges and Karshikoff's paper (Ref. 7) is shown in the fourth column. The fifth column is the nuclear polarizability based on Onsager's model (Ref. 7) in units of \AA^3 .

Amino acid	α	N_s	α_∞	α_d
Ala	1.1(0.4)	25	7.4	1.3
Arg	5.7(0.3)	19	16.7	2.2
Asn	9.8(1.1)	20	11.3	89.1
Asp	4.0(0.3)	13	9.9	29.8
Cys	1.8(0.4)	16	10.5	15.6
Gln	17.7(1.9)	8	14.3	75.2
Glu	6.2(0.5)	15	11.7	31.5
Gly	2.2(0.5)	23	5.6	1.3
His	12.3(5.0)	3	15.7	22.1
Ile	1.1(0.2)	14	12.9	1.3
Leu	1.2(0.6)	23	12.9	1.3
Lys	8.4(0.6)	17	14.3	9.6
Met	2.8(0.7)	3	14.2	30.1
Phe	1.0(0.1)	8	15.0	1.3
Pro	1.3(0.3)	8	10.7	7.1
Ser	7.3(0.5)	16	8.0	28.4
Thr	8.2(0.5)	14	9.8	28.2
Trp	1.8(0.3)	9	21.5	5.0
Tyr	5.3(0.9)	11	16.0	20.8
Val	1.0(0.2)	11	11.1	1.3

Fluctuations of polarizability values for each amino acid arise from the effect of the particular local environment not accounted for within the framework of the model we have adopted. Although it would be of interest to see how the statistics change as we sample more proteins, we expect that the major contribution to the uncertainties observed in Fig. 1 is due to the effect of heterogeneous local environments, which can be captured only partially with the point-dipole picture of protein residues. However, considering the exponential distribution of polarizabilities within 20 amino acids as seen in Fig. 1, and that even a simple dielectric continuum picture already provides a qualitative guide to the problem, it is reasonable to assume that the dielectric properties of a protein on a length scale comparable to or larger than the size of each residue would be relatively insensitive to the uncertainties of the intrinsic polarizabilities.

We also note that the relatively uniform distribution of the polarizabilities of amino acids in logarithmic scale shown in Fig. 1 suggests the possibility that the natural amino acid set has been evolutionarily selected partially to maximize nature's flexibility in protein design in terms of dielectric properties.

As a first step toward an inhomogeneous model on the dielectric response of proteins, it seems the 20 intrinsic polarizabilities of the amino acids did capture the salient features of protein's inhomogeneity. There are many possible ways by which the model can be extended. The model can easily be generalized to the dynamical case.^{11,12} Evaluation of the frequency-dependent intrinsic polarizabilities would allow for predictions of the time-dependent dielectric response of systems such as chromophores embedded in large proteins. In this report all of the residues are neutral and it will be interesting to see how these intrinsic polarizabilities will hold if charge residues are used in the calculation.

The author is grateful for the contributions and discussions of Hyung-June Woo at the initial stage of this work and for the financial support by a Petroleum Research Fund, administered by American Chemical Society.

¹M. F. Perutz, *Science* **201**, 1187 (1978).

²A. Warshel and S. T. Russell, *Q. Rev. Biol.* **17**, 283 (1984); C. N. Schultz and A. Warshel, *Proteins: Struct., Funct., Genet.* **44**, 400 (2001).

³K. A. Sharp and B. Honig, *Annu. Rev. Biophys. Biophys. Chem.* **19**, 301 (1990); T. Simonson, *Int. J. Quantum Chem.* **73**, 45 (1999).

⁴R. A. Marcus and N. Sutin, *Biochim. Biophys. Acta* **811**, 265 (1985).

⁵C. L. Brooks, M. Karplus, and B. M. Pettit, *Adv. Chem. Phys.* **71**, 1 (1987); T. Simonson and C. L. Brooks, *J. Am. Chem. Soc.* **118**, 8452 (1996). Y. Y. Sham, I. Muegge, and A. Warshel, *Biophys. J.* **74**, 1744 (1998).

⁶H. Nakamura, T. Sakamoto, and A. Wada, *Protein Eng.* **2**, 177 (1988); H. Nakamura, *Q. Rev. Biol.* **29**, 1 (1996), and reference therein.

⁷D. Voges and A. Karshikoff, *J. Chem. Phys.* **108**, 2219 (1998).

⁸B. T. Thole, *Chem. Phys.* **59**, 341 (1981); P. Th. van Duijnen and M. Swart, *J. Phys. Chem. A* **102**, 2399 (1998), and references therein.

⁹T. Simonson, D. Perahia, and G. Bricogne, *J. Mol. Biol.* **218**, 859 (1991).

¹⁰D. Chandler, *Phys. Rev. E* **48**, 2898 (1993).

¹¹X. Song, D. Chandler, and R. A. Marcus, *J. Phys. Chem.* **100**, 11954 (1996).

¹²X. Song and D. Chandler, *J. Chem. Phys.* **108**, 2594 (1998).

¹³Protein Data Bank entry codes of the crystal structure data used for bovine pancreatic trypsin inhibitor, ribonuclease inhibitor, and lysozyme were 1BPI, 1BTA, and 135L, respectively.

- ¹⁴W. Jorgensen, J. Chandrasekar, J. Madura, R. Impey, and M. Klein, *J. Chem. Phys.* **79**, 926 (1983).
- ¹⁵B. Brooks, R. Bruccoleri, B. Olafson, D. States, S. Swaminathan, and M. Karplus, *J. Comput. Chem.* **4**, 187 (1983).
- ¹⁶J. Ryckaert, G. Ciccoti, and H. Berendsen, *J. Comput. Phys.* **23**, 327 (1977).
- ¹⁷T. A. Darden, D. M. York, and L. G. Pedersen, *J. Chem. Phys.* **98**, 10089 (1993).
- ¹⁸T. Lazaridis and M. Karplus, *Proteins: Struct., Funct., Genet.* **35**, 133 (1999).
- ¹⁹Analogous summation of the dipole–dipole interaction $\mathbf{T}(\mathbf{r}, \mathbf{r}')$ would have to be done in Eq. (8). This modification is not necessary, since determination of α_i involves the diagonal parts of Eq. (8) only, and the reaction field on a dipole due to its own replicas vanishes.
- ²⁰M. P. Allen and D. J. Tildesley, *Computer Simulation of Liquids* (Oxford University Press, New York, 1994).